



**HAL**  
open science

# Regulation of effector gene expression as concerted waves in *Leptosphaeria maculans*: a two-players game involving a chromatin remodeler and a specific transcription factor

Colin Clairret, Elise J Gay, Antoine Porquier, Françoise Blaise, Claire-line Marais, Thierry Rouxel, Marie-hélène Balesdent, Jessica L. Soyer, Isabelle Fudal

## ► To cite this version:

Colin Clairret, Elise J Gay, Antoine Porquier, Françoise Blaise, Claire-line Marais, et al.. Regulation of effector gene expression as concerted waves in *Leptosphaeria maculans*: a two-players game involving a chromatin remodeler and a specific transcription factor. International Congress of Plant Pathology, Aug 2023, Lyon, France. hal-04565512

**HAL Id: hal-04565512**

**<https://hal.inrae.fr/hal-04565512v1>**

Submitted on 1 May 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Regulation of effector gene expression as concerted waves in *Leptosphaeria maculans*: a two-players game involving a chromatin remodeler and a specific transcription factor**

**Colin Clairet, Elise J. Gay, Antoine Porquier, Françoise Blaise, Claire-Line Marais, Thierry Rouxel, Marie-Hélène Balesdent, Jessica L. Soyer and Isabelle Fudal**

UR BIOGER, INRAE, Université Paris-Saclay, 92120, Palaiseau, France

*Leptosphaeria maculans* is a phytopathogenic fungus responsible for oilseed rape stem canker that displays a particularly complex lifecycle. Different sets of effector-genes are expressed at each stage of oilseed rape infection. Repeat-rich regions of *L. maculans* genome are enriched in effector-genes specifically expressed during biotrophic stages of infection. These regions show a repressed chromatin state during mycelial growth *in vitro*. We showed the importance of chromatin remodeling in the control of effector-genes expression. As such, the repressive histone modification H3K9me3 deposited by the methyltransferase KMT1, is involved in the regulation of these genes not expressed *in vitro* but highly expressed during infection. However, inactivation of *KMT1* did not de-repress the expression of effector-genes *in vitro* at the same level as observed during infection, suggesting additional actors involved, such as transcription factor(s) (TF). We investigated the involvement of Pf2, a fungal specific Zn<sub>2</sub>Cys<sub>6</sub> TF, in the control of effector-gene expression. Deletion of *LmPf2* lead to a non-pathogenic mutant. Its over-expression was not sufficient to express effector-genes *in vitro*. In contrast, its over-expression in a *Kmt1* mutant background induced the expression of effector-genes *in vitro* to the same level as during plant infection. These results demonstrated for the first time a dual control of effector-gene expression involving a chromatin remodeler and an infection specific TF.