



HAL
open science

Natural genetic variation underlying the negative effect of elevated CO₂ on ionome composition in *Arabidopsis thaliana*

Oceane Cassan, Lea-Lou Pimpare, Mozzanino Timothy, Cecile Fizames, Sebastien Devidal, Fabrice Roux, Alexandru Milcu, Sophie Lebre, Alain Gojon, Antoine Martin

► To cite this version:

Oceane Cassan, Lea-Lou Pimpare, Mozzanino Timothy, Cecile Fizames, Sebastien Devidal, et al.. Natural genetic variation underlying the negative effect of elevated CO₂ on ionome composition in *Arabidopsis thaliana*. 2023. hal-04592341v1

HAL Id: hal-04592341

<https://hal.inrae.fr/hal-04592341v1>

Preprint submitted on 8 Jun 2023 (v1), last revised 29 May 2024 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Natural genetic variation underlying the negative effect of elevated CO₂ on ionome composition in *Arabidopsis thaliana*

Océane Cassan¹, Léa-Lou Pimparé¹, Timothy Mozzanino¹, Cécile Fizames¹, Sébastien Devidal², Fabrice Roux³, Alexandru Milcu^{2,4}, Sophie Lèbre⁵, Alain Gojon¹, Antoine Martin^{1*}

¹ IPSiM, Univ Montpellier, CNRS, INRAE, Institut Agro, 34060, Montpellier, France

² Montpellier European Ecotron, Univ Montpellier, CNRS, Campus Baillarguet, 34980, Montferrier-sur-Lez, France

³ Laboratoire des Interactions Plantes-Microbes-Environnement, Institut National de Recherche pour l'Agriculture, l'Alimentation et l'Environnement, CNRS, Université de Toulouse, Castanet-Tolosan, France

⁴ CEFÉ, Univ Montpellier, CNRS, EPHE, IRD, UPVM, SupAgro, INRAE, 34293, Montpellier, France

⁵ IMAG, Montpellier, France

*Author for correspondence: antoine.martin@cirs.fr

1 **Abstract**

2 The elevation of atmospheric CO₂ leads to a decline in the plant mineral content, which poses
3 a major threat to food security in the coming decades. To date, very few genes have been
4 identified as having a role in the negative effect of elevated CO₂ on plant mineral composition.
5 Yet, several studies have shown a certain degree of diversity in the ionome's response to
6 elevated CO₂, associated with genotypic variation. This suggests the existence of genetic
7 factors controlling the effect of CO₂ on ionome composition. However, no large-scale studies
8 have been carried out to date to explore the genetic diversity of the ionome responses to
9 elevated CO₂. Here, we used six hundred *Arabidopsis thaliana* accessions, representing
10 geographical distributions ranging from worldwide to regional and local environments, to
11 analyze the natural genetic variation underlying the negative effect of elevated CO₂ on the
12 ionome composition in plants. We show that the growth under elevated CO₂ leads to a global
13 and important decrease of the ionome content whatever the geographic distribution of the
14 population. We also observed a high range of genetic diversity in the response of the ionome
15 composition to elevated CO₂, and we identified sub-populations, showing effects on their
16 ionome ranging from the most pronounced to resilience or even to a benefit in response to
17 elevated CO₂. Using genome-wide association mapping on the response of each mineral
18 element to elevated CO₂ or on integrative traits, we identified a large set of QTLs and genes
19 associated with the ionome response to elevated CO₂. Finally, we demonstrate that
20 manipulating the function of one of these genes can mitigate the negative effect of elevated
21 CO₂ on the plant mineral composition. Therefore, this resource will contribute to understand
22 the genetic mechanisms underlying the negative effect of elevated CO₂ on the mineral
23 composition of plants, and to the development of biofortified crops adapted to a high-CO₂
24 world.

25 Introduction

26 The elevation of atmospheric CO₂ concentration leads to a decline in the mineral composition
27 of C3 plants ¹. The negative effect of elevated CO₂ on plant mineral composition has been
28 observed worldwide, and alters the content of nutrients that are essential for human
29 nutrition, such as nitrogen (N) and proteins, iron (Fe) or zinc (Zn) ². The rise in atmospheric
30 CO₂ thus poses a major threat to food security in the coming decades. The reasons why
31 elevated CO₂ leads to the degradation of plant mineral composition are far from being well
32 understood. To date, only a few genes with a potential regulatory effect on this mechanism
33 have been identified ³⁻⁷. These elements nevertheless converge towards the fact that the
34 adaptation of plants to future high CO₂ climate can be achieved through the identification and
35 the characterization of genetic mechanisms. In addition to this, several studies suggest that
36 exploring the natural genetic variability of plants represents a major opportunity to
37 understand the mechanisms by which high CO₂ leads to a decline in plant mineral composition
38 ⁸⁻¹⁰. Indeed, a significant diversity in the response of mineral composition to high CO₂ has been
39 observed in several plant species. For protein and therefore N content, as well as for Fe or Zn
40 content, substantial variations have been observed between small panels of genotypes from
41 different species ⁸⁻¹⁰. This implies the presence of a genetic diversity reservoir, which can
42 facilitate the understanding of the ionome's response to high CO₂ and subsequently provide
43 an opportunity to alleviate this negative impact. However, in order to identify the genetic
44 determinants of this negative response of the ionome to high CO₂, large-scale approaches are
45 necessary, but are still lacking for the moment. The objective of this work was to fill the
46 aforementioned knowledge gap by using a large collection of natural genotypes of the model
47 plant *Arabidopsis thaliana* allowing to explore in depth the natural variation of the ionome
48 response to elevated CO₂, and to generate a resource of phenotypic data that can be used in




Figure 1: Elevated CO₂ negatively impacts the ionome content at the population-scale level in *Arabidopsis thaliana*. A. Representation of the experimental design used in this study. The content of eight mineral elements was assessed for around 600 *Arabidopsis thaliana* accessions coming from the REGMAP (B), LANGUEDOC (C) and TOU-A (D) populations. Each dot represents the value of the content of a mineral element for one accession (yellow: ambient CO₂ (aCO₂, ~420 ppm), blue: elevated CO₂ (eCO₂, 900 ppm). N (% of dry weight), Fe ($\mu\text{g}\cdot\text{g}^{-1}$ dry weight), Zn ($\mu\text{g}\cdot\text{g}^{-1}$ dry weight), Cu ($\mu\text{g}\cdot\text{g}^{-1}$ dry weight), Mg ($\mu\text{g}\cdot\text{g}^{-1}$ dry weight), Mn ($\mu\text{g}\cdot\text{g}^{-1}$ dry weight), Na ($\mu\text{g}\cdot\text{g}^{-1}$ dry weight), C (% of dry weight). Asterisks indicate significant differences (Paired Wilcoxon test; *, P < 0.05; **, P < 0.005; ***, P < 0.0005). ns; not significant.

49 association genetics approaches. To this end, we used several hundreds of accessions from
50 different geographic scales of *A. thaliana*, and analyzed their mineral composition under
51 contrasted conditions of CO₂ concentration. This allowed us to extract the general trends in
52 the ionome response to high CO₂, and to identify a large set of genes associated with the
53 variation in the mineral composition of plants in response to high CO₂. By combining this
54 information with genome expression data under elevated CO₂, we end up by functionally
55 validating one of these genes for its importance in the reduction of Zn content under elevated
56 CO₂, and therefore by demonstrating the relevance of this resource for future improvement
57 of plant nutrient content under elevated CO₂.

58

59 **Results**

60 In order to explore the natural variation and identify its underlying genetic basis associated
61 with the negative effect of elevated CO₂ on plant ionome, we used three populations of *A.*
62 *thaliana* representing different geographic scales (i.e., the worldwide REGMAP population,
63 the LANGUEDOC regional population and the local TOU-A population from east of France) and
64 displaying different levels of genetic diversity (Fig. 1A). These populations were grown under
65 ambient or elevated CO₂, and we measured in each accession the composition of their ionome
66 in rosettes, including C, N, Na, Fe, Mg, Mn, Zn and Cu content.

67 **Elevated CO₂ globally decreases ionome content at the population level, whatever the**
68 **geographic scale.**

69 In the three *A. thaliana* populations, we observed a global and important decrease of the
70 ionome content when plants were grown under elevated CO₂ as compared to ambient CO₂.
71 This was particularly the case for N and Fe, for which the decrease in content was very robust
72 and important in each of the population analyzed (Fig. 1B-D). Zn, Cu and Mg content were also




Figure 2: Elevated CO₂ leads to high phenotypic diversity of ionome response in *Arabidopsis thaliana*.

Distributions of the relative change (%) of the content of 8 mineral elements between elevated CO₂ and ambient CO₂, in each population (A: REGMAP, B: LANGUEDOC, C: TOU-A). Each dot represents the value of the relative change of the content a mineral element for one accession. The name of the element appears in bold if the mean of the element in elevated CO₂ is significantly different from the mean of the element in ambient CO₂ (Paired wilcoxon test, significance threshold of 0.05).

73 negatively affected to a significant extent by the growth under elevated CO₂ in the REGMAP
74 and in the TOU-A populations (Fig. 1B, C), although not significantly in the LANGUEDOC
75 population (Fig. 1D). More variability for the effect of elevated CO₂ was observed on Mn and
76 Na content, which were decreased in the REGMAP population, but not significantly changed
77 in the TOU-A and LANGUEDOC populations, respectively. In parallel, the C content of these
78 populations increased under elevated CO₂, by very significant factors for the REGMAP and the
79 LANGUEDOC populations. Altogether, these observations demonstrate that elevated CO₂ has
80 on average a strong negative impact on the mineral content of natural genotypes of *A.*
81 *thaliana* at the population-scale, whatever their geographic distribution.

82

83 **The ionome of *Arabidopsis thaliana* natural accessions displays a high range of phenotypic**
84 **diversity in response to elevated CO₂.**

85 To explore the effect of elevated CO₂ in each accession, we calculated the relative change in
86 nutrient composition of *A. thaliana* accessions from the three populations in response to
87 elevated CO₂. In agreement with the results previously mentioned, we observed that the
88 median relative change of most nutrient content at the population-level was negatively
89 affected by elevated CO₂ (Fig. 2). But the most striking observation was the genetic diversity
90 of ionome response observed in these populations. Indeed, while most the natural accessions
91 were negatively affected by elevated CO₂ (with a negative relative ratio of their nutrient
92 content between ambient and elevated CO₂), a considerable number of accessions were
93 rather not affected by elevated CO₂, or even positively affected, therefore showing an
94 improved nutrient composition under elevated CO₂. For macronutrients like N, the relative
95 change of concentration between ambient and elevated CO₂ varied from 20% to -50%, and for
96 micronutrients like Cu, Fe or Zn, the relative change of concentration between ambient and




Figure 3: Elevated CO₂ results in a general pattern of ionome variation common to most accessions constituting natural populations of *Arabidopsis thaliana*. Principal Component Analysis (PCA) was performed using the variation of each element in response to elevated CO₂. A. Natural accessions were positioned on the PCA and colored based on population. B. Contribution of each element to the PCA axis.

97 elevated CO₂ varied from 100% to -60% (Fig. 2). In addition, some differences among nutrients
98 were observed between populations. For instance, a smaller dispersion of Fe relative change
99 in the LANGUEDOC population, against a higher distribution of Mn relative change.

100 In order to explore the behavior of the different elements in response to elevated CO₂ and to
101 observe the structure of phenotypic variation, we performed a principal component analysis
102 (PCA) of the relative change in the 8 elements for the accessions from the three populations.
103 The accessions from all populations seem to have globally similar responses to elevated CO₂,
104 as suggested by the overlap of the three populations in the two first principal components
105 (Fig. 3A). The first component of the PCA described a clear antagonistic trend between C
106 content and the change of other mineral elements (Fig. 3B), suggesting that most of the
107 variation between accessions in term of mineral response (almost 40%) could be driven by
108 one or a few mechanisms resulting in an inverse variation between the whole ionome and C
109 change (Fig. 3B). Interestingly, the second component, explaining almost 15% of the variation
110 among accessions in term of mineral response, was mainly driven jointly by change in N and
111 C concentration. Altogether, these results show that there is a marked and large variability
112 among accessions in their mineral concentration in response to elevated CO₂, illustrated by
113 accessions negatively affected by elevated CO₂ and others positively affected by elevated CO₂.

114 In order to explore specific behavior of sub-populations, we clustered the accessions from the
115 REGMAP panel via a k-means approach. This multivariate clustering resulted in the
116 partitioning of accessions in three groups (Fig.4 – Suppl. Table 1). Cluster 1 displayed the most
117 negative pattern of ionome response to elevated CO₂. Inversely, accessions included in Cluster
118 2 displayed a globally positive response, with the highest relative change for almost all mineral
119 elements, except for C content. These accessions did not appear to be clustered
120 geographically with respect to their collection origin in the REGMAP panel (Suppl. Fig. 1),




Figure 4: Variation in the response of the ionome to elevated CO₂ identifies contrasting subpopulations inside the REGMAP panel. K-means clustering was performed in the REGMAP accessions to identify different subpopulations. Each accession is represented by a dot, connected by a line between each element. Cluster 1: 65 accessions. Cluster 2: 25 accessions. Cluster 3: 69 accessions.


121 which is in line with the high genetic diversity of response to elevated CO₂ observed at smaller
122 geographical scales (Fig.2). Finally, Cluster 3 displayed a resilient pattern, with accessions
123 showing a globally attenuated response to elevated CO₂. Interestingly, the large phenotypic
124 diversity of the ionome observed in the three populations in response to high CO₂, as well as
125 the presence of contrasted subpopulations in the REGMPA panel, suggests the presence of
126 genetic determinants associated with this response.

127


128 **Genetic architecture of the ionome response to elevated CO₂, and identification of genetic** 129 **determinants**

130 We run Genome-Wide Association (GWA) mapping to describe the genetic architecture of the
131 ionome response to elevated CO₂, and to fine-map candidate genes underlying the detected
132 quantitative trait loci (QTLs). We focused here on the phenotypic data collected on the
133 REGMAP population, and used the sequencing data available for this population ¹¹. We
134 included in this analysis the level of each mineral under ambient and under elevated CO₂, as
135 well as the relative change between ambient and elevated CO₂ for each element. We also
136 included a trait corresponding for each accession to the coordinate on the first and on the
137 second PCA axes (PCA1 and PCA2) explaining collectively more than 50% of ionic variation
138 (Fig. 3). Therefore, these values correspond to traits driving and summarizing a large part of
139 the ionome variation under elevated CO₂. This resulted as a whole in running GWA mapping
140 on 30 different single-trait GWAS. The overall approach was first validated by observing
141 expected results for traits phenotyped under ambient CO₂. For instance, we observed a very
142 strong peak for the Na content at the locus of the *HKT1* gene (Suppl. Fig. 2A), which is known
143 to be involved in the natural genetic variation of Na content in *Arabidopsis thaliana* ¹², or a
144 strong peak for the N content at the locus of the *NIA1* gene (Suppl. Fig. 2B), encoding for an

A Content under eCO₂ Relative change



B Content under eCO₂



C Relative change




Figure 5: Genetic architecture of the response of the ionome to elevated CO₂ in the REGMAP panel of *Arabidopsis thaliana*. A. Manhattan plots for the content of eight mineral elements under elevated CO₂, or for the relative change of the content of mineral elements between elevated CO₂ and ambient CO₂. For each Manhattan plot, SNPs with the 50 most significant P-value, located above the horizontal line, are colored. Bar plots showing the number of SNPs identified by GWAs for traits under elevated CO₂ (B) or for the relative change of the content of mineral elements between elevated CO₂ and ambient CO₂ (C) that are unique to one element or shared between 2 or 3 traits.

145 isoform of the nitrate reductase required for the first step of nitrate reduction and associated
146 with natural genetic variation of N content in *A. thaliana*¹³.

147 GWA mapping revealed a polygenic architecture for each phenotypic trait, although its
148 complexity largely differs among traits. For instance, very few and neat peaks of association
149 were detected Na and Mn content under elevated CO₂, or of Fe and Cu relative change
150 between ambient and elevated CO₂ (Fig.5A, Suppl. Fig. 3). On the other hand, a more complex
151 genetic architecture with the detection of a large number of QTLs was observed for traits
152 related to N or C content (Fig.5A, Suppl. Fig. 3). For each of the traits that have been analyzed
153 under elevated CO₂ or corresponding to the relative change of their content between ambient
154 and elevated CO₂, we isolated the 50 SNPs with the most significant p-value, hereafter named
155 top SNPs (Fig. 5A, Suppl. Tables 2 and 3). In order to identify the overlap between the genetic
156 architecture of each trait, we looked whether some of the top SNPs were shared among traits.
157 While the large majority of SNPs were specific to one trait, 30 and 21 SNPs were shared
158 between two traits for the content under elevated CO₂ or for the relative change between
159 ambient and elevated CO₂, respectively (Fig. 5B and C, Supplemental Tables 2 and 3). In
160 addition, 8 and 2 SNPs were shared between three traits for the content under elevated CO₂
161 or for the relative change between ambient and elevated CO₂, respectively (Fig. 5B and C,
162 Supplemental Tables 2 and 3). Most of the shared SNPs were associated with micronutrients
163 (Fe, Mn, Zn and Mg content) and with N and/or with the first component of the PCA axis. An
164 interesting QTL located on chromosome 1 was notably associated with 6 traits, displaying
165 SNPs shared between Mn, Zn and N relative change and SNPs shared between Mn, N and PC1
166 content under elevated CO₂ (Fig. 5A, Suppl. Tables 2 and 3). Another QTL located on
167 chromosome 3 encompasses SNPs shared between Fe, Zn and PC1 content under elevated
168 CO₂ (Fig. 5A, Suppl. Table 2).




Figure 6: Identification of genes detected by GWA mapping and differentially regulated by elevated CO₂. Intersection between elevated CO₂-DEG in shoot (A) or root (B) and genes identified by GWA mapping. UpSet plots display the number of elevated CO₂-DEG that are associated to a locus identified for the content or the relative change of one or several mineral elements under elevated CO₂. Illustration of the pattern of elevated CO₂-DEG in shoot (C) or root (D) also identified by GWA mapping.

169 We next identified for each trait a list of the genes located at ± 25 kb from the top 50 SNPs,
170 which corresponds to the rough estimate of the decay of linkage disequilibrium identified in
171 *A. thaliana* at the worldwide scale ¹⁴. This resulted in a list of genes for each element, ranging
172 from 154 to 422 genes depending on the element (Suppl. Tables 2 and 3). Among others,
173 several genes associated with top 50 SNPs were identified as obvious candidates of the effect
174 of elevated CO₂ on plant nutrition and ionome content. This was the case of *ZINC INDUCED*
175 *FACILITATOR 1* (*ZIF1*, *AT5G13740*) and *ZIF-LIKE1* (*AT5G13750*), linked with SNPs identified for
176 Zn content under elevated CO₂, and involved Zn sequestration mechanisms ¹⁵. We also noticed
177 the link between SNPs identified for Zn relative change and *TIP2;2* (*AT4G17340*), known to be
178 involved in Zn root-to-shoot translocation ¹⁶. Concerning N relative change, some of the top
179 50 SNPs were linked to *ASN1* (*AT3G47340*), which is an actor of N status and remobilization
180 ^{17,18}. The *H+/CATION EXCHANGER 1* (*CAX1*) gene (*AT2G38170*), involved in the response to Mn
181 deficiency, was also linked to SNPs associated with Mn content under elevated CO₂ ¹⁹. Some
182 of the top 50 SNPs identified for Fe relative change were linked to *MCO2* (*AT5G21100*) and
183 *MCO3* (*AT5G21105*) genes, which have been recently characterized as actors of the regulation
184 of Fe homeostasis ²⁰. Finally, it is interesting to note that the QTL located on chromosome 3
185 mentioned above displaying significant shared SNPs identified for Fe, Zn and PC1 content
186 under elevated CO₂ was associated among other genes with *ISU2* (*AT3G01020*), coding for one
187 of the Fe-S clusters in *Arabidopsis thaliana*, which are known to be essential for
188 photosynthesis and metabolism ²¹. Altogether, this demonstrated that genes identified
189 through this approach represent a large and valuable reservoir of candidates to study and to
190 counteract the effect of elevated CO₂ on plant nutrition and ionome content.

191 To analyze how these genes identified by GWA mapping are regulated by elevated CO₂, we
192 performed RNA-seq from shoots and roots grown under ambient and elevated CO₂.




Figure 7: Natural variation of the *TIP2;2* gene is associated with improved responses of Zn content to elevated CO₂. Manhattan plot of the relative change of Zn content between elevated CO₂ and ambient CO₂ showing the presence of a peak closed to the *TIP2;2* locus. B. Comparison of haplotypes and their relative change of Zn content between elevated CO₂ and ambient CO₂. Three SNPs located at the *TIP2;2* locus are associated to an improvement of Zn content under elevated CO₂ for accessions that possess them (haplotype 1) compared to the rest of the population (haplotype 0). C. Relative expression of *TIP2;2* in the roots under elevated CO₂ for accessions belonging to haplotype 0 or haplotype 1. Relative expression levels were calculated based on *UBQ10* as internal control. Horizontal black line represented the median of each group of haplotypes. ***P < 0,001, unpaired Mann-Whitney test. D. Shoot Zn content under elevated CO₂ for WT (Columbia) and *tip2;2-1* mutant belonging to haplotype 0 or haplotype 1. Data are presented as the mean (with SD) of 5 and 6 biological repeats for the WT and *tip2;2-1*, respectively. *P < 0.05, unpaired Mann-Whitney test.

193 Differentially expressed genes (DEG) associated to the effect of elevated CO₂ were identified
194 from shoots and roots (Suppl. Table 4). We crossed the list of shoots or roots elevated CO₂-
195 DEG with the list of genes identified by GWA mapping for each element, which resulted in a
196 list of 182 genes identified by GWA mapping and differentially regulated by elevated CO₂ in
197 shoot or in roots (Suppl. Table 5), making them strong candidates to be involved in the
198 response of the mineral composition of plants to elevated CO₂. Most of these genes were
199 deregulated by elevated CO₂ in shoot (Fig. 6A, B). In shoot or in roots, these genes mainly
200 showed an association with C-, Mg- or Zn-related traits (Fig. 6A, B). Several of these genes,
201 identified by GWA mapping and whose expression is deregulated in response to high CO₂,
202 were known for their role in nutrient homeostasis. This was the case for the *ASN1* and *DUR3*
203 genes, encoding an asparagine synthase and a urea transporter involved in N metabolism and
204 remobilization, both associated here with a N-related peak of association, and whose
205 expression is modulated by high CO₂ in leaves (Fig. 6C). We also observed in the leaves an
206 interesting profile for several genes related to C metabolism and photosynthesis. This was the
207 case for the *BGP3* gene, involved in chloroplast development, or for the carbonic anhydrase
208 *CA1*, both showing a decreased expression in response to high CO₂ and both associated with
209 a peak in C-related GWA mapping under elevated CO₂ (Fig. 6C). In roots, the gene most
210 deregulated in response to high CO₂ was *AT1G64710*, encoding a GroES-type alcohol
211 dehydrogenase, which interestingly is also deregulated in leaves (Fig. 6D). We also observed
212 in the roots that the expression of the *TIP2;2* gene, associated here with a peak detected for
213 Zn relative change GWA mapping, was deregulated in response to elevated CO₂ (Fig. 6D).

214 To go further, we selected one of the association peaks identified by GWA mapping, and
215 sought to functionally validate the importance of this QTL in response to elevated CO₂, in order
216 to demonstrate the value of our data set and GWA mapping analyses. To do so, we selected

217 an association peak located on chromosome 4 and associated with Zn relative change (Fig.
218 7A). More precisely, this association peak displayed the SNPs with the most significant p -
219 values and more largely three SNPs that fell into the top 10 SNPs of the trait corresponding to
220 the Zn relative change between ambient and elevated CO₂. The SNPs corresponding to the
221 alternative alleles were associated to an increase of Zn content under elevated CO₂ (Fig. 7B).
222 These SNPs are located very close to the *TIP2;2* (*AT4G17340*) gene, which has been recently
223 characterized as an actor of Zn root-to-shoot translocation¹⁶. We thus selected a set of
224 accessions from haplotype 0 (reduced Zn content under elevated CO₂) or haplotype 1
225 (increased Zn content under elevated CO₂), and analyze *TIP2;2* expression in the roots under
226 elevated CO₂. This analysis revealed a haplotype-specific difference in *TIP2;2* expression under
227 elevated CO₂, with accessions from haplotype 1 showing a reduced *TIP2;2* expression in the
228 roots compared to those from haplotype 0 (Fig. 7C), correlated with a higher Zn content in the
229 shoot (Fig. 7B). To validate the effect of *TIP2;2* expression of Zn content under elevated CO₂,
230 we used the *tip2;2-1* knock-out mutant and compared its Zn content under elevated CO₂ to
231 this of the WT. We observed that the *tip2;2-1* mutant line displayed a significant higher Zn
232 content in the shoot under elevated CO₂ compared to the WT, confirming that *TIP2;2*
233 expression determines Zn content under elevated CO₂ (Fig. 7D). Altogether, these results
234 demonstrated that these data sets generated in this study and the associated analyses are a
235 valuable resource to identify genes able to counteract the general negative effect of elevated
236 CO₂ on the mineral composition of plants.

237

238 Discussion

239 **The natural variation of ionome response to elevated CO₂ in *Arabidopsis thaliana* displays a**
240 **high degree of genetic variation**

241 In the present work, we analyzed the diversity of ionome response to elevated CO₂ present in
242 the natural variation of *Arabidopsis thaliana*. In agreement with several other phenotypic
243 traits related to phenology and disease resistance {Brachi, 2013 #1232;Huard-Chauveau, 2013
244 #1247;Roux, 2022 #1248}, we observed a wide range of responses at complementary
245 geographical scales, from accessions with a ionome strongly negatively affected by high CO₂
246 to accessions with a ionome benefiting from high CO₂. This confirms for the first time on large
247 and complementary sets of natural genotypes what has been observed by meta-analysis on
248 isolated groups of plants worldwide ^{2,9}. The global analysis of the distribution of each mineral
249 element studied suggests firstly a trend where the whole ionome would evolve in a unified
250 manner in response to high CO₂, and in an opposite manner to C. This is in line with a number
251 of studies that have proposed that the accumulation of carbohydrates due to the stimulation
252 of photosynthesis by high CO₂ would be the cause of the decrease in plant mineral
253 composition ²²⁻²⁵. However, the reading of the genetic architecture performed here by a
254 genome-wide association genetics approach suggests that the majority of the genetic
255 mechanisms underlying the negative effect of elevated CO₂ on the ionome are specific to each
256 mineral element. Some specific cases, such as the QTL detected on chromosome 1 and
257 associated with the natural genetic variation of 6 traits among the 20 considered, will certainly
258 deserve a more in-depth analysis.

259 By clustering globally distributed accessions according to their ionome sensitivity to high CO₂,
260 we were able to observe that the geographic origin of the accessions likely did not determine
261 their response to CO₂. This suggests that inherent genetic factors, more than those due to
262 local adaption, direct the response of plants to elevated CO₂. This seems consistent since the
263 CO₂ elevation applied here to natural *Arabidopsis thaliana* variants does not correspond to
264 any environment experienced by plants yet, at least for several tens of millions of years ^{26,27}.

265 In this context of brutal and highly impactful environmental change, the presence of cryptic
266 genetic variation often explains the appearance of relatively rapid adaptive mechanisms^{28,29}.
267 Although not formally tested here, it would be interesting to examine whether the variation
268 in the ionome in response to elevated CO₂ shows evidence of cryptic variation. In any case,
269 the presence of high phenotypic diversity in these natural populations of *A. thaliana*
270 demonstrates very clearly the possibility of taking advantage of this genetic variation to
271 understand and alleviate the negative response of plant mineral composition to high CO₂.

272 **GWA mapping of ionome variation under elevated CO₂ identified a large number of genes**
273 **to understand and mitigate the negative effect of high CO₂ on plant mineral composition**

274 In order to understand the genetic mechanisms underlying the effect of high CO₂ on plant
275 mineral composition, and to enable future breeding approaches, we adopted an association
276 genetics approach. This led to the identification of a large number of candidate genes
277 associated to the variation of nutrients under elevated CO₂. Several genes in this list can easily
278 attract attention. In particular, we can note the identification of *ASN1* and *DUR3* genes in two
279 of the loci associated with N content variation under elevated CO₂. *ASN1*, and to a lesser
280 extent *DUR3*, play an important role in the remobilization and the reallocation of N within the
281 plant, and their manipulation can lead to variation in N use efficiency^{17,18,30}. This is interesting
282 because for the moment, root N uptake and N assimilation seemed to be the key targets of
283 the negative effect of high CO₂ on plant N content^{4,31}, but these results suggest that
284 remobilization of N may also be involved. We also identified the *CA1* gene, coding for a
285 carbonic anhydrase, in the vicinity of a QTL associated with C variation under high CO₂. *CA1* is
286 involved in the regulation of stomatal opening by elevated CO₂³², and the β carbonic
287 anhydrase family of which *CA1* belongs is involved in the regulation of photosynthetic
288 efficiency, although *CA1* shows no significant effect under standard conditions³³. It would be

289 therefore interesting to assess the role of *CA1* natural genetic variation under elevated CO₂. If
290 *CA1* regulates the C variation of the ionome under elevated CO₂, this could, according to our
291 observations, significantly influence the global mineral composition of plants. Interestingly,
292 the genes identified by GWA mapping in the ionome response to high CO₂, including those
293 mentioned above, showed substantial variation at the gene expression level. We ended this
294 study with the functional validation of an association peak identified by GWA mapping for the
295 relative change of Zn content between ambient and elevated CO₂. Zn is an essential element
296 for a large number of metabolic processes in humans, and Zn deficiency, found in up to one
297 third of the world's population, leads to severe health problems. We demonstrated that *TIP2;2*
298 gene expression varied in response to CO₂ in a haplotype-specific manner. Consistent with
299 these results, we show that manipulating *TIP2;2* expression with a knock-out mutant can
300 modulate the Zn loss observed under high CO₂. A recent study demonstrated that *TIP2;2* was
301 responsible for Zn retention in the roots {Wang, 2022 #1177}. It therefore seems consistent
302 that natural accessions with the lowest expression levels of this gene are those with the
303 highest Zn content in aerial parts, due to low retention in their roots. This example illustrates
304 the potential of the resource we have generated here towards the development of biofortified
305 plants. The development of biofortified plants represents a considerable challenge in view of
306 the current problems of malnutrition on a global scale, and this challenge becomes even more
307 important in a context of rising atmospheric CO₂ ³⁴. This reservoir of data and genes will
308 certainly contribute to the understanding of the mechanisms underlying the general negative
309 effect of CO₂ on mineral composition, and to the development of crop plants adapted to
310 forthcoming high-CO₂ climate.

311

312 **Methods**

313 *Data and code availability*

314 Data and R notebooks containing the analyses performed in this article can be found at
315 <https://src.koda.cnrs.fr/groups/ipsim/sirene-team>. RNA-seq data generated for this study are
316 available at <https://www.ebi.ac.uk/biostudies/arrayexpress/studies> using the accession no
317 xxx.

318 *Plant Material*

319 A subset of the REGMAP panel, the LANGUEDOC panel and the TOU-A panel were used in this
320 study. These populations were previously described here {Brachi, 2013 #1232;Frachon, 2017
321 #1249;Horton, 2012 #1231}. These populations were grown on Jiffy-7 peat pellets (Jiffy
322 Products International, NL) under ambient (~420 ppm) or elevated (900 ppm) CO₂ in the
323 growth chambers of the Microcosms experimental platform at the Montpellier European
324 Ecotron CNRS. Growth conditions were 6-h/22-h light (22°C) / dark (20°) photoperiod, with
325 200 μmol m⁻² s⁻¹ light intensity and 65% of hygrometry. Plants were watered twice a week
326 with a growth solution containing KH₂PO₄ 1 mM, MgSO₄ 1 mM, K₂SO₄ 250 μM, CaCl₂ 250 μM,
327 Na-Fe-EDTA 100 μM, KNO₃ 10 mM, KCl 50 μM, H₃BO₃ 30 μM, MnSO₄ 5 μM, ZnSO₄ 1 μM, CuSO₄
328 1 μM, (NH₄)₆Mo₇O₂₄ 0,1 μM, as described by³⁵. The entire rosettes were collected three weeks
329 after sowing. The *tip2;2-1* mutant line corresponds to the *SALK_152463* allele¹⁶.

330 *Ionome analysis*

331 From 3 to 5 replicates per accession were used for each ionome analysis. Total C and N content
332 was obtained from dried shoot tissue using an Elementar Pyrocube analyzer. Cu, Fe, Mg, Mn,
333 Na and Zn content was obtained from dry shoot tissue mixed with 750 μl of nitric acid (65%
334 [v/v]) and 250 μl of hydrogen peroxide (30% [v/v]). After one night at room temperature,
335 samples were mineralized at 85°C during 24 hours. Once mineralized, 4 ml of milliQ water was

336 added to each sample. Mineral contents present in the samples were then measured by
337 microwave plasma atomic emission spectroscopy (MP-AES, Agilent Technologies).

338 *Removal of outlier observations*

339 Prior to GWAS and multivariate analyses such as PCA or clustering, mineral composition
340 measures were pre-processed to remove technical outliers. For a given element and CO₂
341 condition, the values positioned more than 5 median absolute deviations away from the
342 median were removed from the dataset.

343 *PCA and Clustering*

344 Principal Component Analysis was performed using the R *ade4* package after the prior scaling
345 of the variables to a z-score. Clustering of the REGMAP panel based on the relative changes of
346 the mineral composition of each accession has been done using a k-means clustering with the
347 R *kmeans* function. For this step, the variables were also scaled to a z-score. The number of
348 clusters in the k-means algorithm was chosen by the elbow method on the criteria of cluster
349 homogeneity (within-sum of squares).

350 *GWAs*

351 Genome-Wide Association mapping was performed using the R *statgenGWAs* package.
352 Genotype data was prepared using the *codeMarkers* function, removing duplicated SNPs and
353 filtering for a minimum allele relative frequency of 0.04. Associations were performed by the
354 *runSingleTraitGwas* function, that implements the EMMA algorithm. Population structure was
355 modeled via a kinship matrix built from the Astle method. Manhattan plots were drawn using
356 the *manPlotFast* function of the *ramwas* R package.

357 *RNA-seq experiments*

358 Plants were grown in hydroponics to have access to the roots in addition to the shoot, as
359 previously described in ⁴. Shoot or root from 5 plants were pooled into one biological replicate,

360 flash frozen in liquid nitrogen, and stored at -80°C. RNA of three biological replicates were
361 extracted from shoot or root tissues using Direct-zol RNA Miniprep (Zymo Research, CA, USA),
362 according to the manufacturer recommendations. RNA-sequencing libraries were done from
363 shoot or root total RNA using standard RNA-Seq protocol method (Poly-A selection for mRNA
364 species) by the Novogene company. RNA-sequencing was performed using Illumina
365 technology on a NovaSeq6000 system providing PE150 reads. The quality control and adapter
366 trimming of raw paired-end fastq files was done with *fastp* and its default parameters.
367 Mapping to the TAIR10 reference genome was performed with STAR, and using the following
368 options:

369 `--outSAMtype BAM SortedByCoordinate`

370 `--outFilterMismatchNmax 1`

371 `--outFilterMismatchNoverLmax 0.15`

372 `--alignIntronMin 30`

373 `--alignIntronMax 5000`

374 Quantification of the bam files against the TAIR10 GFF3 annotation file was done using *htseq-*
375 *count* with options:

376 `-f bam --type gene -r pos`

377 `--idattr=Name --stranded=no`

378 Normalization and differential expression were performed using *DIANE* R package ³⁶, with no
379 fold change constraint, and an adjusted p-value threshold (FDR) of 0.05. Lowly expressed
380 genes with an average value across conditions under 25 reads were excluded from the
381 analysis.

382 *Quantitative real-time PCR*

383 Plants were grown in hydroponics to have access to the roots, as previously described in ⁴.
384 Root tissue from 5 plants were pooled into one biological replicate, flash frozen in liquid
385 nitrogen, and stored at -80°C. RNA were extracted from shoot or root tissues using TRIZOL
386 (Invitrogen, USA), according to the manufacturer recommendations, and DNase treated using
387 RQ1 (Promega, USA). Reverse transcription was achieved from 1 µg of total RNA with M-MLV
388 reverse transcriptase (RNase H minus, Point Mutant, Promega, USA) using an anchored
389 oligo(dT)20 primer. Accumulation of transcripts was measured by qRT-PCR (LightCycler 480,
390 Roche Diagnostics, USA) using the SYBR Premix Ex Taq™ (TaKaRa, Japan). Gene expression
391 was normalized using *UBQ10* and *ACT2* as internal standards. Results are presented as the
392 expression relative to *UBQ10*. Sequences of primers used in RT-qPCR for gene expression
393 analysis are listed in Supplemental Table 6.

394

395 **Acknowledgments**

396 This work was supported by the I-Site Montpellier Université d'Excellence (MUSE; project
397 ECO2THREATS), the CNRS through the Mission for Transversal and Interdisciplinary Initiatives
398 (MITI) 80 PRIME program, and the Plant Biology and Breeding department of INRAE. O.C. was
399 recipient of a PhD fellowship from the CNRS. T.M. was recipient of a PhD fellowship from
400 INRAE and Région Occitanie. We thank Jiping Liu from Cornell University for the gift of *tip2;2-*
401 *1* mutant line. This study benefited from the CNRS human and technical resources allocated
402 to the Ecotron Research Infrastructure and from the state allocation "Investissement
403 d'Avenir" AnaEE France ANR-11-INBS-0001. A CC-BY public copyright license has been applied
404 by the authors to the present document.

405

406 **References**

- 407 1 Gojon, A., Cassan, O., Bach, L., Lejay, L. & Martin, A. The decline of plant mineral
408 nutrition under rising CO₂: physiological and molecular aspects of a bad deal. *Trends*
409 *in Plant Science* **28**, 185-198 (2023). <https://doi.org/10.1016/j.tplants.2022.09.002>
- 410 2 Loladze, I. Hidden shift of the ionome of plants exposed to elevated CO₂ depletes
411 minerals at the base of human nutrition. *Elife* **3**, e02245 (2014).
412 <https://doi.org/10.7554/eLife.02245>
- 413 3 Bouain, N. *et al.* Plant growth stimulation by high CO₂ depends on phosphorus
414 homeostasis in chloroplasts. *Curr Biol* (2022).
415 <https://doi.org/10.1016/j.cub.2022.08.032>
- 416 4 Cassan, O. *et al.* A gene regulatory network in Arabidopsis roots reveals features and
417 regulators of the plant response to elevated CO₂. *New Phytol* (2023).
418 <https://doi.org/10.1111/nph.18788>
- 419 5 Sun, P. *et al.* Countering elevated CO₂ induced Fe and Zn reduction in Arabidopsis
420 seeds. *New Phytol* **235**, 1796-1806 (2022). <https://doi.org/10.1111/nph.18290>
- 421 6 Umnajkitikorn, K., Sade, N., Rubio Wilhelmi, M. D. M., Gilbert, M. E. & Blumwald, E.
422 Silencing of OsCV (chloroplast vesiculation) maintained photorespiration and N
423 assimilation in rice plants grown under elevated CO₂. *Plant Cell Environ* **43**, 920-933
424 (2020). <https://doi.org/10.1111/pce.13723>
- 425 7 Yang, A., Li, Q., Chen, L. & Zhang, W.-H. A rice small GTPase, Rab6a, is involved in the
426 regulation of grain yield and iron nutrition in response to CO₂ enrichment. *Journal of*
427 *Experimental Botany* **71**, 5680-5688 (2020). <https://doi.org/10.1093/jxb/eraa279>

- 428 8 Marcos-Barbero, E. L., Pérez, P., Martínez-Carrasco, R., Arellano, J. B. & Morcuende, R.
429 Screening for Higher Grain Yield and Biomass among Sixty Bread Wheat Genotypes
430 Grown under Elevated CO₂ and High-Temperature Conditions. *Plants* **10**, 1596 (2021).
- 431 9 Myers, S. S. *et al.* Increasing CO₂ threatens human nutrition. *Nature* **510**, 139-142
432 (2014). <https://doi.org/10.1038/nature13179>
- 433 10 Zhu, C. *et al.* Carbon dioxide (CO₂) levels this century will alter the protein,
434 micronutrients, and vitamin content of rice grains with potential health consequences
435 for the poorest rice-dependent countries. *Sci Adv* **4**, eaaq1012 (2018).
436 <https://doi.org/10.1126/sciadv.aaq1012>
- 437 11 Arouisse, B., Korte, A., van Eeuwijk, F. & Kruijer, W. Imputation of 3 million SNPs in the
438 Arabidopsis regional mapping population. *The Plant Journal* **102**, 872-882 (2020).
439 [https://doi.org:https://doi.org/10.1111/tpj.14659](https://doi.org/https://doi.org/10.1111/tpj.14659)
- 440 12 Baxter, I. *et al.* A Coastal Cline in Sodium Accumulation in Arabidopsis thaliana Is Driven
441 by Natural Variation of the Sodium Transporter AtHKT1;1. *PLOS Genetics* **6**, e1001193
442 (2010). <https://doi.org/10.1371/journal.pgen.1001193>
- 443 13 North, K. A., Ehling, B., Koprivova, A., Rennenberg, H. & Kopriva, S. Natural variation
444 in Arabidopsis adaptation to growth at low nitrogen conditions. *Plant Physiology and*
445 *Biochemistry* **47**, 912-918 (2009).
446 [https://doi.org:https://doi.org/10.1016/j.plaphy.2009.06.009](https://doi.org/https://doi.org/10.1016/j.plaphy.2009.06.009)
- 447 14 Kim, S. *et al.* Recombination and linkage disequilibrium in Arabidopsis thaliana. *Nature*
448 *Genetics* **39**, 1151-1155 (2007). <https://doi.org/10.1038/ng2115>
- 449 15 Katz, E. *et al.* Genetic variation underlying differential ammonium and nitrate
450 responses in Arabidopsis thaliana. *The Plant Cell* **34**, 4696-4713 (2022).
451 <https://doi.org/10.1093/plcell/koac279>

- 452 16 Wang, Y. *et al.* AtTIP2;2 facilitates resistance to zinc toxicity via promoting zinc
453 immobilization in the root and limiting root-to-shoot zinc translocation in *Arabidopsis*
454 *thaliana*. *Ecotoxicology and Environmental Safety* **233**, 113333 (2022).
455 [https://doi.org:https://doi.org/10.1016/j.ecoenv.2022.113333](https://doi.org/10.1016/j.ecoenv.2022.113333)
- 456 17 Gaufichon, L. *et al.* ASN1-encoded asparagine synthetase in floral organs contributes
457 to nitrogen filling in *Arabidopsis* seeds. *The Plant Journal* **91**, 371-393 (2017).
458 [https://doi.org:https://doi.org/10.1111/tpj.13567](https://doi.org/10.1111/tpj.13567)
- 459 18 Lam, H.-M. *et al.* Overexpression of the ASN1 Gene Enhances Nitrogen Status in Seeds
460 of *Arabidopsis*. *Plant Physiology* **132**, 926-935 (2003).
461 [https://doi.org:10.1104/pp.103.020123](https://doi.org/10.1104/pp.103.020123)
- 462 19 Li, Y. *et al.* Combined analyses of translome and transcriptome in *Arabidopsis* reveal
463 new players responding to magnesium deficiency. *Journal of integrative plant biology*
464 **63**, 2075-2092 (2021). [https://doi.org:https://doi.org/10.1111/jipb.13169](https://doi.org/10.1111/jipb.13169)
- 465 20 Brun, A. *et al.* MCO1 and MCO3, two putative ascorbate oxidases with ferroxidase
466 activity, new candidates for the regulation of apoplastic iron excess in *Arabidopsis*.
467 *Plant Direct* **6**, e463 (2022). [https://doi.org:10.1002/pld3.463](https://doi.org/10.1002/pld3.463)
- 468 21 Balk, J. & Schaedler, T. A. Iron cofactor assembly in plants. *Annu Rev Plant Biol* **65**, 125-
469 153 (2014). [https://doi.org:10.1146/annurev-arplant-050213-035759](https://doi.org/10.1146/annurev-arplant-050213-035759)
- 470 22 Ainsworth, E. A. & Long, S. P. What have we learned from 15 years of free-air CO₂
471 enrichment (FACE)? A meta-analytic review of the responses of photosynthesis,
472 canopy properties and plant production to rising CO₂. *New Phytol* **165**, 351-371 (2005).
473 [https://doi.org:10.1111/j.1469-8137.2004.01224.x](https://doi.org/10.1111/j.1469-8137.2004.01224.x)

- 474 23 Dusenge, M. E., Duarte, A. G. & Way, D. A. Plant carbon metabolism and climate
475 change: elevated CO₂ and temperature impacts on photosynthesis, photorespiration
476 and respiration. *New Phytol* **221**, 32-49 (2019). <https://doi.org/10.1111/nph.15283>
- 477 24 Tausz-Posch, S., Tausz, M. & Bourgault, M. Elevated [CO₂] effects on crops: Advances
478 in understanding acclimation, nitrogen dynamics and interactions with drought and
479 other organisms. *Plant Biol (Stuttg)* **22 Suppl 1**, 38-51 (2020).
480 <https://doi.org/10.1111/plb.12994>
- 481 25 Thompson, M., Gamage, D., Hirotsu, N., Martin, A. & Seneweera, S. Effects of Elevated
482 Carbon Dioxide on Photosynthesis and Carbon Partitioning: A Perspective on Root
483 Sugar Sensing and Hormonal Crosstalk. *Front Physiol* **8**, 578 (2017).
484 <https://doi.org/10.3389/fphys.2017.00578>
- 485 26 Luthi, D. *et al.* High-resolution carbon dioxide concentration record 650,000-800,000
486 years before present. *Nature* **453**, 379-382 (2008).
487 <https://doi.org/10.1038/nature06949>
- 488 27 Pearson, P. N. & Palmer, M. R. Atmospheric carbon dioxide concentrations over the
489 past 60 million years. *Nature* **406**, 695-699 (2000). <https://doi.org/10.1038/35021000>
- 490 28 Cortés, A. J. & López-Hernández, F. Harnessing Crop Wild Diversity for Climate Change
491 Adaptation. *Genes* **12**, 783 (2021).
- 492 29 Pauls, S. U., Nowak, C., Balint, M. & Pfenninger, M. The impact of global climate change
493 on genetic diversity within populations and species. *Mol Ecol* **22**, 925-946 (2013).
494 <https://doi.org/10.1111/mec.12152>
- 495 30 Bohner, A., Kojima, S., Hajirezaei, M., Melzer, M. & von Wirén, N. Urea retranslocation
496 from senescing Arabidopsis leaves is promoted by DUR3-mediated urea retrieval from

- 497 leaf apoplast. *The Plant Journal* **81**, 377-387 (2015).
498 <https://doi.org/10.1111/tpj.12740>
- 499 31 Bloom, A. J., Burger, M., Rubio Asensio, J. S. & Cousins, A. B. Carbon dioxide enrichment
500 inhibits nitrate assimilation in wheat and Arabidopsis. *Science* **328**, 899-903 (2010).
501 <https://doi.org/10.1126/science.1186440>
- 502 32 Hu, H. *et al.* Distinct Cellular Locations of Carbonic Anhydrases Mediate Carbon Dioxide
503 Control of Stomatal Movements *Plant Physiology* **169**, 1168-1178 (2015).
504 <https://doi.org/10.1104/pp.15.00646>
- 505 33 Sharma, N. *et al.* Arabidopsis stromal carbonic anhydrases exhibit non-overlapping
506 roles in photosynthetic efficiency and development. *Plant J* **n/a** (2023).
507 <https://doi.org/10.1111/tpj.16231>
- 508 34 Shahzad, Z. & Rouached, H. Protecting plant nutrition from the effects of climate
509 change. *Curr Biol* **32**, R725-R727 (2022). <https://doi.org/10.1016/j.cub.2022.05.056>
- 510 35 Gansel, X., Munos, S., Tillard, P. & Gojon, A. Differential regulation of the NO₃⁻ and
511 NH₄⁺ transporter genes AtNrt2.1 and AtAmt1.1 in Arabidopsis: relation with long-
512 distance and local controls by N status of the plant. *Plant J* **26**, 143-155 (2001).
- 513 36 Cassan, O., Lebre, S. & Martin, A. Inferring and analyzing gene regulatory networks
514 from multi-factorial expression data: a complete and interactive suite. *BMC Genomics*
515 **22**, 387 (2021). <https://doi.org/10.1186/s12864-021-07659-2>
516

Figure legends :

Figure 1: Elevated CO₂ negatively impacts the ionome content at the population-scale level in *Arabidopsis thaliana*. A. Representation of the experimental design used in this study. The content of eight mineral elements was assessed for around 600 *Arabidopsis thaliana* accessions coming from the REGMAP (B), LANGUEDOC (C) and TOU-A (D) populations. Each dot represents the value of the content of a mineral element for one accession (yellow: ambient CO₂ (aCO₂, ~420 ppm), blue: elevated CO₂ (eCO₂, 900 ppm). N (% of dry weight), Fe (µg.g⁻¹ dry weight), Zn (µg.g⁻¹ dry weight), Cu (µg.g⁻¹ dry weight), Mg (µg.g⁻¹ dry weight), Mn (µg.g⁻¹ dry weight), Na (µg.g⁻¹ dry weight), C (% of dry weight). Asterisks indicate significant differences (Paired Wilcoxon test; *, P < 0.05; **, P < 0.005; ***, P < 0.0005). ns; not significant.

Figure 2: Elevated CO₂ leads to high phenotypic diversity of ionome response in *Arabidopsis thaliana*. Distributions of the relative change (%) of the content of 8 mineral elements between elevated CO₂ and ambient CO₂, in each population (A: REGMAP, B: LANGUEDOC, C: TOU-A). Each dot represents the value of the relative change of the content a mineral element for one accession. The name of the element appears in bold if the mean of the element in elevated CO₂ is significantly different from the mean of the element in ambient CO₂ (Paired wilcoxon test, significance threshold of 0.05).

Figure 3: Elevated CO₂ results in a general pattern of ionome variation common to most accessions constituting natural populations of *Arabidopsis thaliana*. Principal Component Analysis (PCA) was performed using the variation of each element in response to elevated CO₂.

A. Natural accessions were positioned on the PCA and colored based on population. B. Contribution of each element to the PCA axis.

Figure 4: Variation in the response of the ionome to elevated CO₂ identifies contrasting subpopulations inside the REGMAP panel. K-means clustering was performed in the REGMAP accessions to identify different subpopulations. Each accession is represented by a dot, connected by a line between each element. Cluster 1: 65 accessions. Cluster 2: 25 accessions. Cluster 3: 69 accessions.

Figure 5: Genetic architecture of the response of the ionome to elevated CO₂ in the REGMAP panel of *Arabidopsis thaliana*. A. Manhattan plots for the content of eight mineral elements under elevated CO₂, or for the relative change of the content of mineral elements between elevated CO₂ and ambient CO₂. For each Manhattan plot, SNPs with the 50 most significant P-value, located above the horizontal line, are colored. Bar plots showing the number of SNPs identified by GWAs for traits under elevated CO₂ (B) or for the relative change of the content of mineral elements between elevated CO₂ and ambient CO₂ (C) that are unique to one element or shared between 2 or 3 traits.

Figure 6: Identification of genes detected by GWA mapping and differentially regulated by elevated CO₂. Intersection between elevated CO₂-DEG in shoot (A) or root (B) and genes identified by GWA mapping. UpSet plots display the number of elevated CO₂-DEG that are associated to a locus identified for the content or the relative change of one or several mineral elements under elevated CO₂. Illustration of the pattern of elevated CO₂-DEG in shoot (C) or root (D) also identified by GWA mapping.

Figure 7: Natural variation of the *TIP2;2* gene is associated with improved responses of Zn content to elevated CO₂. Manhattan plot of the relative change of Zn content between elevated CO₂ and ambient CO₂ showing the presence of a peak closed to the *TIP2;2* locus. B. Comparison of haplotypes and their relative change of Zn content between elevated CO₂ and ambient CO₂. Three SNPs located at the *TIP2;2* locus are associated to an improvement of Zn content under elevated CO₂ for accessions that possess them (haplotype 1) compared to the rest of the population (haplotype 0). C. Relative expression of *TIP2;2* in the roots under elevated CO₂ for accessions belonging to haplotype 0 or haplotype 1. Relative expression levels were calculated based on *UBQ10* as internal control. Horizontal black line represented the median of each group of haplotypes. ***P < 0,001, unpaired Mann-Whitney test. D. Shoot Zn content under elevated CO₂ for WT (Columbia) and *tip2;2-1* mutant belonging to haplotype 0 or haplotype 1. Data are presented as the mean (with SD) of 5 and 6 biological repeats for the WT and *tip2;2-1*, respectively. *P < 0.05, unpaired Mann-Whitney test.