



The benefits of a massive harvesting of legacy measured profiles for mapping primary and functional soil properties in the Coastal Plain of Occitanie (Southern France)

Philippe Lagacherie, Léa Courteille, Amine Chemchem

► To cite this version:

Philippe Lagacherie, Léa Courteille, Amine Chemchem. The benefits of a massive harvesting of legacy measured profiles for mapping primary and functional soil properties in the Coastal Plain of Occitanie (Southern France). Centennial of the IUSS, May 2024, Firenze (Florence), Italy. hal-04594527


HAL Id: hal-04594527

<https://hal.inrae.fr/hal-04594527>

Submitted on 30 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Centennial of the IUSS



International Union of Soil Sciences



Florence - Italy
May 19 - 21, 2024

www.centennialiuss2024.org

Philippe Lagacherie, Amine Chemchem & Léa Courteille

The benefits of a massive harvesting of legacy measured profiles for mapping primary and functional soil properties in the Coastal Plain of Occitanie (Southern France)

INRAE

LISAH

Atos

Decision makers needs soil information in the Coastal Plain of Occitanie



Water ressource management



Soil available water capacity



Decision makers needs soil information in the Coastal Plain of Occitanie



Water ressource management



Soil available water capacity



Land use (urban) planning




Soil (potential) quality index



Previous productions of Digital Soil Mapping products in the Coastal plain of Occitanie

Developing GlobalSoilMap products

- 2024 legacy soil profiles in 27,236 km² (1/ 13.5 km²)
- 16 soil covariates
- Quantile Regression Forest



Evaluating Digital Soil Mapping approaches for mapping GlobalSoilMap soil properties from legacy data in Languedoc-Roussillon (France)

K. Vaysse ^{a,b,*}, P. Lagacherie ^a
^a INRA UMR LSMS, Montpellier, France
^b SCG-I, Montpellier, France



Mapping Soil Available Water Capacity

- 2024 legacy soil profiles in 27,236 km² (1/ 13.5 km²)
- 16 soil covariates
- Quantile Regression Forest
- Uncertainty propagation using 1rst order Taylor analysis



Uncertainty assessment of soil available water capacity using error propagation: A test in Languedoc-Roussillon
Styc Quentin ^{a,b,*}, Lagacherie Philippe ^{b,*}



Mapping a Soil Quality (Multifunctionality) index

- 1229 legacy soil profiles in 12,125 km² (1/ 9.9 km²)
- 161 soil covariates
- Regression Forest and cokriging of residuals
- Uncertainty propagation using stochastics simulations

DOI: 10.1111/cjss.13345

ORIGINAL ARTICLE



A multivariate approach for mapping a soil quality index and its uncertainty in southern France

M. E. Angelini ^{1,2} | G. B. M. Heuvelink ^{3,4} | P. Lagacherie ²




Lack of precision for effective use in decision making

Soil information density: A first order limitation of DSM prediction performances

More Data or a Better Model? Figuring Out What Matters Most for the Spatial Prediction of Soil Carbon



P.D.S.N. Somarathna*,
Budiman Minasny,
Brendan P. Malone,
School of Life and Environmental
Sciences
University of Sydney
New South Wales
Australia

Modeling techniques used in digital soil carbon mapping encompass a variety of algorithms to address spatial prediction problems such as spatial non-stationarity, nonlinearity and multi-collinearity. A given study site can inherit one or more such spatial prediction problems, necessitating the use of a combination of statistical learning algorithms to improve the accuracy of predictions. In addition, the minimum sample size may affect the accuracy of the model predictions. The effect of varying sample size on model accuracy has not been widely studied in pedometrics. To help fill this gap, we examined the behavior of multiple linear regression (MLR), geographically weighted regression (GWR), linear mixed models (LMMs), Cubist regression trees, quantile regression forests (QRFs), and extreme learning machine regression (ELMR) under varying sample sizes. The results showed that for the study site in the Hunter Valley, Australia, the accuracy of digital prediction of soil carbon is



Analysing the impact of soil spatial sampling on the performances of Digital Soil Mapping models and their evaluation: A numerical experiment on Quantile Random Forest using clay contents obtained from Vis-NIR-SWIR hyperspectral imagery

P. Lagacherie^{a,*}, D. Arrouays^b, H. Bourennane^c, C. Gomez^{a,d}, L. Nkuba-Kasanda^a



Model
— OK R
— QRF (green)
— QRF (cyan)
— QRF (purple)

Massive harvesting of legacy measured soil profiles

Soil data rescue projects	Profile numbers	Authors of profiles	profile ages
Projet IGCS 2021	1 192	SIGALES, GeoSOLEAU	1985-2006
Projet TerraOccitania	6 752	CNABRL	1955-92

From 1,229 profiles (1/ 9.9. km²) to 9,135 profiles (1/1.3 km²)



A semi-automatic procedure for soil profile entries in a soil database



Scanned soil profiles
records


(4 different formats) Text recognition
model training

(1 per format)



87.6% of
automatic
recognition
with 99.5%
accuracy

Computer-assisted manual entry of the
remaining uncertain fields



Final checking
and field
conversions




A semi-automatic procedure for soil profile entries in a soil database



Scanned soil profiles
records


(4 different formats) Text recognition
model training

(1 per format)



87.6% of
automatic
recognition
with 99.5%
accuracy

Computer-assisted manual entry of the
remaining uncertain fields




Final checking
and field
conversions




Procedure evaluation

	Manual entry	Semi-automatic entry	Gain
1 record	13 mn	3 mn	10 mn
6752 records	182 working days	57 working days	125 working days




Multivariate DSM approach (Angelini et al, 2023)



Regression
Random Forests






14 predicted properties



Linear models of co-regionalisation

Stochastic simulations of residuals and map combinations into maps of indicators

300 simulated maps of soil indicators




mean

Final indicator maps



Predicted values

CI 90%




uncertainties


Accuracy (MEC) increases after a massive harvesting of legacy soil data

Analytical soil properties


CEC




pH



Sand



Silt




Low density soil dataset


High density soil dataset

« Observational » soil properties

Coarse frag.




Soil depth.




Functional soil properties

AWC




Multifunctionnality index




Digital Soil Mapping products for decision makers

Soil Available Water Capacity 0-100 cm (mm)




Soil Multifunctionnality Index



Spatial aggregation for uncertainty visualization adapted for decision making

1. approach


- Principle : Reduce the overall map uncertainty by spatial aggregation
- Aggregated units obtained by applying an agglomerative clustering algorithm (Murtagh, 1985) on predicted raster maps
- Areal metric : proportion of area of high quality soil (> 10/12)



Spatial aggregation for uncertainty visualization adapted for decision making


1. approach

- Principle : Reduce the overall map uncertainty by spatial aggregation
- Aggregated units obtained by applying an agglomerative clustering algorithm (Murtagh, 1985) on predicted raster maps
- Areal metric : proportion of area of high quality soil ($> 10/12$)



Which is the best compromise ??


Users' preferences of spatial aggregations
(201 potential users x 10 choices among 2 maps)




Spatial aggregation for uncertainty visualization adapted for decision making

2. Example of « good » compromise

% of area with high quality soils



Uncertainty (standard error)



5 km

Conclusions

- Digital Soil mapping products should be oriented toward the fullfillment of local users needs
- The decrease in mapping uncertainty provided by a massive harvesting of legacy soil profiles is significant but limited for certain soil properties
- Spatial aggregation of DSM outputs is a promising solution for communicating mapping uncertainties to decision makers
- Local Digital Soil mapping should be promoted as an efficient approach for organising and boosting the collection of soil knowledge within the future soil governance entities (soil districts?)



Soil Mapping for Sustainable Land Use Planning

3RD JOINT WORKSHOP ON DIGITAL SOIL MAPPING & GLOBAL SOIL MAP

21st - 24th January 2025
Bengaluru, India

ICAR-National Bureau of Soil Survey and Land Use Planning, Nagpur

in association with

Indian Society of Soil Survey and Land Use Planning (ISSLUP), Nagpur

The International Working Group of Digital Soil Mapping & The International Workshop Group of GlobalSoilMap

