



HAL
open science

Investigation of caliciviruses and astroviruses in Gabonese rodents: A possible influence of national and international trade on the spread of enteric viruses

Clark Mbou-Boutambe, Illich Manfred Mombo, Virginie Rougeron, Fanny Degrugillier, Philippe Gauthier, Boris Makanga, Barthélémy Ngoubangoye, Eric M. Leroy, Franck Prugnonne, Larson Boundenga

► To cite this version:

Clark Mbou-Boutambe, Illich Manfred Mombo, Virginie Rougeron, Fanny Degrugillier, Philippe Gauthier, et al.. Investigation of caliciviruses and astroviruses in Gabonese rodents: A possible influence of national and international trade on the spread of enteric viruses. *Infection, Genetics and Evolution*, 2024, 122, pp.105607. 10.1016/j.meegid.2024.105607 . hal-04605931

HAL Id: hal-04605931

<https://hal.inrae.fr/hal-04605931v1>

Submitted on 8 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License



Investigation of caliciviruses and astroviruses in Gabonese rodents: A possible influence of national and international trade on the spread of enteric viruses

Clark Mbou-Boutambe^{a,b,*}, Ilich Manfred Mombo^{c,d}, Virginie Rougeron^e, Fanny Degrugillier^d, Philippe Gauthier^f, Boris Makanga^g, Barthélemy Ngoubangoye^h, Eric M. Leroy^d, Franck Prugnonne^e, Larson Boundenga^{a,i}

^a Unité de Recherche en Écologie de la Santé (URES), Centre Interdisciplinaire de Recherches Médicales de Franceville (CIRMF), Franceville, BP 769, Gabon

^b Ecole Doctorale Régionale d'Afrique Centrale en Infectiologie Tropicale (EDR), Franceville, BP 876, Gabon

^c Unité Émergence des Maladies Virales, Centre Interdisciplinaire de Recherches Médicales de Franceville (CIRMF), Franceville, Gabon

^d Institut de Recherche pour le Développement (IRD), Maladies Infectieuses et Vecteurs, Écologie, Génétique, Évolution et Contrôle (MIVEGEC) (Université de Montpellier—IRD 224—CNRS 5290), Montpellier 34394, France

^e International Research Laboratory-REHABS, CNRS-Université Lyon 1-Nelson Mandela University, Nelson Mandela University George Campus, George 6531, South Africa

^f CBGP, IRD, CIRAD, INRA, Montpellier SupAgro, Université de Montpellier, Montpellier, France

^g Institut de Recherche en Écologie Tropicale (IRET/CENAREST), Libreville, BP 13354, Gabon

^h Centre de Primatologie, Centre Interdisciplinaire de Recherches Médicales de Franceville (CIRMF), Franceville, BP 769, Gabon

ⁱ Département d'Anthropologie, Université de Durham, South Road, Durham DH1 3LE, UK.

ARTICLE INFO

Keywords:

Astrovirus
Calicivirus
Rodent
Rural areas
Zoonoses
Gabon

ABSTRACT

Caliciviruses (*Caliciviridae*) and astroviruses (*Astroviridae*) are among the leading cause of non-bacterial food-borne disease and gastroenteritis in human. These non-enveloped RNA viruses infect a wide range of vertebrate species including rodents. Rodents are among the most important hosts of infectious diseases globally and are responsible for over 80 zoonotic pathogens that affect humans. Therefore, screening pathogens in rodents will be necessary to prevent cross-species transmission to prevent zoonotic outbreaks. In the present study, we screened caliciviruses and astroviruses in order to describe their diversity and whether they harbor strains that can infect humans. RNA was then extracted from intestine samples of 245 rodents and retrotranscribed in cDNA to screen caliciviruses and astroviruses by PCRs. All the samples tested negative for caliciviruses and while astroviruses were detected in 18 (7.3%) samples of *Rattus rattus* species. Phylogenetic analyses based on the RdRp gene showed that all the sequences belonged to *Mamastrovirus* genus in which they were genetically related to *R. rattus* related AstVs previously detected in Gabon or in *Rattus* spp. AstV from Kenya and Asia. These findings suggested that transportation such as land and railway, as well national and international trade, are likely to facilitate spread of AstVs by the dissemination of rodents.

1. Introduction

Despite the progress of medicine, infectious diseases remain a significant challenge for humanity. One explanation for that is the frequent emergence of new infectious diseases (Watkins, 2018). While once rare, the frequency of infectious disease emergence has accelerated in the last 60 years. It is worth noting that approximately 75% of emerging

infectious diseases known to affect humans originate from animals, with many of these diseases being related to wildlife species, including rodents (Taylor et al., 2001). Rodents belong to the order *Rodentia*, which is the most diverse mammalian group, with nearly 2552 species divided into 36 families (Burgin et al., 2018). Rodents are among the most important sources of infectious diseases globally, and carry over 80 zoonotic pathogens (Han et al., 2016). Therefore, surveillance of

* Corresponding author at: Unité de Recherche en Écologie de la Santé (URES), Centre Interdisciplinaire de Recherches Médicales de Franceville (CIRMF), Franceville, BP 769, Gabon.

E-mail address: clarkmbou@gmail.com (C. Mbou-Boutambe).

<https://doi.org/10.1016/j.meegid.2024.105607>

Received 4 April 2024; Received in revised form 15 May 2024; Accepted 21 May 2024

Available online 26 May 2024

1567-1348/© 2024 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

pathogens hosted by rodents in areas shared with humans is useful to detect pathogens that could cross the species barrier to infect humans leading to outbreaks.

Enteric viruses are transmitted via fecal-oral route, either directly or indirectly through contaminated food, water, and surfaces (De Graaf et al., 2017). These viruses reach the gastrointestinal tract and replicate in the enterocytes primarily. Most of the RNA enteric viruses including caliciviruses (CaVs) and astroviruses (AstVs) are non-enveloped, very stable in the environment and can survive for long periods of time under different conditions. They infect a large spectrum of hosts such as human, livestock, domesticated and wild animals. They are mainly responsible for gastroenteritis in humans and animals, but have also been associated with mild to severe extra-intestinal diseases such as respiratory diseases, nephritis and neurological disorders (Robilotti et al., 2015; Rosenberg, 2016; De Benedictis et al., 2011; Bank-Wolf, B. R, et al., 2010).

Caliciviruses belong to the family *Caliciviridae*, which contains 11 recognized genera, including *Bavovirus*, *Lagovirus*, *Norovirus*, *Salovirus* and *Sapovirus* (<https://ictv.global/taxonomy>). Only noroviruses (NoVs) and sapoviruses (SaVs) can infect humans. The reverse-zoonotic potential of CaVs has been described. Some NoVs and SaVs found in animals are related to human strains (Mattison et al., 2007; Martella et al., 2008; Mombo et al., 2014). Despite that, there is no evidence of animal-related NoVs and SaVs detected in humans (Mombo et al., 2019; Bohou Kombila et al., 2023).

Astroviruses (AstVs) belong to the genera *Avastrovirus* (AAstVs) for AstVs infecting birds and *Mamastrovirus* (MAstV) for those infecting mammals within the family *Astroviridae*. Since the detection of human astroviruses of the groups HAstV-VA/HMO and HAstV-MLB, which are genetically close to animal AstVs but divergent from classical human astroviruses (HAstV-1 to 8), the zoonotic origin of AstVs has been suggested (Finkbeiner et al., 2008; Kapoor et al., 2009).




Fig. 1. Location of rodent sampling sites in Gabon. On this map, only the main roads (orange) in the sample area have been shown.

NoVs, SaVs and AstVs are among the most common causes of food-borne outbreaks of non-bacterial diarrhea and viral acute gastroenteritis therefore are interest for public health (Segone, 2020, Ibrahim et al., Ibrahim et al., 2022, Kowada et al., 2018, Finkbeiner et al., 2009). In Gabon, AstVs have been widely studied in animals including rodents (Mombo et al., 2019). By contrast, there are no data regarding the diversity and the circulation of CaVs in rodents specifically NoVs and SaVs in this country. Thus, this study aimed to screen caliciviruses and astroviruses in rodents from various locations from Gabon to characterize their diversity and to determine whether they can harbor strains capable of infecting humans.

2. Materials and methods

2.1. Ethical statement

Rodent trapping campaigns were carried out following authorization by the Commission Scientifique d'Examen and the application for an Exploration Licence by Gabon (N°AR005/20/MESRSTT/CENAREST/CG/CST/CSAR). The capture, handling and euthanasia of animals and the transfer of samples from one country to another, were carried out in accordance with the guidelines of the American Society of Mammalogists (<http://www.mammalsociety.org/committees/animal-care-and-use>) and in strict compliance with the recommendations of the National Ethics Committee of Gabon (authorization N°PROT/0020/2013I/S G/CNE).

2.2. Study areas and capture

Between October 2021 and June 2022, sampling of small terrestrial rodents was carried out in several rural areas in Gabon (Hereafter referred as Dienga, Okondja, Lastourville, Booué and Ndjolé, Fig. 1). Inside each of these areas, several sampling sessions were performed in different villages (Table 1). Two types of traps were used: Sherman traps and Tomahawk traps (Granjon and Duplantier, 2009). The traps were set in houses and in the forest (Dienga and Ndjolé only) for three consecutive nights. Rodents caught were sacrificed by cervical dislocation, then measured, sexed and dissected. Tissues (brain, lung, intestine, liver) and blood were collected for the detection of pathogens circulating among small mammals. Samples were then and stored in liquid nitrogen until storage in -80°C freezer for further molecular analyses at CIRMF.

2.3. Rodent species identification

Rodent species were identified by morphometric (Granjon and Duplantier, 2009) and genetic tools (partial *cytochrome b* Sanger sequencing) (Nicolas et al., 2012) to avoid any ambiguity in the identification. Due to the existence of cryptic species in Central African rodents, care was taken to identify most, if not all, of the small mammals captured by molecular analyses, using kidney or phalanx samples.

2.4. RNA extraction and reverse transcription

Of the 717 small mammals captured and for costs limitations a

Table 1
Small mammal species sampled in five locality rural habitats and forest of Gabon.

Localities	Number sites	Number house	Trap nights	Total Captures
Booué	5	42	1314	99
Dienga	1	94	1143	149
Lastourville	5	37	886	122
Ndjolé	12	44	1344	165
Okondja	4	44	714	182
Total	27	261	5401	717

subsample of 245 rodents was chosen to test for the presence of calicivirus and astrovirus. For this purpose, we chose to keep all the rodents captured in Dienga because of the cohabitation between native and invasive species, with a view to determining the exchange of enteric viruses between these two groups of rodents. Finally, we randomly sampled a subset of *Rattus rattus* in the other localities. Approximately 100 mg of intestine of each individual was grinded in 500 μL of cold phosphate- saline buffer (PBS), as previously described by Rougeron et al., 2016 (Rougeron et al., 2016). RNA was extracted from the suspension using the RNA isolation kit Nucleospin® RNA, Macherey-Nagel (Düren, Germany), according to the manufacturer's instructions. The extracted RNA was then reverse-transcribed into cDNA using the Superscript IV Reverse Transcriptase (Invitrogen, Illkirch, France) in a final volume of 20 μL . Specifically, 10 μL of RNA in a mix consisting of 1 μL dNTPs, 1 μL of random hexamer and 1 μL of DNase-free water, was incubated at 65°C for 5 min followed by 1 min on ice. A second mixture consisting of 4 μL of 5 \times Superscript IV Buffer, 1 μL of DTT (100 mM), 0.5 μL of RNase Out, 0.5 μL of enzyme and 1 μL of DNase-free water, was added to the previous mixture. The reverse-transcription program was set to 23°C for 10 min, $50\text{--}55^{\circ}\text{C}$ for 10 min, and 80°C for 10 min.

2.5. Calicivirus and astrovirus detection

Caliciviruses and astroviruses were screened respectively by PCR and nested PCR using Platinum Taq DNA Polymerase (Invitrogen) according to the manufacturer's instructions with primers targeting a fragment of (319–331) bp of the RdRp of CalVs and a fragment of 422 bp of the RdRp of AstVs (Table 2). For CalCVs the PCR mix was performed in a final volume of 25 μL , consisting of 5 μL of cDNA, 2.5 μL of 10 X reaction buffer, 0.75 μL of MgCl_2 (50 mM), 0.5 μL of dNTPs, 1 μL of bovine serum albumin (BSA) (1 $\mu\text{g}/\mu\text{L}$), 1 μL of each primer (10 μM), and 0.1 μL of Platinum Taq Enzyme. Similarly, for the detection of AstVs, both rounds of the nested PCR, were performed with Platinum Taq DNA polymerase using the same components and volumes used for CaClVs with the exception of BSA (not included) in the second round. Amplification programs for AstVs and CaVs were set up as previously described (Jiang et al., 1999; Mombo et al., 2019), without the reverse-transcription step for the first round. The PCR products were visualized on a 1.5% agarose gel after electrophoresis and the amplicons were sent to Eurofins Genomics (Ebersberg, Germany) for Sanger sequencing in both directions.

2.6. Phylogenetic analyses

Phylogenetic analyses were performed on the RdRp gene sequences obtained in this study (Genbank accession numbers PP516503 to PP516520) with those of AstVs referenced on Genbank. Sequences were aligned using the ClustalW algorithm implemented in MEGA11 software package. Phylogenetic trees were constructed by Maximum likelihood (freely available at <http://phylogeny.lirmm.fr>) (Dereeper et al., 2008) using Kimura 2-parameter + Gamma distribution (K2 + G) estimated with MEGA11 (Anisimova and Gascuel, 2006) and 100 bootstrap were used to test for the robustness of each node in the tree.

2.7. Statistic analyses and locality mapping

Statistical analyses were performed using R software (v.3.2.1). A chi-square test was performed to determine the influence of sex and age on AstV infection. Maps were generated using ArcGIS 10.3 software (Environmental Systems Research Institute).

3. Results

3.1. Rodent trapping

A trapping effort of 5401 trap.nights allowed the capture of 717 individuals (Table 1). The 245 rodents analysed belong to eleven species

Table 2
Sequences of primers used for the detection of *Astroviruses* and *Caliciviruses*.

Target Virus Name	Gene	Primer Name	Primer Sequence (50–30)	References
Astrovirus	RdRp	FWD1	GARTTYGATTGGRCKGKTAYGA	422 bp (Chu et al., 2008)
		FWD2	GARTTYGATTGGRCKAGGTAYGA	
		RVS1	GGYTTKACCCACATNCCRAA	
		FWD3	CGKTAYGATGGKACKATHCC	
		FWD4	AGGTAYGATGGKACKATHCC	
Calicivirus	RdRp	P289	TGACAATGTAATCATCACCATA	319–331 bp (Jiang et al., 1999)
		P290	GATTACTCCAAGTGGGACTCCAC	

already described in Gabon ([Mangombi et al., 2021](#); [Duplantier, 1982](#); [Duplantier, 1989](#); [Nicolas et al., 2012](#)). The species *Rattus rattus* was found in all localities: Booué 24.2% (24/99), Dienga 65.8% (98/149), Lastourville 18.8% (23/122), Ndjolé 23% (23/113) and Okondja 15.9% (29/182). In contrast, the 48 native rodents captured in this study came from two localities: Dienga ($N = 46$) and Lastourville ($N = 2$). The native rodents captured in Dienga were: *Hybomys univittatus* ($N = 6$), *Mus minutoides* ($N = 3$), *Hylomyscus aeta* ($N = 1$), *Lophuromys roseveari* ($N = 15$), *Malacomys longipes* ($N = 2$), *Mylomys cf. dybowski* ($N = 1$), *Oenomys hypoxanthus* ($N = 2$), *Praomys petteri* ($N = 1$), *Praomys jacksoni* ($N = 9$) and *Lemniscomys striatus* ($N = 6$). In Lastourville, two native rodents: *Praomys jacksoni* ($N = 1$) and *Lemniscomys striatus* ($N = 1$) were found ([Table 3](#)).

3.2. Caliciviruses and astroviruses detection

A total of 245 intestine samples were tested for the presence RNA of CalCVs and AstVs. None of the samples were tested positive for CalCVs, while AstVs were detected in 18 samples (detection rate of 7.3%). According to the rodent species, *R. rattus* species was the one species tested positive and found in every locality with detection rates according to the localities as 21.7% (5/23) in Lastourville, 16.0% (4/25) in Booué, 13.8% (4/29) in Okondja and 13.0% (3/23) in Ndjolé. No factor significantly influenced astrovirus infection: males with 9.8% (12/122) had an infection rate close to that of females 4.9% (6/123; $\chi^2 = 1.5434$, p -value = 0.2141). Similarly, adult *Rattus rattus* had an AstV infection rate of 7.7% (10/130) similar to the 7.0% (8/115) found in juveniles ($\chi^2 = 8.6037e-31$, p -value = 1).

3.3. Phylogenetic analyses

In order to describe the genetic diversity of AstV strains detected in *R. rattus* in this study, the partial RdRp sequences were compared to reference sequences retrieved from Genbank. Our results showed that all the sequences of this study clustered with other sequences of AstVs detected in rodents ([Fig. 2](#)). Specifically, phylogenetic analyses showed that some sequences are specific to Gabon. Indeed, five sequences

including three from rats trapped in Lastourville (RLT-103, RLT-114, RLT-118), two from Booué (RB-078, RB-082) clustered with AstVs previously described in rats from Franceville (Genbank accession numbers MF741182, MF741183 and MF741185) ([Mombo et al., 2019](#)). The sequences shared nucleotide identities ranging from 89% to 100%. In addition, an AstV sequence found at Dienga (RD-083) was phylogenetically divergent (89% nucleotide identities) from the AstV sequences grouped within this group. Five other sequences (ROK-023, RB-095, RB-098, NDJ-149 and NDJ-164) from rats trapped in Okondja, Booué and Ndjolé, clustered with a *R. rattus*-related AstV (MF741187) previously detected in Makokou. These sequences formed the second clade of rodent-borne AstVs sequences from Gabon that shared 90 to 99% nucleotide identities. Besides, the phylogenetic analyses also showed that some sequences are grouped with AstVs identified outside Gabon. Indeed, three sequences (RD-011, RLT-110 and RLT-112) clustered with sequences of AstVs detected in *Rattus rattus* in Kenya and in Reunion free tailed bats ([Joffrin et al., 2021](#)). Despite, the sequences were divergent showing nucleotide identities ranged from 85% to 100% ([Fig. 2](#)). Moreover, two sequences, one from Ndjolé (NDJ-061) and the other from Okondja (ROK-136), were grouped with AstV sequences found in China in *Rattus tanesumia*, *Rattus norvegicus*, *Rattus losea* (Genbank accession numbers OP482212, KY370002 and HM450382). These sequences showed 88–100% nucleotide identity). Finally, the last two sequences (ROK-041 and ROK-080) also clustered with sequences of AstVs detected in China in various species of *Rattus* genus such as *R. norvegicus*, *R. tanezumi* but also from the insectivorous bat species *Myotis fimbriatus*. The sequences shared 89% to 93% nucleotides identities with sequences of AstVs detected in the other *Rattus* species ([Fig. 2](#)).

4. Discussion

Caliciviruses (CaVs, such as noroviruses and sapoviruses) and astroviruses (AstVs) are common causes of diarrhea in human populations ([Finkbeiner et al., 2009](#)). These viruses infect various vertebrate species including rodents. Rodents, especially in domestic and peri-domestic contexts, could then disseminate these viruses and humans

Table 3
Results of screening for *Astroviruses* in the various locations sampled.

Rodent species	Booué	Dienga	Lastourville	Ndjolé	Okondja	Total
	<i>Astrovirus</i> Positive/Tested					
<i>Rattus rattus</i>	4/24 (16.7%)	2/98 (2%)	5/23 (21.7%)	3/23 (13%)	4/29 (13.8%)	18/197 (9.1%)
<i>Praomys jacksoni</i>	–	0/9 (0%)	0/1 (0%)	–	–	0/10 (0%)
<i>Praomys petteri</i>	0/1 (0%)	–	–	–	–	0/1 (0%)
<i>Mus minutoides</i>	–	0/3 (0%)	–	–	–	0/3 (0%)
<i>Hybomys univittatus</i>	–	0/6 (0%)	–	–	–	0/6 (0%)
<i>Hylomyscus aeta</i>	–	0/1 (0%)	–	–	–	0/1 (0%)
<i>Lophuromys roseveari</i>	–	0/15 (0%)	–	–	–	0/15 (0%)
<i>Lemniscomys striatus</i>	–	0/6 (0%)	0/1 (0%)	–	–	0/7 (0%)
<i>Mylomys dybowski</i>	–	0/1 (0%)	–	–	–	0/1 (0%)
<i>Malacomys longipes</i>	–	0/2 (0%)	–	–	–	0/2 (0%)
<i>Oenomys hypoxanthus</i>	–	0/2 (0%)	–	–	–	0/2 (0%)
Total	4/25 (16.0%)	2/143 (1.4%)	5/25 (20%)	3/23 (13%)	4/29 (13.8%)	18/245 (7.3%)

could be infected directly or indirectly via a zoonotic transmission (Milton et al., 2020). Little is known about the diversity and spillover potential of these viruses carried by rodents on a large scale in Gabon, particularly caliciviruses. In the 245 intestine samples from eleven rodent species trapped in forest and in human houses in five Gabonese localities and screened for the presence of CaVs and AstVs, only AstVs were detected in *R. rattus*. None of the samples tested were positive for CaVs. Among CaVs noroviruses are the one only detected in rodents. However most of the studies have focused on noroviruses due to their significance in public health, and to serve of as model to understand human infection (Manuel et al., 2018; Barclay et al., 2014). Rodent noroviruses were first identified in laboratory mice in 2003 and subsequently in other laboratory and also wild mice (Farkas et al., 2012; Karst et al., 2003). In addition to laboratory mice, NoVs have also been detected in wild rodent such as *R. rattus* and *Apodemus* spp., and recently in *R. norvegicus* (Tsunesumi et al., 2012; Farkas et al., 2012). All NoVs detected were specific to rodents, except in *R. norvegicus* in which human-related strains were also detected (Niendorf et al., 2021). The use of specific primers targeting NoV RNA in previous studies may have favored detection. Thus, the absence of detection in this study could be explained not only by the use of a more generic system, but also by the low number of samples tested.

In our study, 18 rodents (7.3%) were found positive for AstVs, which is a higher detection rate than in previous studies where AstVs were detected in rodents captured in Franceville and Makokou, two urban cities in Gabon (4.6%) (Mombo et al., 2019). In the previous study, AstVs were detected in three species *Hybomys univittatus*, *Mus musculus* and *R. rattus* while only *R. rattus* was found to be positive in our study. For the *R. rattus* species, the prevalence seemed to be relatively the same, i.e. 10.5% in the previous study and 9.1% in the present one. The absence of AstVs in other species could be explained by the small sample sizes obtained during trapping, as well as the fact that no other rodent species were captured in all localities except Dienga. Yet, our findings showed that the prevalence of astroviruses in Dienga was significantly lower than in other localities. The low prevalence of AstV in Dienga could be explain by several factors. Firstly, the sample size taken in this locality was larger than in the other sites. The discrepancy in sample size may have resulted in an underestimation of the true prevalence of infection. Furthermore, Dienga is a remote forested village with limited interactions with other localities. This characteristic could serve to limit the circulation of AstV in rodents in Dienga in comparison with other sites, which would also contribute to the low prevalence observed.

Concerning the age class, the results indicate that the prevalence of AstV between young and adult *R. rattus* does not differ. This suggests that age is not a significant factor in determining the prevalence of Astrovirus infection in *R. rattus* rodents. Nevertheless, studies in humans have demonstrated that astroviruses are responsible for gastroenteritis in young children and the elderly, with varying susceptibility depending on the age group (Korsman et al., 2014; Vu et al., 2017). Although astroviruses are known to affect children and the elderly in humans (Kurtz and Lee, 1978; Nguengkeng Tsague et al., 2020), no study has demonstrated a difference in infection rates between the two age groups. It is unclear whether there is a difference in the rate of infection between young and adult mammals, particularly (Chu et al., 2010; To, K. K. W, et al., 2017). In light of the currently available data, it is not possible to draw any conclusions regarding the influence of age in astrovirus infection in rodents, further studies are then needed.

With regard to sex, although our results are not statistically significant, we nevertheless show that AstV appears to be more prevalent in males than females. This observation is in line with the fact male rats tend to persist longer in foraging areas than females, indicating a sex difference in foraging-related decision making strategies (Garcia et al., 2023).

Phylogenetic analyses based on the RdRp gene revealed that all sequences obtained in this study belong to the genus *Mamastrovirus* with host restriction, as these sequences fell into the large clade of rodent

astroviruses. Moreover in this big clade, all *R. rattus* AstVs sequences always clustered with other rat AstVs. The phylogenetic analyses also revealed that some sequences were specific to Gabon. Firstly, some AstV sequences from rats captured in Lastourville and Booué were genetically close to AstVs previously described in *R. rattus* from Franceville (Mombo et al., 2019). Secondly, some other sequences from Booué, Ndjolé, and Okondja clustered with AstV sequences previously described in *R. rattus* captured in Makokou (Mombo et al., 2019). The phylogenetic groups correspond to the distribution of communities along the single line passing through Booué, Lastourville and Franceville, as well as one of the land routes that includes Ndjolé, Booué, Makokou and Okondja. These routes are crucial for transporting food and supplies to the country's various cities and communities. Therefore, these transport routes could play a significant role in the spread of AstVs, by facilitating the proliferation of rodents. This is consistent with studies conducted in Senegal, which suggest that road transport facilitates the spread of black rats between different areas, and even the dissemination of the pathogens they may harbor (Lombard, 2017; Lucaccioni et al., 2016). In Gabon, during the double epidemic of chikungunya and dengue fever between 2007 and 2010, the railroads may have contributed to the spread of chikungunya and dengue viruses, by enabling infected mosquitoes to spread more rapidly. Cases have appeared in communities located along the railway line over time (Nkoghe et al., 2012). Additionally, an AstV sequence from Dienga (RD-083) was highly divergent sharing 89% nucleotide identities with AstVs specific to Gabon from the AstV sequences grouped together in this group. This suggest that this could be a new genotype. However, a complete analysis of the full genome and capsid is necessary to draw a conclusion.

Phylogenetic analyses also revealed that three sequences (RLT-110, RD-011 and RLT-112) grouped with AstVs of *R. rattus* identified in Kenya which formed a cluster phylogenetically close to AstVs of Reunion fruit bats. The detection of the AstVs strains in different host species (bats and rodents) but phylogenetically close indicates potential exchanges. In Reunion, for example, rodents are seen in the vicinity of caves where bat guanos have been tested positive (Joffrin et al., 2021). Rodents and bats also interact near fruit trees and dwellings, consequently such interactions could favor the exchange of AstVs. These findings raise questions about the nomenclature of AstVs, which are typically named after the host in which they were first detected, which therefore minimize the possibility of passive transmissions of the virus or external contamination. Consequently, Kerstin et al. (2016) recommended carrying out successful virus isolations in these species as evidence of productive infection (Fischer et al., 2016).

Finally, phylogenetic analyses showed that two sequences cluster with a groups of AstVs detected in diverse host species within the genus *Rattus* such as *R. rattus*, *R. norvegicus*, *R. tanezumi*, and *R. losea* from China (Wu et al., 2018). This finding suggests the intra-species transmission of AstVs within the *Rattus* genus that coevolved with their host. Wu et al. (2018) also suggested that frequent host shifts during the viral evolutionary history, which may have created opportunities for the emergence of new viruses that can adapt to new hosts (Wu et al., 2018).

5. Conclusions

This study did not detect CalVs in rodents and a high host restriction for AstVs, which were detected only in black rats *R. rattus*. The dissemination of AstVs could be facilitated by railroads and roads, a hypothesis that could be tested by screening samples from localities outside the main network of circulation. The detection of rodent AstVs phylogenetically close to those infecting bats suggests events of inter-species transmission. This finding challenges the current nomenclature of AstVs, which mainly focuses on the host species. Finally, AstVs may have evolved within the genus *Rattus*, allowing an adaptation to each host species. However, further studies based on the complete genome could bring more information regarding the evolution of rodent AstVs in rodent in Gabon.

CRedit authorship contribution statement

Clark Mbou-Boutambe: Writing – original draft, Validation, Methodology, Investigation, Formal analysis, Data curation. **Illich Manfred Mombo:** Writing – original draft, Visualization, Validation, Supervision, Methodology, Formal analysis, Data curation, Conceptualization. **Virginie Rougeron:** Writing – review & editing, Resources. **Fanny Degrugillier:** Investigation. **Philippe Gauthier:** Investigation. **Boris Makanga:** Investigation. **Barthélemy Ngoubangoye:** Investigation. **Eric M. Leroy:** Investigation. **Franck Prugnolle:** Validation, Resources, Project administration, Funding acquisition. **Larson Boundenga:** Writing – review & editing, Validation, Resources, Project administration, Funding acquisition.

Declaration of competing interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Data availability

The original contributions presented in the study are included in the article, further inquiries can be directed to the corresponding author. The partial genome/coding sequences of study viruses can be found in the National Center for Biotechnology Information Database/ GenBank under the accession numbers PP516503 to PP516520. Access to the fasta files can be provided upon request.

Acknowledgements

This study has been funded by ANR MICETRAL (Project number ANR-19-35CE-0010) and Word Organization for animal Health (WOHA) through the European Union (Capacity building and surveillance for Ebola Virus Disease, EBO-SURSY: FOOD/2016/379-660). We thank the Gabonese Government and the National Research Agency (NRA) for their financial support. We are also grateful to people who consented the captures of rodents in their houses. We would also like to thank the reviewers whose critical comments helped improve this document.

References

- Anisimova, M., Gascuel, O., 2006. Approximate likelihood-ratio test for branches: a fast, accurate, and powerful alternative. *Syst. Biol.* 55, 539–552.
- Bank-Wolf, B. R., König, M., Thiel, H.-J., 2010. Zoonotic aspects of infections with noroviruses and sapoviruses. *Vet. Microbiol.* 140, 204–212.
- Barclay, L., Park, G., Vega, E., Hall, A., Parashar, U., Vinjé, J., Lopman, B., 2014. Infection control for norovirus. *Clin. Microbiol. Infect.* 20, 731–740.
- Bohou Kombila, L., N'dilimabaka, N., Garcia, D., Rieu, O., Engone Ondo, J.D., Ndong Mebale, T., Boundenga, L., Fritz, M., Lenguiya, L.H., Maganga, G.D., 2023. Molecular identification of enteric viruses in domestic animals in northeastern Gabon. *Central Africa. Animals* 13, 2512.
- Burgin, C.J., Colella, J.P., Kahn, P.L., Upham, N.S., 2018. How many species of mammals are there? *J. Mammal.* 99, 1–14.
- Chu, D., Poon, L., Guan, Y., Peiris, J., 2008. Novel astroviruses in insectivorous bats. *J. Virol.* 82, 9107–9114.
- Chu, D.K., Chin, A.W., Smith, G.J., Chan, K.H., Guan, Y., Peiris, J.S., Poon, L.L., 2010. Detection of novel astroviruses in urban brown rats and previously known astroviruses in humans. *J. Gen. Virol.* 91, 2457–2462.
- de Benedictis, P., Schultz-Cherry, S., Burnham, A., Cattoli, G., 2011. Astrovirus infections in humans and animals—molecular biology, genetic diversity, and interspecies transmissions. *Infect. Genet. Evol.* 11, 1529–1544.
- De Graaf, M., Beck, R., Caccio, S.M., Duim, B., Fraaij, P.L., Le Guyader, F.S., Lecuit, M., Le Pendu, J., de Wit, E., Schultze, C., 2017. Sustained fecal-oral human-to-human transmission following a zoonotic event. *Curr. Opin. Virol.* 22, 1–6.
- Dereeper, A., Guignon, V., Blanc, G., Audic, S., Buffet, S., Chevenet, F., Dufayard, J.-F., Guindon, S., Lefort, V., Lescot, M., 2008. Phylogeny. Fr: robust phylogenetic analysis for the non-specialist. *Nucleic Acids Res.* 36, W465–W469.
- Duplantier, J.-M., 1982. Les rongeurs myomorphes forestiers du nord-est du Gabon. *Univ. de Montpellier* 1.
- Duplantier, J.-M., 1989. Les rongeurs myomorphes forestiers du nord-est du Gabon: structure du peuplement, démographie, domaines vitaux. *Revue d'Ecologie, Terre et Vie* 44, 329–346.

- Farkas, T., Fey, B., Keller, G., Martella, V., Egyed, L., 2012. Molecular detection of murine noroviruses in laboratory and wild mice. *Vet. Microbiol.* 160, 463–467.
- Finkbeiner, S.R., Kirkwood, C.D., Wang, D., 2008. Complete genome sequence of a highly divergent astrovirus isolated from a child with acute diarrhoea. *Virol. J.* 5, 1–7.
- Finkbeiner, S.R., Holtz, L.R., Jiang, Y., Rajendran, P., Franz, C.J., Zhao, G., Kang, G., Wang, D., 2009. Human stool contains a previously unrecognized diversity of novel astroviruses. *Virol. J.* 6, 161.
- Fischer, K., Zeus, V., Kwasnitschka, L., Kerth, G., Haase, M., Groschup, M.H., Balkema-Buschmann, A., 2016. Insectivorous bats carry host specific astroviruses and coronaviruses across different regions in Germany. *Infect. Genet. Evol.* 37, 108–116.
- Garcia, M., Gupta, S., Wikenheiser, A.M., 2023. Sex differences in patch-leaving foraging decisions in rats. *Oxf. Open Neurosci* 2, kvad011.
- Granjon, L., Duplantier, J.-M., 2009. Les rongeurs de l'Afrique sahélo-soudanienne. *IRD Marseille, French*.
- Han, B.A., Kramer, A.M., Drake, J.M., 2016. Global patterns of zoonotic disease in mammals. *Trends Parasitol.* 32, 565–577.
- Ibrahim, Y.M., Werid, G.M., Zhang, H., Fu, L., Wang, W., Chen, H., Wang, Y., 2022. Potential zoonotic swine enteric viruses: the risk ignored for public health. *Virus Res.* 315, 198767.
- Jiang, X., Huang, P., Zhong, W., Farkas, T., Cubitt, D., Matson, D., 1999. Design and evaluation of a primer pair that detects both Norwalk-and Sapporo-like caliciviruses by RT-PCR. *J. Virol. Methods* 83, 145–154.
- Joffrin, L., Hoarau, A.O., Lagadec, E., Köster, M., Ramanantsalama, R.V., Mavingui, P., Lebarbenchon, C., 2021. Astrovirus in Reunion free-tailed bat (*Mormopterus francoismoutoui*). *Viruses* 13, 1524.
- Kapoor, A., Li, L., Victoria, J., Oderinde, B., Mason, C., Pandey, P., Zaidi, S., Delwart, E., 2009. Multiple novel astrovirus species in human stool. *J. Gen. Virol.* 90, 2965.
- Karst, S.M., Wobus, C.E., Lay, M., Davidson, J., Virgin IV, H. W., 2003. STAT1-dependent innate immunity to a Norwalk-like virus. *Science* 299, 1575–1578.
- Korsman, S.N.J., Van Zyl, G., Nutt, L., Andersson, M.I., Preiser, W., 2014. *Virologia* (Elsevier Editora Ltda).
- Kowada, K., Takeuchi, K., Hirano, E., Toho, M., Sada, K., 2018. Development of a multiplex real-time PCR assay for detection of human enteric viruses other than norovirus using samples collected from gastroenteritis patients in Fukui prefecture, Japan. *J. Med. Virol.* 90, 67–75.
- Kurtz, J., Lee, T., 1978. Astrovirus gastroenteritis age distribution of antibody. *Med. Microbiol. Immunol.* 166, 227–230.
- Lombard, J., 2017. *Le monde des transports sénégalais: ancrage local et développement international* (IRD Éditions).
- Lucaccioni, H., Granjon, L., Dalecky, A., Fossati, O., Le Fur, J., Duplantier, J.-M., Handschumacher, P., 2016. From human geography to biological invasions: the black rat distribution in the changing southeastern of Senegal. *PLoS One* 11, e0163547.
- Mangombi, J.B., N'dilimabaka, N., Lekana-Douki, J.-B., Banga, O., Maghendji-Nzondo, S., Bourgarel, M., Leroy, E., Fenollar, F., Mediannikov, O., 2021. First investigation of pathogenic bacteria, protozoa and viruses in rodents and shrews in context of forest-savannah-urban areas interface in the city of Franceville (Gabon). *PLoS One* 16, e0248244.
- Manuel, C.S., Moore, M.D., Jaykus, L.-A., 2018. Predicting human norovirus infectivity-relevant advances and continued challenges. *Food Microbiol.* 76, 337–345.
- Martella, V., Lorusso, E., Banyai, K., Decaro, N., Corrente, M., Elia, G., Cavalli, A., Radogna, A., Costantini, V., Saif, L., 2008. Identification of a porcine calicivirus related genetically to human sapoviruses. *J. Clin. Microbiol.* 46, 1907–1913.
- Mattison, K., Shukla, A., Cook, A., Pollari, F., Friendship, R., Kelton, D., Bidawid, S., Farber, J.M., 2007. Human noroviruses in swine and cattle. *Emerg. Infect. Dis.* 13, 1184.
- Milton, A.A.P., Priya, G.B., Ghatak, S., Das, S., 2020. Viral Zoonoses: wildlife perspectives. *Animal-Origin Viral Zoonoses* 339–378.
- Mombo, I.M., Berthet, N., Bouchier, C., Fair, J.N., Schneider, B.S., Renaud, F., Leroy, E. M., Rougeron, V., 2014. Characterization of a genogroup I sapovirus isolated from chimpanzees in the republic of Congo. *Genome Announc.* 2 <https://doi.org/10.1128/genomea.00680-14>.
- Mombo, I.M., Suquet, E., Boundenga, L., Mveang-Nzoghé, A., Maganga-Mboga, C., Arnathau, C., Sidobre, C., Leroy, E.M., Rougeron, V., 2019. Detection of novel astroviruses among rodents of Gabon, Central Africa. *Infect. Genet. Evol.* 68, 43–46.
- Nguekeng Tsague, B., Mikounou Louya, V., Ntoumi, F., Adedoja, A., Vouvongui, C.J., Peko, S.M., Abena, A.A., 2020. Occurrence of human astrovirus associated with gastroenteritis among Congolese children in Brazzaville, republic of Congo. *Int. J. Infect. Dis.* 95, 142–147.
- Nicolas, V., Schaeffer, B., Missoup, A.D., Kennis, J., Colyn, M., Denys, C., Tatard, C., Cruaud, C., Laredo, C., 2012. Assessment of three mitochondrial genes (16S, Cytb, CO1) for identifying species in the Praomyini tribe (Rodentia: Muridae). *PLoS One* 7, e36586.
- Niendorf, S., Harms, D., Hellendahl, K.F., Heuser, E., Böttcher, S., Jacobsen, S., Beck, C.-T., Ulrich, R.G., 2021. Presence and diversity of different enteric viruses in wild Norway rats (*Rattus norvegicus*). *Viruses* 13, 992.
- Nkoghe, D., Kassa, R.F., Caron, M., Grand, G., Mombo, I., Bikié, B., Paupy, C., Becquart, P., Bisvigou, U., Leroy, E.M., 2012. Clinical forms of chikungunya in Gabon, 2010. *PLoS Negl. Trop. Dis.* 6, e1517.
- Robilotti, E., Deresinski, S., Pinsky, B.A., 2015. Norovirus. *Clin. Microbiol. Rev.* 28, 134–164.
- Rosenberg, J.J., 2016. *Norovirus and Other Human Calicivirus Infections*. American Academy of Pediatrics.
- Rougeron, V., Suquet, E., Maganga, G.D., Jiolle, D., Mombo, I.M., Bourgarel, M., Motsch, P., Arnathau, C., Durand, P., Drexler, F., Drosten, C., Renaud, F.,

- Prugnolle, F., Leroy, E.M., 2016. Characterization and phylogenetic analysis of new bat astroviruses detected in Gabon, Central Africa. *Acta Virol.* 60, 386–392.
- Segone, N., 2020. Zoonotic Potential of Viral Enteric Pathogens in South African Domesticated Animals.
- Taylor, L.H., Latham, S.M., Woolhouse, M.E., 2001. Risk factors for human disease emergence. *Philosophical transactions of the Royal Society of London. Series B: Biological Sciences* 356, 983–989.
- To, K.K.W., Chan, W.M., Li, K.S.M., Lam, C.S.F., Chen, Z., Tse, H., Lau, S.K.P., Woo, P.C.Y., Yuen, K.Y., 2017. High prevalence of four novel astrovirus genotype species identified from rodents in China. *J. Gen. Virol.* 98, 1004–1015.
- Tsunesumi, N., Sato, G., Iwasa, M., Kabeya, H., Maruyama, S., Tohya, Y., 2012. Novel murine norovirus-like genes in wild rodents in Japan. *J. Vet. Med. Sci.* 74, 1221–1224.
- Vu, D.L., Bosch, A., Pintó, R.M., Guix, S., 2017. Epidemiology of classic and novel human Astrovirus: gastroenteritis and beyond. *Viruses* 9.
- Watkins, K., 2018. Emerging infectious diseases: a review. *Curr. Emerg. Hosp. Med. Rep.* 6, 86–93.
- Wu, Z., Lu, L., Du, J., Yang, L., Ren, X., Liu, B., Jiang, J., Yang, J., Dong, J., Sun, L., 2018. Comparative analysis of rodent and small mammal viromes to better understand the wildlife origin of emerging infectious diseases. *Microbiome* 6, 178.