



HAL
open science

Climate change altered the dynamics of stand dominant height in forests during the past century

Matthieu Combaud, Thomas Cordonnier, Sylvain Dupire, Patrick Vallet

► To cite this version:

Matthieu Combaud, Thomas Cordonnier, Sylvain Dupire, Patrick Vallet. Climate change altered the dynamics of stand dominant height in forests during the past century. British Ecological Society annual meeting 2023, BES, Dec 2023, Belfast, United Kingdom. pp.1-36. hal-04606325

HAL Id: hal-04606325

<https://hal.inrae.fr/hal-04606325>

Submitted on 10 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License



Métaprogramme CLIMAE

➤ Climate change altered the dynamics of stand dominant height in forests during the past century


Analysis of 20 European tree species

Matthieu Combaud, Thomas Cordonnier, Sylvain Dupire, Patrick Vallet

BES Annual Meeting, Belfast, 13th December 2023



Need to anticipate forest dynamics response to climate change (CC)




Need to anticipate forest dynamics response to climate change (CC)

Impact of CC



Focus on even-aged pure stands




Stand = uniform group of trees

Pure stand = a single species

Even-aged = similar ages for all trees

Focus on even-aged pure stands



Stand = uniform group of trees


Pure stand = a single species

Even-aged = similar ages for all trees

Two key indicators

- **Dominant height** = height of the biggest trees
- **Site index** = dominant height at 70 yrs

Focus on even-aged pure stands



Stand = uniform group of trees

Pure stand = a single species

Even-aged = similar ages for all trees

Two key indicators

- **Dominant height** = height of the biggest trees


- **Site index** = dominant height at 70 yrs

Informs on
productivity

Little influenced
by density





Influences response
to disturbances

Literature on climate impacts on dominant height and site index



← Impact of mean climate on site index and dominant height

Temporal trends in height growth dynamics



CC impact depends on stand climatic conditions


How has climate change over the past century modified dominant height dynamics and site index for 20 common European tree species ?

Focus on

- Temperature and precipitation changes
- Height growth process
- Even-aged pure stand

Modeling dominant height as a function of annual climate, per species

- Large climatic gradient, large species number, large set of control variables



→ French National Forest Inventory (NFI)


From Joly et al., 2010

Modeling dominant height as a function of annual climate, per species

- Large climatic gradient, large species number, large set of control variables
- Integration of an annual increment equation


$$\text{Dominant height increment}_t = f(\text{dominant height}_t, \text{envi}_t)$$

Not observed



Modeling dominant height as a function of annual climate, per species


- Large climatic gradient, large species number, large set of control variables
- Integration of an annual increment equation



Modeling dominant height as a function of annual climate, per species


- Large climatic gradient, large species number, large set of control variables
- Integration of an annual increment equation
- Climatic data with high temporal depth and precise spatial resolution

→ FYRE database (and Safran)




Simulations for all stands

- Impact of each climate variable: simulations of site index for different values of the variable






Simulations for all stands

- Impact of each climate variable: simulations of site index for different values of the variable




- Impact of CC: comparison of site index simulated under a reference climate (average 1891-1920) and actual climate (1950-2020)




Response to each climate variables

Temperature




→ Positive impact until saturation

Precipitation




→ Positive impact until saturation

Water balance



→ Positive impact until saturation

Impact of CC on site index




3 species show strong positive impact

3 species show negative impact


Within-species variations

Climate change impact on site index

Fagus sylvatica



Picea abies



+ Positive impact on cold stands
- Negative impact on warm stands

Stand average temperature 1891-1920

Combaud et al., 2024, FEM

- Method able to study CC long term impact on height increment ...
... even without height increment data !
- Positive impact of CC so far ...
... that could turn into negative in the future
- Strong variabilities in the between and within species response to CC...
... that should be taken into account to design adaptation strategies

References

- Albert, M., and M. Schmidt. 2010. "Climate-Sensitive Modelling of Site-Productivity Relationships for Norway Spruce (*Picea Abies* (L.) Karst.) and Common Beech (*Fagus Sylvatica* L.)." *Forest Ecology and Management, Adaptation of Forests and Forest Management to Changing Climate*, 259 (4): 739–49. <https://doi.org/10.1016/j.foreco.2009.04.039>.
- Assmann, Ernst, and P. W. Davis. 1970. *Principles of Forest Yield Study*. Pergamon.
- Bontemps, Jean-Daniel, Jean-Christophe Hervé, and Jean-François Dhôte. 2009. "Long-Term Changes in Forest Productivity: A Consistent Assessment in Even-Aged Stands." *Forest Science* 55 (6): 549–64. <https://doi.org/10.1093/forestscience/55.6.549>.
- Bontemps, Jean-Daniel, Jean-Christophe Herve, Pierre Duplat, and Jean-François Dhôte. 2012. "Shifts in the Height-Related Competitiveness of Tree Species Following Recent Climate Warming and Implications for Tree Community Composition: The Case of Common Beech and Sessile Oak as Predominant Broadleaved Species in Europe." *Oikos* 121 (8): 1287–99. <https://doi.org/10.1111/j.1600-0706.2011.20080.x>.
- Brandl, S., T. Mette, W. Falk, Patrick Vallet, T. Rötzer, and H. Pretzsch. 2018. "Static Site Indices from Different National Forest Inventories: Harmonization and Prediction from Site Conditions." *Annals of Forest Science* 75 (2). <https://doi.org/10.1007/s13595-018-0737-3>.
- Devers, Alexandre, Jean-Philippe Vidal, Claire Lauvernet, and Olivier Vannier. 2021. "FYRE Climate: A High-Resolution Reanalysis of Daily Precipitation and Temperature in France from 1871 to 2012." *Climate of the Past* 17 (5): 1857–79. <https://doi.org/10.5194/cp-17-1857-2021>.
- IGN. 2022. "Données Brutes de l'inventaire Forestier Mises En Ligne Sur DataIFN - Version 2.0." <https://inventaire-forestier.ign.fr/dataIFN/>.
- Lindner, Marcus, Michael Maroschek, Sigrid Netherer, Antoine Kremer, Anna Barbati, Jordi Garcia-Gonzalo, Rupert Seidl, et al. 2010. "Climate Change Impacts, Adaptive Capacity, and Vulnerability of European Forest Ecosystems." *Forest Ecology and Management, Adaptation of Forests and Forest Management to Changing Climate*, 259 (4): 698–709. <https://doi.org/10.1016/j.foreco.2009.09.023>
- Skovsgaard, J. P., and J. K. Vanclay. 2008. "Forest Site Productivity: A Review of the Evolution of Dendrometric Concepts for Even-Aged Stands." *Forestry: An International Journal of Forest Research* 81 (1): 13–31. <https://doi.org/10.1093/forestry/cpm041>.
- Zeide, Boris. 1993. "Analysis of Growth Equations." *Forest Science* 39 (3): 594–616. <https://doi.org/10.1093/forestscience/39.3.594>.

Thanks to

ONF

CLIMAE Metaprogram

IGN-IFN

Jean-Claude Gégout

Antoine Devers & Jean-Philippe Vidal

Björn Reineking

Jordan Bello

Mathieu Jonard

Xavier Morin

François Morneau

Nathéo Beauchamp

Anne Baranger

Maxime Jaunâtre


Carine Babusiaux

Compléments


Hypotheses

- Higher temperature, precipitation and water balance during spring and summer favor site index, but these positive effects saturate
- Climate change has had a different effect depending on the species and, for a given species, depending on the stand context

IFN measurement strategy




IFN measurement strategy




Dendrometric data			SDH (m)		Age (year)				Stand establishment date	
	Species	Number of stands	Mean	s.d	Mean	s.d.	Min	Max	Min	Max
	Abies alba Mill., 1768	817	25.5	6.2	72.7	33.6	9	147	1871	2002
	Betula pendula Roth, 1788	106	16.0	4.5	31.2	16.5	4	83	1932	2009
	Carpinus betulus L., 1753	97	19.5	4.9	63.4	25.5	11	125	1885	2003
	Castanea sativa Mill., 1768	440	17.7	4.8	43.9	26.1	5	141	1871	2013
	Fagus sylvatica L., 1753	1,470	23.4	7.2	83.1	35.0	8	147	1871	2006
	Fraxinus excelsior L., 1753	334	21.6	6.9	50.9	28.2	6	131	1875	2011
	Larix decidua subsp. decidua Mill., 1768	151	19.0	6.1	67.5	36.0	7	137	1873	2009
	Picea abies subsp. abies (L.) H.Karst., 1881	1,338	22.9	6.6	42.7	22.5	7	145	1871	2011
	Picea sitchensis (Bong.) Carrière, 1855	143	22.3	6.9	31.0	10.3	5	63	1944	2013
	Pinus halepensis Mill., 1768	344	12.5	3.9	50.1	23.6	6	137	1875	2010
	Pinus nigra subsp. nigra J.F.Arnold, 1785	413	15.0	5.3	53.2	29.3	6	135	1881	2012
	Pinus nigra var. corsicana (Loudon) Hyl., 1913	526	16.0	5.9	30.6	20.3	4	146	1871	2015
	Pinus pinaster subsp. pinaster Aiton, 1789	2,424	16.8	6.5	30.3	20.8	2	132	1876	2018
	Pinus sylvestris L., 1753	1,551	15.8	5.9	59.3	27.6	5	144	1873	2011
	Pseudotsuga menziesii (Mirb.) Franco, 1950	1,449	24.1	8.6	30.6	13.0	4	110	1910	2015
	Quercus petraea subsp. petraea (Matt.) Liebl., 1784	2,472	22.2	6.1	80.6	33.4	7	149	1871	2010
	Quercus pubescens Willd., 1805	1,019	13.9	4.3	67.6	25.0	7	145	1871	2006
	Quercus robur var. robur L., 1753	2,105	20.7	5.4	70.3	30.9	8	149	1871	2006
BES 2023 – 13 th December 2023	Quercus rubra L., 1753	114	18.5	6.2	26.4	16.3	4	81	1930	2006
Climate change altered the dynamics of st	Robinia pseudoacacia L., 1753	149	18.0	5.1	32.3	18.0	5	94	1916	2014

Modeling strategy



Model selection and parameter estimation

$$DH^{obs} = 1,3 + \sum \Delta DH_t + \epsilon$$



$$f_{exp}(X_t^{exp}) = A_0 \cdot \frac{\exp(\alpha \cdot X_t^{exp})}{1 + \exp(\alpha \cdot X_t^{exp})} \text{ (eq. 1)}$$


$$f_{decline}(X_t^{decl}) = C_0 \cdot \frac{\exp(\gamma \cdot X_t^{decl})}{1 + \exp(\gamma \cdot X_t^{decl})} \text{ (eq. 2)}$$

The final increment equation is equation (3). In this equation, SDH_t is SDH at the beginning of year t and ΔSDH_t is the SDH variation between the beginning of years t and $t + 1$.


$$\Delta SDH_t = \max[0, f_{exp}(X_t^{exp}) \cdot SDH_t^{\beta_0} - f_{decl}(X_t^{decl}) \cdot SDH_t] \text{ (eq. 3)}$$

$$SDH_i^{obs} = 1.3 + \sum_{t=t_i^{ori}}^{t_i^{obs}-1} \Delta SDH_{t,i} + \epsilon_i \text{ (eq. 4)}$$

Reference climate (average 1891- 1920)




Climate anomaly compared to reference climate (average 1891-1920)



Climate anomaly ■ positive ■ negative


Climate anomaly ■ positive ■ negative

Climate evolution (1891-1920 vs 1991-2020)




Simulations for all stands


- Impact of each climate variable: simulations of site index for different values of the variable



- Impact of CC: comparison of site index simulated under a reference climate (average 1891-1920) and actual climate (1950-2020)




Partial impact



- Abies alba (Ab.al) ● Betula pendula (Be.pe) ● Carpinus betulus (Ca.be) ● Castanea sativa (Ca.sa) ● Fagus sylvatica (Fa.sy) ● Fraxinus excelsior (Fr.ex)
- Pinus halepensis (Pi.ha) ● Pinus sylvestris (Pi.sy) ● Pseudotsuga menziesii (Ps.me) ● Quercus pubescens (Qu.pu) ● Quercus rubra (Qu.ru) ● Robinia pseudoacacia (Ro.ps)
- Larix decidua (La.de) ● Picea abies (Pi.ab) ● Pinus pinaster (Pi.pi) ● Quercus petraea (Qu.pe) ● Quercus robur (Qu.ro) ● Pinus nigra var. corsicana (Pi.ni.co)


Response to temperature and to precipitation

Temperature



→ Positive impact until saturation

Precipitation




→ Positive impact until saturation


Impact of CC on dominant height dynamics

Dominant height
Abies alba (m)

French temperature
anomalies (°C)



Within-species impact



Quality metrics

species	RMSE (m)	RMSPE (%)	bias (m)	optimism (%)
Abies alba	3.6	22.0	0.0	3.8
Betula pendula	2.5	21.0	0.0	34.7
Carpinus betulus	2.9	18.9	0.0	18.7
Castanea sativa	3.1	20.0	0.0	7.0
Fagus sylvatica	3.8	21.2	0.0	2.9
Fraxinus excelsior	3.7	21.3	0.0	6.5
Larix decidua	3.1	21.0	0.0	25.6
Picea abies	2.9	15.0	0.0	5.9
Picea sitchensis	3.5	23.5	0.0	9.3
Pinus halepensis	2.6	24.5	0.0	6.8
Pinus nigra subsp. nigra	3.2	28.5	0.0	9.6
Pinus nigra var. corsicana	2.4	17.4	0.0	7.1
Pinus pinaster	2.5	17.8	0.0	3.8
Pinus sylvestris	3.2	29.9	0.0	3.1
Pseudotsuga menziesii	2.9	14.3	0.0	3.6
Quercus petraea	3.2	17.6	0.0	2.1
Quercus pubescens	2.9	26.7	0.0	3.1
Quercus robur	3.4	19.8	0.0	2.9
Quercus rubra	3.0	18.3	-0.1	22.6
Robinia pseudoacacia	3.4	21.7	0.0	13.5