



**HAL**  
open science

## Designing innovative spatial strategies to control black leaf streak disease of banana through modelling approach

Catherine Abadie, Eduardo Corrales, Béranger Decouture, Marta Zaffaroni, Jean-François Rey, José Guzmán, Edgar Valverde, Nancy Chaves, Loup Rimbaud

► **To cite this version:**

Catherine Abadie, Eduardo Corrales, Béranger Decouture, Marta Zaffaroni, Jean-François Rey, et al.. Designing innovative spatial strategies to control black leaf streak disease of banana through modelling approach. XXIV Congreso Internacional Acobat, Apr 2024, Mexico, Mexico. hal-04608136

**HAL Id: hal-04608136**

**<https://hal.inrae.fr/hal-04608136>**

Submitted on 11 Jun 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



## DESIGNING INNOVATIVE SPATIAL STRATEGIES TO CONTROL BLACK LEAF STREAK DISEASE OF BANANA THROUGH MODELLING APPROACH


Catherine Abadie<sup>1,2,5</sup>, Eduardo Corrales<sup>2</sup>, Béranger Decouture<sup>1,2</sup>, Marta Zaffaroni<sup>3</sup>, Jean-François Rey<sup>3</sup>, José Guzmán<sup>4</sup>, Edgar Valverde<sup>4</sup>, Nancy Chaves<sup>4</sup>, Loup Rimbaud<sup>3</sup>. [catherine.abadie@cirad.fr](mailto:catherine.abadie@cirad.fr)

<sup>1</sup>CIRAD-PHIM, Costa Rica; <sup>2</sup>CATIE, Turrialba, Costa-Rica; <sup>3</sup>INRAE, Avignon, France; <sup>4</sup>CORBANA, Costa-Rica; <sup>5</sup>PHIM, University of Montpellier, CIRAD, INRAE, Institut Agro, IRD, F-34398 Montpellier, France

**Objectives.** As experimentations are not possible at large scale, the aim of the study is to assess the efficacy of innovative strategies to reduce fungicide use to manage Black Leaf Streak Disease (BLSD) of banana by using a modeling approach. Two different spatio-temporal strategies are evaluated on a real production landscape : (i) one directly based on fungicide strategies ; (ii) one based on the cultivation of resistant cultivars

### Methods


The generic epidemiological spatially-explicit model *landsepi* (Rimbaud *et al.*, 2018) was adapted to simulate BLSD by modifying plant growth to banana and 30 BLSD parameters. 52 spatial-temporal strategies were defined (Figure 1) to **reduce the index of fungicide treatments**: (1) with a **direct fungicide reduction** by reducing application frequency (5,10,15,20 days), the dose (25,50,75, 100%) or the spatial coverage (in line, at 25, 50, 75, 100%) representing 160 combinations of scenarios ; (2) by replacing the susceptible commercial variety by **resistant varieties** in 3 spatial allocations (line, mosaic or mixture) and modifying the 4 resistance genes efficacies representing 288 combinations. Each combinations was simulated 30 times **on a real 347-ha banana production basin** (Finca San Pablo in Costa-Rica, Figure 2). The exported yield and BLSD severity (Area Under Disease Curve Progress) provided by the simulated scenarios during 10 years will be compared.




**Fig. 1. 488 combinations of scenarios** to reduce fungicides (i) by reducing fungicide coverage, application frequency and fungicide doses, (ii) by using resistant varieties in line, mosaic or in mixture with susceptible variety. White and orange polygons represent non-treated and treated plots (with fungicide application or using cultivar resistance).

### Results. The model simulates realistic curves.

The parametrization of the *landsepi* model for BLSD allowed to obtain realistic curves of dynamics of the 4 compartments (Healthy, Latent, Infectious, Removed) with BLSD severity of 0,38 and an exported yield of 0,2 t.ha<sup>-1</sup>.season<sup>-1</sup> for 100% susceptible variety without any fungicide (Fig. 3)



**Fig 3: BLSD dynamics in 347 ha of susceptible banana without control during 5 years.**



**Fig. 2: Aerial photography of the Finca San Pablo with 295 plots (347ha), Costa-Rica**

**Perspectives.** The **13440 simulations outputs** are statistically analysed to allow to identify at 350 ha scale the **best scenarios of direct and indirect fungicide reduction** to manage BLSD while sustaining acceptable yields.