



HAL
open science

A new machine-learning model to partition soil organic carbon into its centennially stable and active fractions based on Rock-Eval(r) thermal analysis

Marija Stojanova, Pierre Barré, Hugues Clivot, Lauric Cécillon, François Baudin, Thomas Kätterer, Bent T. Christensen, Claire Chenu, Ines Merbach, Adrián Andriulo, et al.

► To cite this version:

Marija Stojanova, Pierre Barré, Hugues Clivot, Lauric Cécillon, François Baudin, et al.. A new machine-learning model to partition soil organic carbon into its centennially stable and active fractions based on Rock-Eval(r) thermal analysis. EGU General Assembly 2024, Apr 2024, Vienne, Austria. p. EGU24-11107, 2024, 10.5194/egusphere-egu24-11107 . hal-04612125

HAL Id: hal-04612125

<https://hal.inrae.fr/hal-04612125>

Submitted on 17 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

PARTYsoc v3: A new machine-learning model to partition soil organic carbon into its centennially stable and active fractions based on Rock-Eval® thermal analysis

Marija Stojanova [◇] Pierre Barré [◇] Hugues Clivot [♠] Adrian Andriulo [#] François Baudin [♣] Lauric Cécillon [▽] Claire Chenu [♡] Bent T Christensen [△] Sabine Houot [♡] Thomas Kätterer [○] Ines Merbach [▽] Fabien Ferchaud [†]

[◇]Laboratoire de Géologie, École Normale Supérieure, CNRS, PSL University, IPSL, Paris, France [♠]Université de Reims Champagne-Ardenne, INRAE, FARE, UMR A 614, Reims, France [#] Instituto Nacional de Tecnología Agropecuaria, Buenos Aires [♣]ISTeP, UMR 7193 Sorbonne Université, CNRS, Paris [▽]Research and innovation office, Ministry of agriculture and food sovereignty, Paris, France [♡]UMR ECOSYS, INRAE, AgroParisTech, Université Paris Saclay, Palaiseau, France [△]Aarhus University, Department of Agroecology - Soil Fertility [○]Swedish University of Agricultural Sciences [▽]Helmholtz-Centre for Environmental Research [†]UMR Eco&Sols, Univ Montpellier, CIRAD, INRAE, IRD, InstitutAgro Montpellier, Montpellier


Introduction

- Accurately simulating soil organic carbon (SOC) stock evolution is crucial for assessing the role of soil in climate regulation.
- SOC evolution is often modeled by dividing SOC into kinetic pools with contrasting residence times. **Initializing such kinetic pools remains a complex task** and a source of uncertainty in SOC simulations.
- In a previous study, Cécillon et al. [1] developed a machine-learning model (PARTYsoc v2) that uses Rock-Eval® thermal analysis results as input features to quantify the proportion of centennially stable and active SOC fractions.
- These proportions have been shown to be particularly effective for initializing the AMG model, enabling very accurate simulations of SOC stock evolutions for a dozen French agricultural long-term experiments [2].
- This present work builds a **new version of PARTYsoc**, validated on a larger data set, and extends the usefulness of the AMG model initialized with PARTYsoc to different parts of the world.

Materials and methods

Data


- Data collection of 18 sites with known crop yields and SOC stock evolutions and archived soil samples available for Rock-Eval® characterization.
- Sites are located in Europe and South America.
- The list of Rock-Eval® parameters [3] represents our potential **set of features**.
- The stable SOC stock size for each site is determined by optimization of the AMG model as in Clivot et al. (2019) [4].




- Using the optimal stable SOC stock values, we calculate the stable SOC proportion of each sample from each site, representing our **target variable**.
- We reserve 1/3 of the data for testing, and use 2/3 for training.

Feature engineering and model training


- We remove the highly correlated Rock-Eval® features using Spearman's correlation and a threshold of 0.9.
- We use AutoFeatureCorrelation and permutation importance to **rank the features according to their impact on the predicted target**.
- We run 30 repetitions of 5-fold cross-validation to find the **optimal number of features** for prediction on the test set and in Leave-One-Site-Out (LOSO) validation.
- The final model is a **combination of a Support Vector Machine (SVM) and a Beta regression**, trained using 10 Rock-Eval® features.



Results



- On average, we obtain an $R^2 = 0.78$ and $RMSE = 0.07$ in internal validation, and $R^2 = 0.59$ and $RMSE = 0.09$ in LOSO validation.



- Using PARTYsoc v3 (LOSO) values to initialize AMG simulations offers comparable results to initializing with optimal AMG values.

Conclusions and forthcoming research

- The proposed new PARTYsoc v3 builds upon and improves the original PARTYsoc v2 by incorporating a larger data set covering a **wider geographic area**, and a **more parsimonious machine-learning model** (10 vs. 18 Rock-Eval® features in PARTYsoc v2).
- We are currently working on extending the data set as well as stabilizing the processes of feature selection and model parameterization.

References

- L. Cécillon, F. Baudin, C. Chenu, B. T. Christensen, U. Franko, S. Houot, E. Kanari, T. Kätterer, I. Merbach, F. Van Oort, et al., "Partitioning soil organic carbon into its centennially stable and active fractions with machine-learning models based on Rock-Eval (R) thermal analysis," *Geoscientific Model Development*, 2021.
- E. Kanari, L. Cécillon, F. Baudin, H. Clivot, F. Ferchaud, S. Houot, F. Levasseur, B. Mary, L. Soucémarianadin, C. Chenu, et al., "A robust initialization method for accurate soil organic carbon simulations," *Biogeosciences*, vol. 19, no. 2, pp. 375–387, 2022.
- L. Pacini, T. Adatte, P. Barré, M. Boussafir, N. Bouton, L. Cécillon, V. Lamoureux-Var, D. Sebag, E. Verrecchia, A. Watrignon, et al., "Reproducibility of Rock-Eval® thermal analysis for soil organic matter characterization," *Organic Geochemistry*, vol. 186, p. 104687, 2023.
- H. Clivot, J.-C. Mouny, A. Duparque, J.-L. Dinh, P. Denoroy, S. Houot, F. Vertès, R. Trochard, A. Bouthier, S. Sagot, et al., "Modeling soil organic carbon evolution in long-term arable experiments with AMG model," *Environmental modelling & software*, 2019.

Acknowledgements

We thank Florence Savignac for her help in conducting Rock-Eval measurements. This work would not have been possible without all the people who helped initiate and maintain these invaluable long-term trials. The analyses have been funded by the CNRS-INRAE "IMMORTAL" project.