



HAL
open science

Adding to the diversity of a latest Devonian gondwanan plant assemblage: a new anatomically preserved axis from Barraba, Australia

Brigitte Meyer-Berthaud, Antoine Champreux, Anne-Laure Decombeix

► To cite this version:

Brigitte Meyer-Berthaud, Antoine Champreux, Anne-Laure Decombeix. Adding to the diversity of a latest Devonian gondwanan plant assemblage: a new anatomically preserved axis from Barraba, Australia. XI International Organisation Palaeobotany Conference, May 2024, Prague (CZ), Czech Republic. hal-04612426

HAL Id: hal-04612426

<https://hal.inrae.fr/hal-04612426>

Submitted on 14 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

XIth International Organization of Palaeobotany Congress, Prague, Czech Rep., May-27-June 1 2024

Adding to the diversity of a latest Devonian gondwanan plant assemblage : a new anatomically preserved axis from Barraba, Australia

Brigitte Meyer Berthaud¹, Antoine Champreux², Anne-Laure Decombeix¹

¹*CNRS, AMAP - Botany and modeling of plant architecture and vegetation, Montpellier, France*

²*Flinders University, ARC Centre of Excellence for Australian Biodiversity and Heritage, Adelaide, Australia*

Documenting a wide paleogeographic range of Famennian (latest Devonian) floras is essential to better understand the state of land plant diversity between the Kellwasser and the Hangenberg bio-events, two first-order global crises for the marine biota but whose impact on terrestrial organisms needs to be clarified.

There are few well-documented plant assemblages of Famennian age in Gondwana. One of them occurs in the Mandowa Mudstone Formation at Barraba, in New South Wales (Australia). The plant remains recovered at Barraba are dated from the late to latest Famennian and are preserved both as adpressions and permineralizations. A recent assessment of the permineralized assemblage indicates that the spore-producing plants are dominant and include lycopsids, cladoxylopsids, and one iridopterid (*Keraphyton*).

This presentation describes a second anatomically preserved iridopteridalean plant, represented by a 50 mm long and 7.7 mm wide fragment of axis with no branch or appendage bases attached along its length. Its tissues are entirely primary. The vascular system is actinostelic. In transverse section, it shows three fundamental ribs that divide twice slightly unequally. The protoxylem strands occur at the rib tips. Metaxylem tracheids are scalariform. The cortex is aerenchymatous in its innermost part.

This new plant is the first iridopterid to show an adaptation to a wetland habitat. Together with *Keraphyton*, it confirms that a group of Iridopteridales characterized by a complex vascular system occurred beyond the Frasnian in Eastern Gondwana. This discovery is consistent with a limited impact of the Kellwasser bio-event on land plant diversity prior to the Devonian-Carboniferous boundary.