



Evolution under climate change of the resilience of the services provided by the cultivated areas of the Pays de Fougères

Anne-Isabelle Graux, Patrick Chabrier, Eric Casellas, Klervi Le-Floch, Patrice Lecharpentier, Renan Le Roux, Fabien Ferchaud

► To cite this version:

Anne-Isabelle Graux, Patrick Chabrier, Eric Casellas, Klervi Le-Floch, Patrice Lecharpentier, et al.. Evolution under climate change of the resilience of the services provided by the cultivated areas of the Pays de Fougères. XIII STICS Seminar, Nov 2023, Latresne, France. hal-04637949

HAL Id: hal-04637949

<https://hal.inrae.fr/hal-04637949v1>

Submitted on 8 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License


➤ Evolution under climate change of the resilience of the services provided by the cultivated areas of the Pays de Fougères

REDELAC (2023 -2024) - Resilience And Sustainability Of Lowland Dairy Farms To Climate Hazards

Graux Anne-Isabelle, Patrick Chabrier, Eric Casellas, Klervi Le-Floch, Patrice Lecharpentier, Renan Le-Roux, Fabien Ferchaud


INRAE

INRAE, UMR PEGASE, anne-isabelle.graux@inrae.fr



➤ Background


Climate change
(climate hazards)



Dairy farming provides services



Food production, which depends
on **feed production**



Mitigate GHG emissions
by storing C

Impacts and adaptations depend on the forage system, local climatic conditions and seasons, as well as the public climate policies

➤ REDELAC objectives

- Study the **impact** of future climate and anticipate the **adaptive evolution** of dairy farms
- **Test a methodology** based on models/tools



Indicators



- **availability of feeds** and the evolution of **soil C stocks**
- evolution of **climate and feasibility** of crop/grassland **management** practices

AQAL-farm



- **adaptation of farms to feed resources**, and the consequences for **milk production** and **forage autonomy**

CAP'2ER[®]




AE

- **changes in the environmental footprint of farms** under climate change

> Questions

- How can **feed production** and **C storage services** evolve on the scale of the farm's cultivated area?
- What **differences** can be expected between
 - **farms** with **different forage systems**?
 - future **climates scenarios** and **time horizons**?
 - **locations** within a territory?
- Will dairy farms experience **more or less years** when the **herd's feed requirements** are **not met**? And where **C stock** is **moving away** from its trend?
- **What climatic conditions** explain these exceptional years in terms of feed production? And C stock change ?

➤ REDELAC's study is limited to



- A small (940 km²) area in Brittany: the Pays de Fougères
 - High **dairy farm density**, interest shown by **local stakeholders** in our questions
 - **Agricultural** territory, **soil and** (oceanic temperate) **climate favorable** to production
 - Drawn up a **territorial CAEP** => halve agricultural GHG emissions by 2050



➤ REDELAC's study is limited to

- 3 dairy farms representative of farms in Brittany (not real farms)
- ≠ forage systems




Farm	Description	 Corn in forage area (%)
cc1	Conventional, corn all year round	48
cc2	Conventional, corn silo closed 3 months	29
cc8	Organic, all-grass	0

- Well described by the “Chambre d’Agriculture de Bretagne”



➤ The land on each farm


- Cultivated in **1 or 2 rotations**, with a **few ha in perm. grassland**




 **CC1 farm**



 **CC2 farm**



 **CC8 farm**




- Most crops/grasslands are used to produce **feed for the herd**

- Corn  => corn silage 
- Grass  => Grazing  hay  or grass silage 
- Wheat  => sale, farm concentrate 
- Meslin  => farm concentrate 

➤ STICS simulations




- **Research version** derived from v10.0, which fixes some bugs
- **Improved parameterisation** of grasslands 
 - **BNF activation** to simulate **white clover** in ryegrass-clover associations

- **Resolution:**

- **Pedoclimatic units (PCU)**

= intersection of climate & soil resolution

grays polygons



- **30-year time horizons**
 - past — **reference : 1976-2005**
 - future
 - H1 : 2021-2050**
 - H2 : 2041-2070**
 - H3 : 2071-2100**

➤ STICS simulations


- Simulations of rotations and permanent grasslands for
 - **1 PCU** (soil with WHC=80mm <= geographic database of French soils + previous studies)
 - **1 climate scenario** <= DRIAS-2020 dataset
 - 1 global-regional climate model pair: **CNRM-CM5/ALADIN63**
 - 1 GHG emission scenario: **rcp8.5** (no climate regulation, +5°C by 2100)

WARNING

Preliminary
results


Each head of the rotation
⇒ Crop/grassland yield each year

2 uses of temporary grassland
⇒ grazing, hay/ silage



➤ Analysis of results

- Calculation of annual feed and soil C stocks at farm scale
 - Based on **areas** allocated to each rotation and **grassland/crop yield** or **soil C stock simulated** by STICS





- **Losses** from field to animal, excepted from grazed grass
- Other resilience indicators
 - **Theoretical herd feed requirements** based on herd size/composition and theoretical feed intake of heifers/cows
 - **“Deficit” years** = years when feed stocks from the year's production < feed requirements
 - **“Resistance”** = feed stocks from the year's production / feed requirements in deficit years

➤ Evolution of feed production service on the scale of the farm's cultivated area


- **Feed stocks** ↗ by **+15%** in h1, **+30%** in h2 and by **+40%** in h3
- **Fewer years** for which production does not meet the herd's DM feed requirements (↘ **deficit years** and **resistance** ↗)

Crop valorization




➤ Evolution of C storage service on the scale of the farm's cultivated area

- **Ref. period:** the **all-grass** organic farm **stores C**, unlike the other 2 farms whose C stocks are stable
- **Future:** trend towards **soil C depletion** or lower stock increase for the all-grass organic farm



➤ What explains these evolutions at farm scale ?

- ↗ in feed stocks




- **Slight** ↗ in annual **rainfall** (changes in distribution)
- ↗ in **temperature**
- ↗ in PET but **stable actual ET** due to CO₂ effect

Period	R (mm)	T (°C)	PET (mm)	CO ₂ (ppm)
ref	1015	10.8	805	354
h1	+15	+0.9	+78	+119
h2	+29	+1.7	+113	+224
h3	+55	+3.3	+200	+453

➤ What explains these evolutions at farm scale ?

- Trend towards lower soil C stocks



- ↗ **SOC mineralization** not fully offset by ↗ **C inputs** (with only a fraction stabilised in SOM)

cc1 

Period	Min. (t C/ha)	Inputs (t C/ha)
ref	3.5	4.0
h1	+ 0.4	+ 0.4
h2	+ 0.8	+ 0.9
h3	+ 1.3	+ 1.4

cc2 

Period	Min. (t C/ha)	Inputs (t C/ha)
ref	3.6	4.1
h1	+ 0.5	+ 0.5
h2	+ 0.9	+ 1.1
h3	+ 1.4	+ 1.6

cc8 

Period	Min. (t C/ha)	Inputs (t C/ha)
ref	4.5	5.4
h1	+ 0.5	+ 0.6
h2	+ 1.0	+ 1.2
h3	+ 1.7	+ 1.9

➤ Conclusions

Based on the simulated example (one PCU and one climate scenario), in the Pays de Fougères:


- **Climate still favorable** to production in the future
- **Increase in overall feed stocks** and better coverage of herd feed requirements
- **C destocking** or lower C stocking for all-grass organic farms
- **N₂O emissions slightly enhanced** by climate change

⇒ Possible **antagonism** between **food security/local consumption** issues and **C footprint reduction**?


⇒ Possible evolution of farms towards **greater proportion of grass** and **fewer concentrates** in the animal diet?

➤ Prospects

- Extension of STICS simulations to the entire plan
 - Simulation for **all PCU** in the territory and **all future climate scenarios**



➤ Prospects



- Additional analysis of results
 - **Resilience of soil C storage**
 - Changes in **forage quality**
 - **Accessibility of grass** for grazing (soil bearing capacity)
 - Animal **heat stress**
- Supply of information required for AQAL-farm model simulations
 - **Grass growth and accessibility**
 - Annual **feed stocks**
 - Forecast **feeding plan**, etc.

➤ Thank you for your attention



Photo : L. Delaby



INRAE

Anne-Isabelle Graux

15 November 2023