



HAL
open science

Light strategy regulation enhances single-cell protein production and nutrient recovery from wastewater by photosynthetic bacteria

Xiaodan Wang, Haifeng Lu, Guangming Zhang, Jean-Philippe Steyer, Gabriel Capson-Tojo

► **To cite this version:**

Xiaodan Wang, Haifeng Lu, Guangming Zhang, Jean-Philippe Steyer, Gabriel Capson-Tojo. Light strategy regulation enhances single-cell protein production and nutrient recovery from wastewater by photosynthetic bacteria. 1st International Conference on Novel Photorefineries for Resource Recovery, Sep 2024, Valladolid, Spain. hal-04644543

HAL Id: hal-04644543

<https://hal.inrae.fr/hal-04644543v1>

Submitted on 11 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Light strategy regulation enhances single-cell protein production and nutrient recovery from wastewater by photosynthetic bacteria

Xiaodan Wang^{1,2}, Haifeng Lu^{1*}, Guangming Zhang³, Jean-Philippe Steyer², Gabriel Capson-Tojo²

¹ College of Water Resources and Civil Engineering, China Agricultural University, Beijing 100083, China.

² INRAE, Univ Montpellier, LBE, 102 Avenue des Etangs, Narbonne 11100, France.

³ School of Energy and Environmental Engineering, Hebei University of Technology, Tianjin 300401, China.

*Corresponding author: haifenglu@cau.edu.cn

Keywords: light intensity; photoperiod; light spectrum; single cell protein; resource recovery

Introduction

The increasing production of wastewater and the lack of sufficient protein supply worldwide are two major challenges caused by rapid population growth and increasing environmental awareness (Li et al., 2023). The utilization of wastewater streams to produce single-cell protein (SCP) can be a promising method to solve the above problems simultaneously. Purple non-sulfur bacteria (PNSB) are a promising SCP mediator due to their unique metabolic features, mostly thanks to their high biomass yields (Capson-Tojo et al., 2020). In addition, PNSB can exhibit tolerance to high organic loads and utilize the organic constituents present in the wastewater for growth (Lu et al., 2019). Furthermore, in comparison to many other microbial SCP sources, PNSB biomass contains value-added biomolecules, such as carotenoids, bacteriochlorophyll, coenzyme Q10, and a broader scope of essential amino acids (Zhi et al., 2020), which improve the nutritional value of SCP. Light plays a critical role on PNSB growth, being the energy source during phototrophic metabolism. Light factors include light intensity, photoperiod, and light spectrum. Previous studies on the impact of light regulation on PNSB have primarily examined its effects on biomass growth and pigment production (Zhou et al., 2015; Kuo et al., 2012). It has been found that light regulation has a significant impact on PNSB metabolism, especially on carbon and nitrogen assimilation. SCP, as a major component of PNSB organisms, is susceptible to the influence of the light environment. However, comprehensive studies on light strategy regulation of SCP synthesis in wastewater systems with photosynthetic bacteria are currently lacking. This study investigates the effects of light intensity, photoperiod and light spectrum on biomass production, SCP synthesis, pollutant removal, nutrient transformation efficiency and the correlation between these factors in a PNSB wastewater resource recovery system.

Materials and Methods

Artificial sugar wastewater was used in the experiment, made from glucose (1.6 g/L), malic acid (1.6 g/L), sodium acetate (1.6 g/L), ammonium sulfate (1 g/L), potassium dihydrogen phosphate (0.2 g/L), and magnesium sulfate heptahydrate (0.1 g/L). The initial concentrations of chemical oxygen demand (COD) and ammonium nitrogen (NH₄⁺-N) were 4000 and 250 mg/L, respectively. The initial pH was adjusted to 7.0 using HCl (0.5 mol/L) and NaOH (0.5 mol/L) solution.

The inoculated PNSB was a strain of *Rhodospseudomonas palustris* (*R. palustris*) and was purchased from the China BeNa Culture Collection. The initial inoculum of *R. palustris* had a concentration of 360 mg/L (dry weight). All experiments were conducted on a 2.5 m × 1.2 m × 1.5 m (length × width × height) experimental stand. The setup included an illumination system (either incandescent lamps or LEDs), batch photobioreactors (1,000 mL high-silica glass bottles with a working volume of 600 mL) and a magnetic stirrer (operating at 120 rpm) for mixing. Blackout curtains were used to create light-tight enclosures between different treatment groups to prevent interference between light sources. Each group consisted of three replicates. Three different lighting factors were modified, namely light intensity, photoperiod and light spectrum. To control for a single variable, the light intensity was maintained at a constant level of 120 μmol/m²/s in the experiments involving photoperiod and light spectrum (see Table 1).

Table 1. Experimental design of light supply strategies

Light strategies	Control groups	Experimental groups
Light intensity (μmol/m ² /s)	0	40, 80, 120, 160, 200, 240

Photoperiod (L/D) *	24/0	3/21, 9/15, 12/12, 15/9, 18/6
Light wavelength (nm)	Incandescent lamp (380~780)	White light (380~780), Red light (557~711), Yellow light (542~665), Blue light (404~528), Green light (542~665)

* L corresponds to light (illuminated); D corresponds to dark (non-illuminated). The values indicate the number of hours for L and D

Results and discussion

The results indicated that, under conditions of 120 $\mu\text{mol}/\text{m}^2/\text{s}$ light intensity, L/D equal to 18/6, and incandescent light, *R. palustris* exhibited the highest biomass concentrations and daily bacterial production, reaching 1,140.5 \pm 19.7 mg/L and 0.32 \pm 0.02 g/L/d, respectively. This represents an increase of 17.06% ~ 93.21% and 54.43%~299.93% compared to the L/D of 24/0, 3/21 and 9/15, respectively ($P < 0.05$). Moreover, under 120 $\mu\text{mol}/\text{m}^2/\text{s}$ light intensity, L/D equal to 3/21, and incandescent light, *R. palustris* displayed the highest protein content, at 67.47%. This protein content was significantly greater, showing an increase of 21.96%~44.54% compared to all other experimental groups ($P < 0.05$). Additionally, COD and $\text{NH}_4^+\text{-N}$ removal efficiencies reached 72.03%~78.40% under these optimal conditions (L/D = 18/6), showing an increase of 20.26%~40.97% compared to other experimental groups (L/D = 3/21) ($P < 0.05$). Correlation analysis revealed significant negative associations between light intensity and light spectrum with protein content and concentration, respectively. Conversely, photoperiod exhibited a significant positive correlation with protein concentration (Fig. 1). Consequently, photoperiod is an effective regulatory strategy for enhancing the production of SCP by *R. palustris*. This research provides innovative methods and insights for enhancing SCP synthesis in wastewater systems using photosynthetic bacteria.




Fig. 1. Correlation analysis under three light supply strategies: (a) light intensity; (b) photoperiod; (c) light spectrum (* $p < 0.05$ ** $p < 0.01$ *** $p < 0.001$)

Acknowledgments

This research has received funding from the Project of Sanya Yazhou Bay Science and Technology City (No.SCKJ-JYRC-2022-34).

References

- Li R., Fan X., Jiang Y., Wang R., Guo R., Zhang Y., Fu S. 2023. From anaerobic digestion to single cell protein synthesis: A promising route beyond biogas utilization. *Water Res*, **243**, 120417.
- Capson-Tojo G., Batstone D.J., Grassino M., Vlaeminck S.E., Puyol D., Verstraete W., Kleerebezem R., Oehmen A., Ghimire A., Pikaar I., Lema J.M., Hulsen T. 2020. Purple phototrophic bacteria for resource recovery: Challenges and opportunities. *Biotechnol Adv*, **43**, 107567.
- Lu H., Zhang G., Zheng Z., Meng F., Du T., He S. 2019. Bio-conversion of photosynthetic bacteria from non-toxic wastewater to realize wastewater treatment and bioresource recovery: A review. *Bioresour Technol*, **278**, 383-399.
- Zhi R., Cao K., Zhang G., Zhu J., Xian G. 2020. Zero excess sludge wastewater treatment with value-added substances recovery using photosynthetic bacteria. *Journal of Cleaner Production*, **250**.
- Zhou Q., Zhang P., Zhang G., Peng M. Biomass and pigments production in photosynthetic bacteria wastewater treatment: Effects of photoperiod[J]. *Bioresour Technol*, 2015,**190**, 196-200.
- Kuo F., Chien Y., Chen C. Effects of light sources on growth and carotenoid content of photosynthetic bacteria *Rhodospseudomonas palustris*[J]. *Bioresour Technol*, 2012,**113**, 315-318.