



**HAL**  
open science

# Discovering how heat stress and water deficit, alone or in combination reconfigure interactions between plant and microbial communities

Corentin Maslard, Mustapha Arkoun, Christophe Salon, Fanny Leroy, Lun Jing, Jingjing Peng, Marion Prudent

## ► To cite this version:

Corentin Maslard, Mustapha Arkoun, Christophe Salon, Fanny Leroy, Lun Jing, et al.. Discovering how heat stress and water deficit, alone or in combination reconfigure interactions between plant and microbial communities. International Society of Root Research 12th International Symposium Roots [& Roads] to a sustainable future. 12ISRR 2024, Jun 2024, Leipzig, Germany. hal-04645313

**HAL Id: hal-04645313**

**<https://hal.inrae.fr/hal-04645313v1>**

Submitted on 11 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

## 2.4 A holistic view on root systems - integrating the data from roots, root biota, and soils

### Discovering how heat stress and water deficit, alone or in combination reconfigure interactions between plant and microbial communities

Corentin Maslard<sup>1</sup>; Mustapha Arkoun<sup>2</sup>; Christophe Salon<sup>1</sup>; Fanny Leroy<sup>3</sup>; Lun Jing<sup>2</sup>; Jingjing Peng<sup>4</sup>; Marion Prudent<sup>1</sup>

<sup>1</sup> INRAE; <sup>2</sup> TIMAC AGRO; <sup>3</sup> Normandie Université, Unicaen; <sup>4</sup> China Agricultural University (CAU)

In the context of climate change, more frequent drought events and heatwaves predict a significant decrease in soybean yields. As the world's most widely grown legume crop, there's an urgent need to enhance its resilience to ensure productivity. This study explores how heat and water stress, alone or combined, impact soybean nutrition and growth, including root morphology, nutrient uptake and efficiency, and the surrounding microbial environment. The aim was to identify if root architecture can confer resistance to these stresses and identify the structural and functional changes implemented by the plant during these stresses.

Two soybean genotypes with contrasted root architectures were grown in the 4PMI high-throughput phenotyping platform during their vegetative stage, under various climate conditions. Conditions included control, heatwaves, water deficit, and combined heatwaves and water deficit. A holistic approach was adopted, encompassing plant ecophysiological, metabolic, and transcriptional measurements and rhizospheric microbiome analysis. At transcriptomic level, the effect of stress was much greater than the effect of genotype. Moreover, combined stresses provoked a specific response that was not simply the sum of the effects of water deficit and heat stress. Among the many stress-induced deregulations, we observed a concomitant increase in the level of expression of root sulphur (S) transporters, and in the concentration of S in shoots. In case of combined stresses, arginine, and asparagine (osmoprotective amino acids) concentrations increased significantly in roots and exudates, as did the presence of actinobacteria in the root endosphere. We will discuss the close relationship between compounds produced by the plants in response to stress and the modulation of associated microbial communities in the rhizosphere.

This study highlights the need to consider plant interactions with micro-organisms to understand and enhance its resilience to stress.