



**HAL**  
open science

## Review in plant -“flavescence dorée” phytoplasma interactions

Sandrine Eveillard

► **To cite this version:**

Sandrine Eveillard. Review in plant -“flavescence dorée” phytoplasma interactions. European workshop on “Flavescence dorée” - Recent acquisitions and management strategies, Assunta Bertaccini, Nicola Mori, Elisa Angelini and Annalisa Polverari, Jan 2024, Verona, Italy, Italy. pp.11. hal-04650242

**HAL Id: hal-04650242**

**<https://hal.inrae.fr/hal-04650242v1>**

Submitted on 16 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Review in plant-“flavescence dorée” phytoplasma interactions

Sandrine Eveillard

INRAE, Université de Bordeaux, UMR BFP, 33882, Villenave d'Ornon, France

“Flavescence dorée” (FD) phytoplasma is responsible for an epidemic quarantine disease of grapevine in Europe and control methods are mandatory. Nevertheless, the disease still progress, and alternative control methods are needed implying a better knowledge of the plant-FD phytoplasma interactions. It is known that grapevine have various susceptibilities to “flavescence dorée” in terms of percentage of infected plants, symptoms severity and phytoplasma titers, with all the *Vitis vinifera* being susceptible although at different degrees. This was shown in vineyard and in controlled conditions, for example for Tocai friulano, Nebbiolo and Merlot (Ripamonti *et al.*, 2021; Eveillard *et al.*, 2016).

Omics studies were then undertaken to determine the mechanism of response of the plant to the FD phytoplasma infection. Highly and poorly susceptible varieties were studied to identify resistance or susceptibility factors. Indeed, transcriptomic changes were observed in infected plants for several genes in various pathways (Casarin *et al.*, 2023; Bodin *et al.*, 2023). Metabolomic changes were also observed between healthy and infected grapevine, but also between highly and poorly susceptible cultivars (Deborde *et al.*, 2023; Teixeira *et al.*, 2020, 2023; Davosir *et al.*, 2023). Moreover, these changes could be linked to genes overexpression or repression. In addition to poorly and highly susceptible cultivars, transcriptomic analysis were also done on recovered grapevine showing specificities (Pacífico *et al.*, 2018). In parallel to these researches concerning the plant response, effectors secreted by phytoplasmas have been shown to interfere with host plant physiology, presumably in favor of phytoplasma multiplication or dissemination. Most of the recent studies of phytoplasma effectors have focused on a few proteins that are not encoded in the “flavescence dorée” genome but one (Debonneville *et al.*, 2023). These researches aim to identify grapevine genes, metabolites or pathways associated to FD resistance or susceptibility that could be used as biomarkers or targets to inactivate or enhanced to help the plant fighting against the disease.

## Literature cited

- Bodin E., Jollard C., Dassé A., Dufour MC., Razan F., Desqué D., Corio-Costet MF, Malembic-Maher S and Eveillard S 2023. Insight into grapevine gene responses to “flavescence dorée” phytoplasma. *Phytopathogenic Mollicutes*, 13(1): 19-20.
- Casarin S, Vincenzi S, Esposito A, Filippin L, Forte V, Angelini E and Bertazzon N. 2023. A successful defence strategy in grapevine cultivar Tocai friulano provides compartmentation of grapevine “flavescence dorée” phytoplasma. *BMC Plant Biology*, 23: 161.
- Debonneville C, Mandelli L, Brodard J, Groux R, Roquis D and Schumpp O 2022. The complete genome of the “flavescence dorée” phytoplasma reveals characteristics of low genome plasticity. *Biology*, 11(7): 953.
- Deborde C, Bernillon S, Valls-Fonayet J, Jacob D, Malembic-Maher S, Desqué D, Lusseau T, Moing A and Eveillard S 2023. Towards a metabolomic characterization of the grapevine response to “flavescence dorée” infection by NMR and LC-MS profiling. *Phytopathogenic Mollicutes*, 13(1): 17-18.
- Davosir D, Šola I, Ludwig-Müller J and Šeruga Musić M. 2023. “Flavescence dorée” strain-specific impact on phenolic metabolism dynamics in grapevine (*Vitis vinifera*) throughout the development of phytoplasma infection. *Journal of Agricultural Food Chemistry*. <https://doi.org/10.1021/acs.jafc.3c06501>.
- Eveillard S, Jollard C, Labroussaa F, Khalil D, Perrin M, Desqué D, Salar P, Razan F, Hévin C, Bordenave L, Foissac X, Masson JE and Malembic-Maher S 2016. Contrasting susceptibilities to “flavescence dorée” in *Vitis vinifera*, rootstocks and wild *Vitis* species. *Frontiers in Plant Science*, 7: 1762
- Pacifico D, Margaria P, Galetto L, Legovich M, Abbà S, Veratti F, Marzachi C and Palmano S. 2018. Differential gene expression in two grapevine cultivars recovered from “flavescence dorée”. *Microbiological Research*, 220:72-82.
- Ripamonti M, Pegoraro M, Morabito C, Gribaudo I, Schubert A, Bosco D and Marzachi C 2021. Susceptibility to “flavescence dorée” of different *Vitis vinifera* genotypes from north-western Italy. *Plant Pathology*, 70(3): 511–520.
- Teixeira A, Martins V, Frusciantè S, Cruz T, Noronha H, Diretto G and Gerós H. 2020. “Flavescence dorée”-derived leaf yellowing in grapevine (*Vitis vinifera* L.) is associated to a general repression of isoprenoid biosynthetic pathways. *Frontiers in Plant Science*, 17(11): 896
- Teixeira A, Noronha H, Frusciantè S, Diretto G and Gerós H 2023. The grapevine metabolite profile of phloem sap is modified by “flavescence dorée”. *OENO One*, 57(1): 307-320.