



**HAL**  
open science

## Insights into physiological responses of different grapevine varieties to Flavescence Dorée infection: an integrated approach

Loreen Larroquet, Enora Bodin, Cédric Cassan, Camille Jollard, Marie-Cecile Dufour, Frédérique Razan, Yves Gibon, M.-F. Corio-Costet, Sylvie Malembic-Maher, Sandrine Eveillard

### ► To cite this version:

Loreen Larroquet, Enora Bodin, Cédric Cassan, Camille Jollard, Marie-Cecile Dufour, et al.. Insights into physiological responses of different grapevine varieties to Flavescence Dorée infection: an integrated approach. 6. European Bois Noir workshop and 1st International Pro-AECOLOGY conference, UMR-1332 Fruit Biology and Pathology, University of Bordeaux and INRAE, May 2024, Bordeaux (France), France. hal-04650385

**HAL Id: hal-04650385**

**<https://hal.inrae.fr/hal-04650385>**

Submitted on 16 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Insights into physiological responses of different grapevine varieties to Flavescence Dorée infection: an integrated approach

Loreen Larroquet<sup>1#</sup>, Enora Bodin<sup>1#</sup>, Cédric Cassan<sup>1</sup>, Camille Jollard<sup>1</sup>, Marie-Cécile Dufour<sup>2</sup>, Frédérique Razan<sup>1</sup>, Yves Gibon<sup>1</sup>, Marie-France Corio-Costet<sup>2</sup>, Sylvie Malembic-Maher<sup>1</sup>, Sandrine Eveillard<sup>1\*</sup>

# both authors contributed equally to this work

<sup>1</sup>University of Bordeaux, INRAE, BFP, Villenave d'Ornon, France;

<sup>2</sup>INRAE, SAVE, Villenave d'Ornon, France.

\*Corresponding author(s): [Sandrine.eveillard@inrae.fr](mailto:Sandrine.eveillard@inrae.fr)

### INTRODUCTION

Flavescence dorée (FD) is a severe and widespread quarantine disease that affects grapevines in various European viticultural regions (Boudon-Padieu *et al.*, 2002). It is caused by a phytoplasma (FDp), and grapevine varieties exhibit different levels of susceptibility to the disease. Cabernet Sauvignon (CS) is highly susceptible whilst Merlot (M) is less susceptible (Eveillard *et al.*, 2016). Previous studies have shown that phytoplasma infections lead to significant transcriptomic and metabolomic changes in grapevines (Bertazzon *et al.*, 2019; Dermastia, 2019; Margaria *et al.*, 2014; Prezelj *et al.*, 2016). The purpose of this study is to better decipher the differences of response to FDp infection of various grapevine cultivars, and identify biomarkers associated with reduced susceptibility. This was achieved under controlled conditions integrating transcriptomics, metabolic profiling and physiological studies.

### MATERIALS AND METHODS

For all experiments, grapevine plants were inoculated with FDp strain FDPEY-05 (Papura *et al.*, 2009) in a high-confinement greenhouse as described in Eveillard *et al.* (2016). FDp-infected or non-infected *Scaphoideus titanus* were placed for a one-week transmission on the fifth leaf from the apex. One week post-inoculation (wpi) and ten wpi, leaves were collected from each plant and immediately frozen in liquid nitrogen. Plant DNA extractions and phytoplasma absolute qPCR quantifications were performed as in Eveillard *et al.*, 2016. For transcriptomic analysis, RNA extraction and reverse transcription were performed as in Dufour *et al.* (2016) followed by a high-throughput qPCR method (Fluidigm ©) with the pathways-targeted NeoViGen and Biostim chips (Bodin *et al.*, 2020; Bodin *et al.*, 2023; Dufour *et al.*, 2016). For metabolic profiling, ethanolic extraction was performed on lyophilized samples. Soluble sugars, organic acids, free total amino acids, starch, proteins and cellular residues were quantified through enzymatic methods and via spectrophotometric/fluorescent assays. Physiological data, *i.e.* stomatal conductance and chlorophyll a fluorescence, have been acquired with the LI-COR 600 System in automatic mode. All statistical analyses were performed using R Studio (version 4.3.2.).

### RESULTS AND DISCUSSION

No phytoplasma was detected in the leaves of the grapevines that were exposed to non-infected insects. Phytoplasma titers at 1 wpi were non statistically different among the grape varieties tested. However, titers were significantly higher in CS than in M at 10 wpi.

Transcriptomic analysis revealed specific gene expressions for each plant condition. Indeed, principal component analysis (PCA) separated infected and healthy plants on the first dimension and varieties on the second dimension. For example, infected-M showed overexpression of genes involved in jasmonic acid pathways, while healthy-M modulated the genes related to salicylic acid and ethylene pathways.

Results using the LI-COR 600 system also suggested different responses depending on the

susceptibility of grapevine varieties. Indeed, in M and CS, chlorophyll fluorescence tended to increase  
Proceedings of the 6th European Bois Noir workshop and 1st International Pro-AECOLOGY conference  
14-16 may, 2024

109

in M but decreased in CS in infected leaves. The stomatal conductance significantly decreased in  
infected leaves at 10 wpi, as compared to leaves exposed to non-infected insects.

Finally, preliminary enzymatic assays indicate that, at 1 wpi, carbohydrate levels increased in infected  
CS as compared to healthy ones, while they decreased in infected M.

Altogether, these results are in line with studies showing significant transcriptomic differences  
between healthy and phytoplasma-infected plants, as well as between highly susceptible and poorly  
susceptible varieties (Bertazzon *et al.*, 2019; Margaria *et al.*, 2014). Additionally, the reduction in  
chlorophyll fluorescence in infected cultivars, directly linked to photosystem II activity, is in  
agreement with findings indicating a decrease in chlorophyll content (Teixeira *et al.*, 2020).

#### ACKNOWLEDGEMENTS

We thank our colleagues T. Lusseau, D. Lacaze and T. Mauduit for production of insect and plant  
material. We thank also J. Masson and M. Perrin for supplying *in vitro* plantlets. This research was  
funded by the ‘Conseil Interprofessionnel du Vin de Bordeaux’, ‘France Agrimer’, the Nouvelle  
Aquitaine region, ‘Plan National Dépérissement du Vignoble’ and “Grand Programme de Recherche  
– Bordeaux Plant Science” of the University of Bordeaux.

#### REFERENCES

- Bodin E., Bellée, A., Dufour, M. C., André, O., & Corio-Costet, M. F. (2020). Grapevine Stimulation: A  
Multidisciplinary Approach to Investigate the Effects of Biostimulants and a Plant Defense Stimulator.  
Journal  
of Agricultural and Food Chemistry, 68(51), 15085–15096. <https://doi.org/10.1021/acs.jafc.0c05849>
- Bodin, E., Jollard, C., Dasse, A., Dufour, M. C., Razan, F., Desqué, D., Corio-Costet, M. F., Malembic-Maher  
S, &  
Eveillard, S. (2023). Insights into grapevine gene responses to ‘flavescence  
dorée’ phytoplasma. *Phytopathogenic Mollicutes*, 13(1), 19-20.
- Boudon-Padiou E 2002. Flavescence dorée of the grapevine: knowledge and new developments in  
epidemiology, etiology  
and diagnosis. *ATTI Giornate Fitopatologica*, 1:15-34.
- Bertazzon N, Bagnaresi P, Forte V, Mazzucotelli E, Filippin L, Guerra D, Zechini A, Cattivelli L, Angelini E  
2019.  
Grapevine comparative early transcriptomic profiling suggests that Flavescence dorée phytoplasma represses  
plant responses induced by vector feeding in susceptible varieties. *BMC Genomics*, 20(1):526.
- Dermastia M 2019. Plant Hormones in Phytoplasma Infected Plants. *Frontiers in Plant Science*, 10:477.
- Dufour M.-C., Magnin, N., Dumas, B., Vergnes, S., & Corio-Costet, M.-F. (2016). High-throughput gene-  
expression  
quantification of grapevine defense responses in the field using microfluidic dynamic arrays. *BMC Genomics*,  
17(1), 957. <https://doi.org/10.1186/s12864-016-3304>.
- Eveillard S, Jollard C, Labroussaa F, Khalil D, Perrin M, Desqué D, Salar P, Razan F, Hévin C, Bordenave L,  
Foissac X,  
Masson JE, Malembic-Maher S 2016. Contrasting Susceptibilities to Flavescence Dorée in *Vitis vinifera*,  
Rootstocks and Wild Vitis Species. *Frontiers in Plant Science*, 7:1762.
- Margaria P, Ferrandino A, Caciagli P, Kedrina O, Schubert A, Palmano S 2014. Metabolic and transcript  
analysis of the  
flavonoid pathway in diseased and recovered Nebbiolo and Barbera grapevines (*Vitis vinifera* L.) following  
infection by Flavescence dorée phytoplasma. *Plant Cell and Environment*, 37:2183-200.
- Papura D, Delmotte F, Giresse X, Salar P, Danet J-L, Van Helden M, Foissac X, Malembic-Maher S. (2009).  
Comparing  
the spatial genetic structures of the Flavescence dorée phytoplasma and its leafhopper vector *Scaphoideus*

*titanus*.

*Infection, Genetics and Evolution*, 9 (5), 867-876.

Prezelj N, Covington E, Roitsch T, Gruden K, Fragner L, Weckwerth W, Chersicola M, Vodopivec M, Dermastia M 2016.

Metabolic Consequences of Infection of Grapevine (*Vitis vinifera* L.) cv. “Modra frankinja” with Flavescence Dorée Phytoplasma. *Frontiers in Plant Science*, 7 :711

Teixeira A., Martins V., Frusciante S., Cruz T., Noronha H., Diretto G., Gerós H. (2020). Flavescence Dorée-Derived Leaf

Yellowing in Grapevine (*Vitis vinifera* L.) Is Associated to a General Repression of Isoprenoid Biosynthetic Pathways. *Frontiers in Plant Science*, 17;11: 896. doi: 10.3389/fpls.2020.00896. PMID: 32625230; PMCID: PMC7311760.