



HAL
open science

Undetection of vector-borne viruses in equids of Galapagos Islands

Gina Zanella, Cécile Beck, José-Carlos Vallé-Casuso, Madeline Anthony, Marilyn Cruz, Alberto Vélez, Rommel Lenin Vinueza, Gaëlle Gonzalez

► **To cite this version:**

Gina Zanella, Cécile Beck, José-Carlos Vallé-Casuso, Madeline Anthony, Marilyn Cruz, et al.. Undetection of vector-borne viruses in equids of Galapagos Islands. *Frontiers in Veterinary Science*, 2024, 11, 10.3389/fvets.2024.1411624 . hal-04654492

HAL Id: hal-04654492

<https://hal.inrae.fr/hal-04654492>

Submitted on 19 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



OPEN ACCESS

EDITED BY

Heinzpeter Schwermer,
Federal Food Safety and Veterinary Office
(FSVO), Switzerland

REVIEWED BY

Serafeim Christos Chaintoutis,
Aristotle University of Thessaloniki, Greece
Zhaoyang Tang,
Huzhou University, China

*CORRESPONDENCE

Gina Zanella
✉ gina.zanella@anses.fr

RECEIVED 03 April 2024

ACCEPTED 29 May 2024

PUBLISHED 07 June 2024

CITATION

Zanella G, Beck C, Valle-Casuso J-C,
Anthony M, Cruz M, Vélez A,
Vinueza RL and Gonzalez G (2024)
Undetection of vector-borne viruses in equids
of Galapagos Islands.
Front. Vet. Sci. 11:1411624.
doi: 10.3389/fvets.2024.1411624

COPYRIGHT

© 2024 Zanella, Beck, Valle-Casuso, Anthony,
Cruz, Vélez, Vinueza and Gonzalez. This is an
open-access article distributed under the
terms of the [Creative Commons Attribution
License \(CC BY\)](#). The use, distribution or
reproduction in other forums is permitted,
provided the original author(s) and the
copyright owner(s) are credited and that the
original publication in this journal is cited, in
accordance with accepted academic
practice. No use, distribution or reproduction
is permitted which does not comply with
these terms.

Undetection of vector-borne viruses in equids of Galapagos Islands

Gina Zanella^{1*}, Cécile Beck², José-Carlos Valle-Casuso^{3,4},
Madeline Anthony^{3,4}, Marilyn Cruz⁵, Alberto Vélez⁵,
Rommel Lenin Vinueza⁶ and Gaëlle Gonzalez²

¹Epidemiology Unit (EPIMIM), Laboratoire de Santé Animale, ANSES, Ecole Nationale Vétérinaire d'Alfort, Maisons-Alfort, France, ²ANSES, INRAE, Ecole Nationale Vétérinaire d'Alfort, UMR 1161 Virologie, Laboratoire de Santé Animale, Maisons-Alfort, France, ³ANSES, Laboratory for Animal Health in Normandy, Physiopathology and Epidemiology of Equine Diseases Unit, Goustranville, France, ⁴Mixed Technological Unit "Equine Health and Welfare - Organisation and Traceability of the Equine Industry" (UMT SABOT), Normandie, France, ⁵Agencia de Regulación y Control de la Bioseguridad y Cuarentena para Galápagos (ABG), Puerto Ayora, Ecuador, ⁶Escuela de Medicina Veterinaria, Universidad San Francisco de Quito, Quito, Ecuador

Domestic species, including equids, were introduced in the Galapagos Islands in the XIX century. Equine vector-borne diseases are circulating in South America but their occurrence in the Galapagos Island was unknown. The objective of this study was to detect the occurrence of West Nile virus (WNV), Usutu virus (USUV) and equine infectious anemia virus (EIAV) in the four Galapagos Islands raising equids if they were present at a prevalence >1%. Serum samples were collected from 411 equids belonging to 124 owners from April to July 2019. All the results were negative to the ELISA tests used suggesting that WNV, USUV and EIAV are not circulating in the equine population of the Galapagos Islands.

KEYWORDS

WNV, USUV, EIAV, West Nile, Usutu, equine infectious anemia, Galapagos Islands, Ecuador

Introduction

The Galapagos Islands are a set of insular territories 960 km off the coast of Ecuador and an Ecuadorian province. They have been listed as a UNESCO World Heritage Site. Only 3% of their territory is colonized while the remaining 97% belongs to the Galapagos National Park. The introduction of livestock into the islands dates back from the beginning of the XIX century. In 1832, there was a massive introduction of productive domestic species, including equids, which may have carried parasites or pathogens (1). In 2003, animal importation from mainland Ecuador was banned, except for day-old chicks. Equids are present among the four colonized islands: Santa Cruz, Isabela, San Cristóbal and Floreana. Equid movements are allowed between the Galapagos Islands for animal genetic improvement. Surveillance of viral equid disease through laboratory diagnosis had never been conducted. Considering the movement of animals between mainland Ecuador and the Galapagos Islands in the past, the likelihood of the occurrence of viruses affecting the equids into the islands could not be excluded. Vector-borne viruses could have also been introduced by vectors coming from mainland.

West Nile virus (WNV) and Usutu virus (USUV) are vector-borne viruses that can affect equids and humans (2). They belong to the family *Flaviviridae*, genus *Flavivirus*, and are

TABLE 1 Number of farms and equids sampled in the Galapagos Islands.

Island	No. of farms sampled	No. of equids sampled	Number of equids sampled per farm (min.–max.)	Age (min.–max.)
Santa Cruz	64	140	1–10	1 year–30 years
Isabela	25	171	1–36	3 months–30 years
San Cristóbal	29	80	1–10	3 months–20 years
Floreana	6	20	1–11	3 months–19 years
Total	124	411	1–36	3 months–30 years

responsible for multiple outbreaks of disease in different countries of the world. WNV and USUV transmission cycles include wild birds as amplifying hosts and ornithophilic mosquitoes as vectors but can also infect and cause disease in horses and humans, which serve as incidental dead-end hosts (3, 4). WNV is one of the pathogenic agents that can lead to equine neurological clinical signs, although the infection is not usually accompanied by presentation of clinical illness (5). WNV is endemic in parts of Africa, Europe, the Middle East, and Asia, and since 1999 has spread to North America, Mexico, South America, and the Caribbean (6). WNV has been detected in mainland Ecuador through serological surveys in equids (7) and also in Colombia, an Ecuadorian neighboring country (8). USUV is particularly pathogenic in a few species of birds and has been detected in many mammalian species, including equids, considered to be dead-end hosts (9). USUV has not been reported in Latin America but in view of the emergence of different arboviral diseases, such as chikungunya and Zika, it can be considered as a new emerging threat (10).

Equine infectious anemia (EIA) is caused by a lentivirus and can be transmitted mechanically by biting flies (the virus does not replicate in the vector) (11). EIA virus (EIAV) infection leads to recurring episodes of fever in equids, thrombocytopenia, and wasting symptoms. EIA has been detected in different regions in Brazil (12–14) and Argentina (15) and several outbreaks were reported in different regions in mainland Ecuador (16).

The present study aimed to detect the occurrence of WNV, USUV and EIAV in the Galapagos Islands where equids are raised.

Method

Study population

The Galapagos veterinary services had estimated that around 650 equids are present in the islands of Santa Cruz (~ 230 equids), Isabela (~ 270 equids), San Cristóbal (~ 100 equids) and Floreana (~ 30 equids). The equids often belong to farmers who raise other species, mainly cattle, and are used for farm work, transportation or leisure purposes. There are approximately 132 equid owners in the four

islands (63 in Santa Cruz, 34 in Isabela, 26 in San Cristobal, 9 in Floreana) and between one and 15 equids per farm.

Sampling

For each island, the sample size was calculated to detect a 1% seroprevalence with a 95% level of confidence at the level of the equid population using Epitools.¹ This yielded animal sample sizes of 144 in Santa Cruz, 171 in Isabela, 79 in San Cristobal and 24 in Floreana. A mean number of three equids per farm was used to estimate the total number of farms to be included in the study. A minimum of 10 animals were sampled per farm and in farms with less than 10 animals, all animals were tested. The farms and the animals were randomly selected.

Laboratory tests

Sera were tested using the commercial ID Screen West Nile competition multi-species ELISA kit (Innovative Diagnostics, Montpellier, France) according to manufacturer's instructions. This method, that uses pre-coated plates with the envelop (E) protein of WNV, estimates the competition between antibodies present in the animal serum and monoclonal anti-WNV E antibody conjugated to horseradish peroxidase (HRP). This ELISA, through cross-reactions, allows the detection of other flaviviruses belonging to the Japanese encephalitis virus serocomplex as USUV (17). Assays were interpreted according to Beck et al. (18).

To detect antibodies against EIAV, serum samples were screened using a commercial ELISA, Equine Infectious Anemia Virus Antibody Test Kit, ELISA v2 (VMRD, Pullman, USA). Assays were performed according to the manufacturer's instructions.

Results


Serum samples were collected from 411 equids (367 horses, 20 donkeys and 24 mules) in 124 farms in the four islands (Table 1; Figure 1) from April to July 2019. The sampled animals were between 3 months and 30 years old.

All animals had negative results in the serological tests for the Flavivirus genus and EIAV.

Discussion

The negative serological results obtained in this study suggest that the flavivirus that could affect equids, such WNV or USUV, and EIAV were not circulating in the Galapagos Islands above the prevalence value used to calculate the sample size for this study. This absence could go up to 30 years considering the age of some of the sampled equids.

¹ <https://epitools.ausvet.com.au/>



The Galapagos Islands fill the climatic conditions favorable for the circulation of flaviviruses or EIAV provided the presence of competent vectors. The islands has three mosquito vectors capable of transmitting WNV and USUV (*Culex quinquefasciatus*, *Aedes aegypti*, and *Aedes taeniorhynchus*) (19, 20), bird species related to competent avian hosts (*Dendroica petechia*, *Mimus* spp) (21) and horseflies that can transmit EIAV (20). Since the climatic and vector conditions are filled, it could be argued that the prevalence viruses could be <1% in the equid population. However, in territories where those viruses are circulating, the prevalence values were higher than 1%. For example, WNV has been reported to be at seroprevalence values of 11% in horses in equids in Algeria (serums collected from 2015 to 2017) (22), 44.9% in Spain further to an outbreak in 2020 (23) or 31.6% in asymptomatic horses in Mexico (24). Analysis of 28,089 equids included in a systematic review and meta-analysis of seroprevalence studies of WNV in equids from 16 European countries between 2001 and 2018 revealed a pooled seroprevalence of 8% (25). In Morocco, prevalence of USUV was estimated to be 4% in a survey conducted in military working horses (26). In Spain, of 341 feral horses tested from 2010 to 2020, 9% were found seropositive to USUV (27). Regarding EIAV, in countries where this virus is circulating, prevalence values can be as high as 44% [survey conducted in Argentina (28)] but has also be found at lower values [2% in horses of the Para State in Brazil (14)]. On the other hand, according to the local veterinary services, there had been no reports of disease in equids in the Galapagos Islands consistent with WNV, USUV or EIAV or mortalities linked with WNV in birds.

The fact that the Galapagos Islands have semi-isolated conditions and the ban on animal importation do not provide sufficient guarantees to prevent the introduction of these viruses. Their introduction could arise by windborne transportation of infected vectors, infected migratory birds or infected vectors carried by airplanes or ships. These introduction routes were explored by Kilpatrick et al. (29) through a quantitative risk assessment to predict the WNV introduction into Galapagos Islands. These authors also included day-old chicks imports and mosquitoes in the larval stage present in shipment of tires as likely sources of WNV infection. They found that mosquitoes transported on airplanes carrying tourists represented the highest risk of WNV reaching the Galapagos Islands by a vector pathway and that the risk of introduction through migratory birds was lower but non-negligible. Moreover, it has been estimated that 107 Diptera species have arrived on the Galapagos Islands through human introductions and that 42 species have naturally colonized the islands from mainland Americas (20). Bataille et al. (30) showed from the monitoring of aeroplanes and genetic analysis that *C. quinquefasciatus* was regularly introduced via aircraft into the Galapagos islands. Therefore, effective control methods should be implemented by the national authorities to minimize the likelihood of introduction of insects through these routes that could put in danger endemic wildlife of Galapagos as well as domestic animal populations. Those measures could be combined with surveillance measures to detect any possible introduction. As the equid population is naïve to WNV and EIAV it is possible that that they could display clinical signs and, hence, equid owners should

be made aware of the importance to report the occurrence of unexpected clinical cases. Future serum sampling carried out for other diseases in equids could also be used to detect flavivirus or EIAV. Hence, equids could act as sentinels to provide early warning of virus circulation.

Data availability statement

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

Ethics statement

The animal studies were approved by Agencia de Regulación y Control de la Bioseguridad y Cuarentena para Galápagos (ABG), Puerto Ayora, Ecuador. The studies were conducted in accordance with the local legislation and institutional requirements. Written informed consent was obtained from the owners for the participation of their animals in this study.

Author contributions

GZ: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Validation, Writing – original draft, Writing – review & editing. CB: Formal analysis, Resources, Writing – review & editing. J-CV-C: Formal analysis, Resources, Writing – review & editing. MA: Formal analysis, Writing – review & editing. MC: Resources, Supervision, Validation, Writing – review & editing. AV: Resources, Supervision, Validation, Writing – review & editing.

References

- Toral-Granda MV, Causton CE, Jager H, Trueman M, Izurieta JC, Araujo E, et al. Alien species pathways to the Galapagos Islands, Ecuador. *PLoS One*. (2017) 12:e0184379. doi: 10.1371/journal.pone.0184379
- Agliani G, Giglia G, Marshall EM, Grone A, Rockx BHG, van den Brand JMA. Pathological features of West Nile and Usutu virus natural infections in wild and domestic animals and in humans: A comparative review. *One Health*. (2023) 16:100525. doi: 10.1016/j.onehlt.2023.100525
- Lim SM, Koraka P, Osterhaus AD, Martina BE. West Nile virus: immunity and pathogenesis. *Viruses*. (2011) 3:811–28. doi: 10.3390/v3060811
- Golding M, Seechurn N, Baylis M, Johnson N. JMM Profile: Usutu virus. *J Med Microbiol*. (2023) 72:001652. doi: 10.1099/jmm.0.001652
- Castillo-Olivares J, Wood J. West Nile virus infection of horses. *Vet Res*. (2004) 35:467–83. doi: 10.1051/vetres:2004022
- Charrel RN, de Lamballerie X. West Nile virus, an emerging arbovirus. *Presse Med*. (2004) 33:1521–6. doi: 10.1016/S0755-4982(04)98977-4
- Coello-Peralta R. Serological test for West Nile virus in horses of Los Ríos, Ecuador. *Revista Ciencia UNEMI*. (2016) 9:59–62. doi: 10.29076/issn.2528-7737vol9iss20.2016pp59-62p
- Mattar S, Komar N, Young G, Alvarez J, Gonzalez M. Seroconversion for West Nile and St. Louis encephalitis viruses among sentinel horses in Colombia. *Mem Inst Oswaldo Cruz*. (2011) 106:976–9. doi: 10.1590/S0074-02762011000800012
- Roesch F, Fajardo A, Moratorio G, Vignuzzi M. Usutu virus: an arbovirus on the rise. *Viruses*. (2019) 11:640. doi: 10.3390/v11070640
- Paniz-Mondolfi AE, Villamil-Gomez WE, Rodriguez-Morales AJ. Usutu virus infection in Latin America: A new emerging threat. *Travel Med Infect Dis*. (2016) 14:641–3. doi: 10.1016/j.tmaid.2016.08.004
- Cook RF, Leroux C, Issel CJ. Equine infectious anemia and equine infectious anemia virus in 2013: a review. *Vet Microbiol*. (2013) 167:181–204. doi: 10.1016/j.vetmic.2013.09.031
- Barros ML, Borges AMC, Oliveira De ACS, Lacerda W, Souza De A, Aguiar DM. Spatial distribution and risk factors for equine infectious anaemia in the state of Mato Grosso. *Brazil Rev Sci Tech*. (2018) 37:971–83. doi: 10.20506/rst.37.3.2900
- Cursino AE, Vilela APP, Franco-Luiz APM, de Oliveira JG, Nogueira MF, Junior JPA, et al. Equine infectious anemia virus in naturally infected horses from the Brazilian Pantanal. *Arch Virol*. (2018) 163:2385–94. doi: 10.1007/s00705-018-3877-8
- de Padua BR, Dias RA, Fioravanti MCS, Borsanelli AC. Seroprevalence and risk factors associated with equine infectious anemia in the state of Goiás, Brazil. *Prev Vet Med*. (2022) 209:105781. doi: 10.1016/j.prevetmed.2022.105781
- Hebert L, Polledo G, Lecouturier F, Giorgi M, Beck C, Lowenski S, et al. Serological evidence of equine infectious anaemia, West Nile fever, Surra and equine piroplasmiasis in a herd of horses in northern Argentina. *Vet Parasitol Reg Stud Reports*. (2021) 24:100566. doi: 10.1016/j.vprsr.2021.100566
- Agrocalidad. *Programa nacional sanitario equino*. Ministerio de agricultura, ganadería acuicultura y pesca del Ecuador (2016).
- Chevalier V, Marsot M, Molia S, Rasamoelina H, Rakotondravao R, Pedrono M, et al. Serological evidence of West Nile and Usutu viruses circulation in domestic and wild birds in wetlands of Mali and Madagascar in 2008. *Int J Environ Res Public Health*. (2020) 17:1998. doi: 10.3390/ijerph17061998
- Beck C, Leparac-Goffart I, Desoutter D, Deberge E, Bichet H, Lowenski S, et al. Serological evidence of infection with dengue and Zika viruses in horses on French Pacific Islands. *PLoS Negl Trop Dis*. (2019) 13:e0007162. doi: 10.1371/journal.pntd.0007162
- Peck SB, Heraty J, Landry B, Sinclair BJ. Introduced insect fauna of an oceanic archipelago: the Galapagos Islands, Ecuador. *Am Entomol*. (1998) 44:218–37. doi: 10.1093/ae/44.4.218
- Sinclair BJ. An annotated checklist of the Diptera of the Galapagos archipelago (Ecuador). *Zootaxa*. (2023) 5283:1–102. doi: 10.11646/zootaxa.5283.1.1
- Eastwood G, Goodman SJ, Hilgert N, Cruz M, Kramer LD, Cunningham AA. Using avian surveillance in Ecuador to assess the imminence of West Nile virus incursion to Galapagos. *EcoHealth*. (2014) 11:53–62. doi: 10.1007/s10393-014-0911-5

RV: Data curation, Validation, Writing – review & editing. GG: Formal analysis, Resources, Validation, Writing – review & editing.

Funding

The author(s) declare that no financial support was received for the research, authorship, and/or publication of this article.

Acknowledgments

We would like to thank the equid owners who accepted to participate in the study. Thanks also go to technicians from Agencia de Regulación y Control de la Bioseguridad y Cuarentena para Galápagos (ABG).

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

22. Laabassi F, Dheilly N, Beck C, Amaral R, Gonzalez G, Gaudaire D, et al. Serological evidence of circulation of West Nile virus in equids in Algerian eastern drylands and its epidemiological risk factors. *Comp Immunol Microbiol Infect Dis.* (2023) 94:101947. doi: 10.1016/j.cimid.2023.101947
23. Garcia-Bocanegra I, Franco JJ, Leon CI, Barbero-Moyano J, Garcia-Mina MV, Fernandez-Molera V, et al. High exposure of West Nile virus in equid and wild bird populations in Spain following the epidemic outbreak in 2020. *Transbound Emerg Dis.* (2022) 69:3624–36. doi: 10.1111/tbed.14733
24. Alonso-Padilla J, Loza-Rubio E, Escribano-Romero E, Cordoba L, Cuevas S, Mejia F, et al. The continuous spread of West Nile virus (WNV): seroprevalence in asymptomatic horses. *Epidemiol Infect.* (2009) 137:1163–8. doi: 10.1017/S0950268809002325
25. Metz MBC, Olufemi OT, Daly JM, Barba M. Systematic review and meta-analysis of seroprevalence studies of West Nile virus in equids in Europe between 2001 and 2018. *Transbound Emerg Dis.* (2021) 68:1814–23. doi: 10.1111/tbed.13866
26. Durand B, Haskouri H, Lowenski S, Vachiere N, Beck C, Lecollinet S. Seroprevalence of West Nile and Usutu viruses in military working horses and dogs, Morocco, 2012: dog as an alternative WNV sentinel species? *Epidemiol Infect.* (2016) 144:1857–64. doi: 10.1017/S095026881600011X
27. Magallanes S, Llorente F, Ruiz-Lopez MJ, Martinez-de la Puente J, Soriguer R, Calderon J, et al. Long-term serological surveillance for West Nile and Usutu virus in horses in south-West Spain. *One Health.* (2023) 17:100578. doi: 10.1016/j.onehlt.2023.100578
28. Ricotti S, Garcia MI, Veaute C, Bailat A, Lucca E, Cook RF, et al. Serologically silent, occult equine infectious anemia virus (EIAV) infections in horses. *Vet Microbiol.* (2016) 187:41–9. doi: 10.1016/j.vetmic.2016.03.007
29. Kilpatrick AM, Daszak P, Goodman SJ, Rogg H, Kramer LD, Cedeno V, et al. Predicting pathogen introduction: West Nile virus spread to Galapagos. *Conserv Biol.* (2006) 20:1224–31. doi: 10.1111/j.1523-1739.2006.00423.x
30. Bataille A, Cunningham AA, Cedeno V, Cruz M, Eastwood G, Fonseca DM, et al. Evidence for regular ongoing introductions of mosquito disease vectors into the Galapagos Islands. *Proc Biol Sci.* (2009) 276:3769–75. doi: 10.1098/rspb.2009.0998