



**HAL**  
open science

## **Plant mutations: slaying beautiful hypotheses by surprising evidence**

Niklas Tysklind, Sylvain Schmitt, Patrick Heuret, Valérie Troispoux, Mélanie Beraud, Jocelyne Cazal, Emilie Chancerel, Charlotte Cravero, Erwan Guichoux, Olivier Lepais, et al.

### ► **To cite this version:**

Niklas Tysklind, Sylvain Schmitt, Patrick Heuret, Valérie Troispoux, Mélanie Beraud, et al.. Plant mutations: slaying beautiful hypotheses by surprising evidence. EVOLTREE Conference 2023 - Resilient Forests for the Future, UNITBV, Braşov, Romania, Sep 2023, Brasov, Romania. 10.31926/evoltree.2023 . hal-04664198

**HAL Id: hal-04664198**

**<https://hal.inrae.fr/hal-04664198v1>**

Submitted on 29 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# **Resilient Forests for the Future**

**EVOLTREE Conference 2023**

**12 – 15 September 2023**

**UNITBV, Braşov, Romania**

**Book of Abstracts**



Transilvania University of Braşov

## Session 3:

### Evolutionary genomics

Wednesday 13 September 16:20 - 18:55

Chair: Dr. Annika Perry, UK Centre for Ecology & Hydrology, UK

### Oral presentations

- 16:20        **Keynote**  
**Lind, Brandon M.:**  
The limits of predicting maladaptation to future climate using genomic data
- 17:05        **Gonzalez-Martinez, Santiago C.:**  
Genetic variation within and among populations underlies adaptation to climate in a Mediterranean pine species
- 17:25        **Tysklind, Niklas:**  
Plant mutations: slaying beautiful hypotheses by surprising evidence
- 17:45               ***Short Break***
- 17:55        **Păun, Ovidiu:**  
The drivers of the adaptive radiation of persimmon trees on New Caledonia, a biodiversity hotspot
- 18:15        **Gathercole, Louise:**  
Genomic evidence for extensive introgression amongst British oaks and assessment of mal-adaptation
- 18:35        **Paulo, Octávio S.:**  
A landscape genomic approach on the cork oak and holm oak in the Western Mediterranean: insights into the species' capacity to respond to climate change

# Plant mutations: slaying beautiful hypotheses by surprising evidence

Contribution ID: 141

**Tysklind, Niklas; Schmitt, Sylvain; Heuret, Patrick; Troispoux, Valérie; Beraud, Mélanie; Cazal, Jocelyn; Chancerel, Émilie; Cravero, Charlotte; Guichoux, Erwan; Lepais, Olivier; Loureiro, João; Marande, William; Martin, Olivier; Vincent, Gregoire; Chave, Jérôme; Plomion, Christophe; Leroy, Thibault; Heuertz, Myriam**

Mutation is the source of all genetic diversity, on which selection can act allowing adaptation. Early differentiation between soma and the germline in animal development protects the germline, and thus the offspring, from the accumulation of somatic mutations. In plants, somatic mutations are expected to occur and accumulate along the branches and the differentiation between soma and germline is debated. There are several predictions on the fate of somatic mutations occurring in plant tissues: exposure to UV radiation increases mutagenesis; mutations have generally low frequency in plant tissues; branching topology of the tree dictates mutation distribution; and mutations at high frequency have a higher chance of being transmitted to the next generation. We produced a unique plant dataset of 60 high-coverage whole-genome sequences of two tropical tree species (*Dicorynia guianensis* and *Sextonia rubra*) and identified 18,274 de novo somatic mutations on branches exposed to high and low light, almost all at low frequency in tissues. We demonstrate that: 1) mutation rates and mutation spectra are not demonstrably affected by differences in UV exposure; 2) mutation phylogenies deviate from the branching topology of the tree; and 3) low-frequency mutations are transmitted to the next generation. Altogether, our results suggest far more complex links between plant growth, ageing, UV exposure, and mutation rates than commonly thought. Specially, the transmissibility of low frequency mutations, which are the most abundant, indicates a non-negligible role of low frequency somatic mutations as heritable mutations fuelling plant evolution.

Keywords: mutation, tropical trees, genomics, evolution, heritability