



HAL
open science

Quantification of zoonotic pathogenic bacteria in manures and raw digestates of three biogas plants

Christine Ziebal, Martine Denis, Céline Druilhe, Caroline Le Maréchal, Lorine Derongs, Elisabeth Répérant, Sandra Rouxel, Lorette Heurtevent, Typhaine Poezevara, Bérengère Nagard, et al.

► To cite this version:

Christine Ziebal, Martine Denis, Céline Druilhe, Caroline Le Maréchal, Lorine Derongs, et al.. Quantification of zoonotic pathogenic bacteria in manures and raw digestates of three biogas plants. *Animal Hygiene as a Fundament of One Health and Welfare improving biosecurity, environment and food quality*, ISAH, Sep 2019, Wroclaw, Poland. hal-04673878

HAL Id: hal-04673878

<https://hal.inrae.fr/hal-04673878v1>

Submitted on 20 Aug 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Quantification of zoonotic pathogenic bacteria in manures and raw digestates of three biogas plants

Ziebal Christine^{1,3}, Denis Martine^{2,3}, Druilhe Céline^{1,3}, Le Maréchal Caroline^{2,3}, Lorine Derongs^{1,3}, Sophie Michel-Leroux^{1,3}, Repérant Elisabeth^{2,3}, Sandra Rouxel^{2,3}, Lorette Heurtevent^{1,3}, Typhaine Poëzévara^{2,3}, Boschet Evelyne^{2,3}, Nagard Bérengère^{2,3}, Martin Laure^{2,3}, Catherine Houdayer^{2,3}, Frédéric Barbut^{2,3}, Pourcher Anne-Marie^{1,3}

1. Irstea, Rennes, France

2. Anses, Unité HQPAP, F-22440 Ploufragan, France

3. Université Européenne de Bretagne, Rennes, France

Introduction

On-farm anaerobic digestion (AD) which has expanded in Europe, produced biogas and a by-product (digestate) used as fertilizer. However, very few studies have been devoted to the effect of AD on the level of pathogenic bacteria. The study aims to evaluate the impact of mesophilic AD on the concentrations of zoonotic pathogenic bacteria (*Listeria monocytogenes*, *Salmonella* spp, thermotolerant campylobacters) and on indicator bacteria (*E. coli* and enterococci) in manures and in raw digestates sampled in three biogas plants (BGP).

Materials and Methods

Samples were collected in storage tank (manures) and at the outlet valve of the digester (raw digestates) from three BGP during a one-year period (8 sampling dates per BGP, 3 replicates per sampling date). Two digesters were fed with pig manure. The third digester was essentially fed with cattle manure. Indicator bacteria were enumerated on specific agar media. Pathogenic bacteria were quantified after an enrichment step by MPN. A total of 144 samples (72 inputs, 72 digestates) were analyzed.

Results

The pathogens have been detected in manure at a frequency ranged between 92 and 96%. They were also present in raw digestate at a frequency of 38% (thermotolerant campylobacters), 83% (*L. monocytogenes*) and 88 % (*Salmonella* spp.). The concentrations were lower in raw digestates than in manures. The average concentrations of thermotolerant campylobacters, *L. monocytogenes* and *Salmonella* spp. in raw digestates were 2, 23 and 5 MPN/g, respectively. The average levels of *E.coli* and enterococci (10^4 - 10^5 cfu/g in manure) decreased by 1 to 2 Log₁₀ during AD.

Conclusion

Although mesophilic AD did not eliminate the pathogenic bacteria, it reduced their concentration. Raw digestates are not directly spread on agricultural land. It is thus likely that their post-treatment (eg. storage or post-digestion) also contribute to limit the spread of pathogens together with the digestate onto agricultural land.