

Adoption of cereal-legume intercropping in France: a matter of outlets?

Elodie Yan, Marco Carozzi, Philippe Martin

Université Paris-Saclay, INRAE, AgroParisTech, UMR SADAPT, F-91120, Palaiseau, France

Contact: elodie.yan@inrae.fr



INRAE



ESA
EUROPEAN SOCIETY FOR AGRONOMY

Benefits of cereal-legume intercropping

CEREAL-LEGUME INTERCROPS CONSIDERED

Cereal(s) + legume(s)

On the same plot

At the same time

All harvested as principal crops

*Example of a cereal-legume intercrop:
wheat-faba bean intercrop
© E. Yan*




- **Better use of resources** (Brooker et al., 2015)
 - Stability of yields or even higher yields (e.g. Malézieux et al., 2009; Lithourgidis et al., 2011; Bedoussac et al., 2015)
 - Quality of production (e.g. Malézieux et al., 2009; Lithourgidis et al., 2011; Bedoussac et al., 2015; Li et al., 2023)
 - **Control of pests and diseases** (e.g. Ratnadass et al., 2012; Beillouin et al., 2021)
 - **Competition against weeds** (e.g. Corre-Hellou et al., 2011)
- Reduction of chemical inputs on farms (Yan et al., 2024)

Low adoption rate but local dynamics

- Cereal-legume intercrops: <1% French arable lands in 2022 (IGN, 2023)
 - Technical, material, and economic barriers (Himanen et al., 2016; Mamine & Farès, 2020; Timaeus et al., 2022)
- Increasing dynamic and diffusion from hotspots
 - Adoption linked to organic farming, animal husbandry? (Verret et al., 2020; Timaeus et al., 2022)
 - Potential roles of local groups? e.g. machinery cooperatives, exchanges between farmers (Himanen et al., 2016; Casagrande et al., 2017; Ha et al., 2023)

Hypothesis formulated from case studies



Share of cereal-legume intercrops on total departmental arable lands in 2015, 2018 and 2021, in French departments.

Source of the data: French Land Parcel Identification System (Yan et al., in revision)


AIM OF THE STUDY

Identifying factors favouring the adoption of cereal-legume intercrops

- **National level:** check the hypothesis formulated from the literature and based on case studies
- **Local level:** spot the possible particularities in areas with contrasted agricultural contexts


Material & methods

Quantitative study based on exhaustive data



2020 French agricultural census – Exhaustive survey on French farms (once every ten years)


- 43,968 farms, representative of the French arable crop, livestock and mixed crop–livestock farming systems
- 42 variables (*literature review*)



Farms' characteristics at national scale

Variable ranking and selection with a random forest

1



National-level random forest variable importance plot.


Importance is scaled from 0 to 100 relative to the first most important variable (Yan et al., in revision).

Farms' characteristics at national scale

2

Interpretation with a Classification And Regression Tree

UAA: useable arable area







Values: probability of presence of cereal-legume intercrops

Farms' characteristics: local particularities

- 1'
- 3

- Globally same characteristics than at national level
- Organic farming: less important in the Southern area
 - Cereal-legume intercropping long-established for livestock feed autonomy (Clouet et al., 1986)
→ On-farm grain storage: indicator for feed storage?




Random forest variable importance plots for the Western, Eastern, and Southern areas.

In each plot, the importance is scaled from 0% to 100% relative to the first most important variable.

Variables in bold are the common variables between the top 10 at the national level and the top 10 at the area level (Yan et al., in revision).

Potential links between the main identified factors and outlets for cereal-legume intercrops



Take-home messages

- Factors identified in the literature: confirmed at national and local scales
 - Organic farming, livestock and feed autonomy
 - Farm machinery cooperatives: may provide equipment to overcome material barriers, but also technical and psychological barriers (peer exchanges, learning)
- How to foster the adoption of cereal-legume intercropping?
 - Short-term dynamics: local collective actions
 - Long-term dynamics: securing the outlets, developing market opportunities
- Quantitative study based on farm characteristics: limited understanding of farmers' objectives
 - Relevance of a mixed approach with on-farm surveys
- Methodology applicable to other practices

Thank you!

Authors: [Elodie Yan](#), Marco Carozzi, Philippe Martin

Contact: elodie.yan@inrae.fr

Access to some confidential data, on which is based this work, has been made possible within a secure environment offered by CASD – Centre d'accès sécurisé aux données (Ref. 10.34724/CASD)



References (1/2)

- Bedoussac, L., Journet, E.-P., Hauggaard-Nielsen, H., Naudin, C., Corre-Hellou, G., Jensen, E. S., Prieur, L., & Justes, E. (2015). Ecological principles underlying the increase of productivity achieved by cereal-grain legume intercrops in organic farming. A review. *Agronomy for Sustainable Development*, 35(3), 911-935. <https://doi.org/10.1007/s13593-014-0277-7>
- Beillouin, D., Ben-Ari, T., Malézieux, E., Seufert, V., & Makowski, D. (2021). Positive but variable effects of crop diversification on biodiversity and ecosystem services. *Global Change Biology*, 27(19), 4697-4710. <https://doi.org/10.1111/gcb.15747>
- Brooker, R. W., Bennett, A. E., Cong, W.-F., Daniell, T. J., George, T. S., Hallett, P. D., Hawes, C., Iannetta, P. P. M., Jones, H. G., Karley, A. J., Li, L., McKenzie, B. M., Pakeman, R. J., Paterson, E., Schöb, C., Shen, J., Squire, G., Watson, C. A., Zhang, C., ... White, P. J. (2015). Improving intercropping : A synthesis of research in agronomy, plant physiology and ecology. *New Phytologist*, 206(1), 107-117. <https://doi.org/10.1111/nph.13132>
- Casagrande, M., Alletto, L., Naudin, C., Lenoir, A., Siah, A., & Celette, F. (2017). Enhancing planned and associated biodiversity in French farming systems. *Agronomy for Sustainable Development*, 37(6), 57. <https://doi.org/10.1007/s13593-017-0463-5>
- Clouet, Y., Guilloneau, A., & Ruf, T. (1986). *Diagnostic du système agraire et des systèmes de production en Ségala Aveyronnais*.
- Corre-Hellou, G., Dibet, A., Hauggaard-Nielsen, H., Crozat, Y., Gooding, M., Ambus, P., Dahlmann, C., von Fragstein, P., Pristeri, A., Monti, M., & Jensen, E. S. (2011). The competitive ability of pea-barley intercrops against weeds and the interactions with crop productivity and soil N availability. *Field Crops Research*, 122(3), 264-272. <https://doi.org/10.1016/j.fcr.2011.04.004>
- Ha, T. M., Manevska-Tasevska, G., Jäck, O., Weih, M., & Hansson, H. (2023). Farmers' intention towards intercropping adoption : The role of socioeconomic and behavioural drivers. *International Journal of Agricultural Sustainability*, 21(1), 2270222. <https://doi.org/10.1080/14735903.2023.2270222>
- Himanen, S. J., Mäkinen, H., Rimhanen, K., & Savikko, R. (2016). Engaging Farmers in Climate Change Adaptation Planning : Assessing Intercropping as a Means to Support Farm Adaptive Capacity. *Agriculture*, 6(3), Article 3. <https://doi.org/10.3390/agriculture6030034>
- IGN. (2023). *Institut National de l'Information Géographique et Forestière—IGN. Base de Données*. <https://geoservices.ign.fr/telechargement>
- Lithourgidis, A., Dordas, C., Damalas, C., & Vlachostergios, D. (2011). Annual intercrops : An alternative pathway for sustainable agriculture. *Australian Journal of Crop Science*, 5, 396-410.
- Malézieux, E., Crozat, Y., Dupraz, C., Laurans, M., Makowski, D., Ozier-Lafontaine, H., Rapidel, B., Tourdonnet, S., & Valantin-Morison, M. (2009). Mixing plant species in cropping systems : Concepts, tools and models. A review. *Agronomy for Sustainable Development*, 29(1), 43-62. <https://doi.org/10.1051/agro:2007057>

References (2/2)

- Mamine, F., & Farès, M. (2020). Barriers and Levers to Developing Wheat–Pea Intercropping in Europe : A Review. *Sustainability*, 12(17), Article 17. <https://doi.org/10.3390/su12176962>
- Ratnadass, A., Fernandes, P., Avelino, J., & Habib, R. (2012). Plant species diversity for sustainable management of crop pests and diseases in agroecosystems : A review. *Agronomy for Sustainable Development*, 32(1), 273-303. <https://doi.org/10.1007/s13593-011-0022-4>
- Timaeus, J., Ruigrok, T., Siegmeier, T., & Finckh, M. R. (2022). Adoption of Food Species Mixtures from Farmers' Perspectives in Germany : Managing Complexity and Harnessing Advantages. *Agriculture*, 12(5), Article 5. <https://doi.org/10.3390/agriculture12050697>
- Verret, V., Pelzer, E., Bedoussac, L., & Jeuffroy, M.-H. (2020). Tracking on-farm innovative practices to support crop mixture design : The case of annual mixtures including a legume crop. *European Journal of Agronomy*, 115, 126018. <https://doi.org/10.1016/j.eja.2020.126018>
- Yan, E., Munier-Jolain, N., Martin, P., & Carozzi, M. (2024). Intercropping on French farms : Reducing pesticide and N fertiliser use while maintaining gross margins. *European Journal of Agronomy*, 152, 127036. <https://doi.org/10.1016/j.eja.2023.127036>