



**HAL**  
open science

## A New Derivative of Retro-2 Displays Antiviral Activity against Respiratory Syncytial Virus

Adrien Le Rouzic, Jenna Fix, Robin Vinck, Sandrine Kappler-Gratias,  
Romain Volmer, Franck Gallardo, Jean-François Éléouët, Mathilde Keck,  
Jean-Christophe Cintrat, Julien Barbier, et al.

### ► To cite this version:

Adrien Le Rouzic, Jenna Fix, Robin Vinck, Sandrine Kappler-Gratias, Romain Volmer, et al.. A  
New Derivative of Retro-2 Displays Antiviral Activity against Respiratory Syncytial Virus. 2024.  
hal-04684074

**HAL Id: hal-04684074**

**<https://hal.inrae.fr/hal-04684074v1>**

Preprint submitted on 2 Sep 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

---

# A New Derivative of Retro-2 Displays Antiviral Activity against Respiratory Syncytial Virus

---

[Adrien Le Rouzic](#) , Jenna Fix , [Robin Vinck](#) , Sandrine Kappler-Gratias , [Romain Volmer](#) , [Franck Gallardo](#) , [Jean-François Eléouët](#) , [Mathilde Keck](#) , [Jean-Christophe Cintrat](#) , [Julien Barbier](#) , [Daniel Gillet](#) <sup>\*</sup> , [Marie Galloux](#) <sup>\*</sup>

Posted Date: 26 October 2023

doi: 10.20944/preprints202310.1733.v1

Keywords: Respiratory syncytial virus; Retro-2.2; fusion protein; retrograde transport; antiviral



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

# A New Derivative of Retro-2 Displays Antiviral Activity against Respiratory Syncytial Virus

Adrien Le Rouzic <sup>1,2</sup>, Jenna Fix <sup>1</sup>, Robin Vinck <sup>2,3</sup>, Sandrine Kappler-Gratias <sup>4</sup>, Romain Volmer <sup>5</sup>, Franck Gallardo <sup>4</sup>, Jean-François Eléouët <sup>1</sup>, Mathilde Keck <sup>2</sup>, Jean-Christophe Cintrat <sup>3</sup>, Julien Barbier <sup>2</sup>, Daniel Gillet <sup>2,\*</sup> and Marie Galloux <sup>1,\*</sup>

<sup>1</sup> Université Paris-Saclay-Versailles St Quentin, INRAE, Unité de Virologie et Immunologie Moléculaires (VIM), F-78350 Jouy-en-Josas, France; adrien.le-rouzic@inrae.fr (A.L.R.); jenna.fix@inrae.fr (J.F.); jean-francois.eleouet@inrae.fr (J.-F.E.)

<sup>2</sup> Université Paris-Saclay, CEA, INRAE, Département Médicaments et Technologies pour la Santé (DMTS), SIMoS, 91191 Gif-sur-Yvette, France; robin.vinck13@gmail.com (R.V.); mathilde.keck@cea.fr (M.K.); julien.barbier@cea.fr (J.B.)

<sup>3</sup> Université Paris-Saclay, CEA, INRAE, Département Médicaments et Technologies pour la Santé (DMTS), SCBM, 91191 Gif-sur-Yvette, France; jean-christophe.cintrat@cea.fr

<sup>4</sup> NeoVirTech SAS, 1 Place Pierre Potier, 31000 Toulouse, France; skappler@neovirtech.com (S.K.-G.); fgallardo@neovirtech.com (F.G.)

<sup>5</sup> ENVT Université de Toulouse, INRAE, IHAP, UMR 1225, 31300 Toulouse, France; romain.volmer@envt.fr

\* Correspondence: daniel.gillet@cea.fr (D.G.); marie.galloux@inrae.fr (M.G.)

**Abstract:** Human respiratory syncytial virus (hRSV) is the most common cause of bronchiolitis and pneumonia in new-borns, with all children being infected before the age of two. Reinfections are very common throughout life, and can cause severe respiratory infections in the elderly and immunocompromised adults. Although vaccines and preventive antibodies have recently been licensed for use in specific subpopulations of patients, there is still no therapeutic treatment commonly available for these infections. Here, we investigated the potential antiviral activity of Retro-2.2, a derivative of the cellular retrograde transport inhibitor Retro-2, against hRSV. We show that Retro-2.2 inhibits hRSV replication in cell culture, and impairs the ability of hRSV to form syncytia. Our results suggest that Retro-2.2 treatment affects virus spread by disrupting the trafficking of the viral *de novo* synthesized F and G glycoproteins to the plasma membrane, leading to a defect in virion morphogenesis. Taken together, our data show that targeting intracellular transport may be an effective strategy against hRSV infection.

**Keywords:** Respiratory syncytial virus; Retro-2.2; fusion protein; retrograde transport; antiviral

## 1. Introduction

The human respiratory syncytial virus (hRSV) is the leading cause of severe lower respiratory tract infection in young children worldwide, and the first cause of hospitalization of new-borns under 6 months of age [1]. A systemic multisite study has shown that hRSV is the first etiological agent responsible for severe pneumonia (more than 30%) in hospitalized children in Asia and Africa [2]. It is noteworthy that hRSV is also a frequent cause of otitis in infants, and that children who suffer from severe hRSV infection are at risk of developing further respiratory complications such as asthma [3,4]. Re-infections occur throughout life with symptoms of common cold in healthy adults, but lead to severe lower respiratory infections in immunocompromised and elderly people [5]. Despite the high impact of hRSV infections on public health, the first vaccines to prevent severe hRSV disease in the elderly were approved by the Food and Drug Administration only in 2023 (GSK, Pfizer) [6][7], followed by a vaccine dedicated to pregnant women (Pfizer) [8]. The only specific treatment commercialized for new-borns remains humanized monoclonal antibodies directed against the fusion protein F responsible for viral entry. The first commercialized antibody (Palivizumab, Synagis®) displays poor benefit-cost ratio, limiting its use to prophylaxis of severe infection in pre-term and high-risk infants [9]. The new prophylactic antibody targeting the pre-fusion form of F

(Nirsevimab, Beyfortus®) has been very recently approved in the European Union for the prevention of hRSV lower respiratory tract disease in newborns and infants during their first hRSV season [10,11].

hRSV belongs to the order *Mononegavirales* and the family *Pneumoviridae*, which are enveloped viruses with a single strand negative sense RNA genome [12]. The attachment of the viral particle to the cell surface and the entry into the host cell are mediated by the viral glycoproteins G and F. More specifically, virus entry involves the fusion between viral and cellular membranes. This fusion is pH-independent [13], mediated by F, and could occur at the plasma membrane or in endosomes after macropinocytosis [14]. Of note, in cell culture the F protein is responsible for the fusion between cells, leading to the formation of syncytia which characterises hRSV infection [15][16]. The fusion allows the release into the cell cytoplasm of the ribonucleoprotein complex (RNP), composed of the viral genome encapsidated by the nucleoprotein N, the viral polymerase L, its cofactor the phosphoprotein P, and the M2-1 protein (transcription factor). Transcription and replication of the viral genome are initiated by the viral polymerase L which is responsible for all RNAs synthesis activities (RNA-dependent RNA polymerase, mRNAs capping and methyltransferase). These steps take place in viral factories that are membraneless liquid organelles, called inclusion bodies (IBs) [17,18]. The viral mRNAs are then exported from IBs to the cytoplasm where viral proteins are expressed. At the final stages of the viral cycle, RNPs and structural viral proteins are addressed to the plasma membrane where assembly occurs, leading to the formation of new filamentous virions which are released by membrane fission.

To date, most developed antiviral strategies aim at targeting the viral F protein to impair virus entry, or the viral polymerase L [19–22]. However, none of these compounds have been commercialized yet, with the main limitation being the emergence of escape mutants upon treatment. In this context, the alternative of targeting cellular pathways involved in the viral cycle may represent a new antiviral approach.

Here, we assessed the potential antiviral activity against hRSV of Retro-2.2, a new optimized derivative of the cellular retrograde transport inhibitor Retro-2 [23]. Retro-2 was initially identified as an inhibitor of the ricin toxin [23]. This molecule and its optimized derivative Retro-2.1 were subsequently shown to display an antiviral activity against various viruses such as polyomaviruses and papillomaviruses [24–26], adeno-associated viruses [27], poxviruses [28,29], Ebola and Marburg viruses [30], and more recently SARS-CoV-2 [31]. Here, our results reveal that hRSV replication is inhibited by Retro-2.2 treatment in infected cell cultures. The treatment induced a defect in the formation of syncytia among hRSV infected cells that correlated with a defect of virions release. Further investigation revealed that Retro-2.2 affects the trafficking of the viral membrane proteins F and G to the plasma membrane. Altogether, our results highlight a new antiviral approach against hRSV, based on the inhibition of the final stages of virion assembly and release.

## 2. Results

### 2.1. Retro-2.2 antiviral activity against hRSV


Following *in vitro* screening of Retro-2.1 and different optimized Retro-2 analogues (EP3886859) on DNA and RNA viruses, we investigated the antiviral activity of the Retro-2.2 molecule (Figure 1A) against hRSV virus.

The antiviral activity of Retro-2.2 against hRSV was first assessed on HEp-2 cells infected with a recombinant hRSV expressing the mCherry reporter gene (mCherry-rRSV), which allows to quantify the inhibitory activity of compounds by fluorescence measurement [32–34]. Cells were infected with mCherry-rRSV either in the absence or the presence of serial dilutions of Retro-2.2 for 2 h. After infection, the medium was replaced by medium containing serial dilutions of Retro-2.2, and fluorescence intensity was measured 48 h post-infection. In parallel, the potential cytotoxicity of Retro-2.2 in the same conditions was evaluated on non-infected cells. As shown on Figure 1B, similar results were obtained when the cells were treated during or only after infection, with half-maximal inhibitory concentration (IC<sub>50</sub>) close to 1.6  $\mu$ M. These results suggest that Retro-2.2 has no impact on



viral particles or on the early steps of virus entry. A cytotoxic activity of Retro-2.2 was observed, with a calculated half-maximal cytotoxic concentration ( $CC_{50}$ ) close to 15  $\mu$ M. The impact of Retro-2.2 on hRSV replication was also validated on different cell lines usually used as a model for hRSV studies, i.e. A549, BEAS-2B, and 293T, at a concentration of 3  $\mu$ M (Supplementary Figure S1). We then further evaluated the effect of Retro-2.2 treatment on viral shedding by quantifying virions in the culture medium of infected cells, by plaque formation assay. Our results showed that treatment with 2  $\mu$ M Retro-2.2 induces a  $\log_{10}$  decrease in viral titer compared to untreated cells, as well as a reduction of plaque size (Figure 1C). A direct observation of infected cells by fluorescence also revealed a strong effect of Retro-2.2 treatment on the size of the syncytia formed during hRSV infection, with the presence of large syncytia in the control condition compared to numerous but small syncytia in the presence of Retro-2.2 (Figure 1D).

Altogether, these results show that Retro-2.2 inhibits hRSV replication in cell culture. This inhibition is characterized by a defect of syncytia formation and of viral particles release.



**Figure 1.** Impact of Retro-2.2 treatment on hRSV replication on Hep-2 cells. **(A)** Scheme of the synthesis of Retro-2.2 compound. **(B)** Cells were infected for 2 h with mCherry-rRSV at MOI 0.2, in the presence (blue curve) or the absence (red curve) of serial dilutions of Retro-2.2. The medium was then replaced to incubate cells in the presence of serial dilutions of Retro-2.2 for 48 h. The viral replication was quantified by measurement of the mCherry fluorescence. In parallel, cell viability


upon treatment with Retro-2.2 was quantified in non-infected HEp-2 cells (black curve). Error bars are standard deviations from duplicates. Data are representative of three experiments. The curves were fitted in Graph Pad software using a four parameters logistic (4PL) regression.  $IC_{50}$  and  $CC_{50}$  are indicated. **(C)** Titration of the virions released into the culture media (diluted 10 times) from non-infected cells (NI) or cells infected with mCherry-rRSV, and treated with DMSO (control) or Retro-2.2 at 2  $\mu$ M, by plaque formation assay. Calculated viral titers are indicated in pfu/ml. **(D)** Representative images of HEp-2 cell cultures infected with mCherry-rRSV at MOI 0.2, 48 hours post-infection in the absence or presence of Retro-2.2 at 1.25 and 10  $\mu$ M. Upper panel, mCherry fluorescence of infected cells; Lower panel, nuclei colored with Hoechst. Scale bars, 500  $\mu$ m.

## 2.2. Retro-2.2 effect on the formation of IBs and virions assembly in hRSV infected cells

In order to characterize the inhibitory effect of Retro-2.2 against hRSV, we then studied the impact of Retro-2.2 treatment on the formation of IBs and on viral assembly. BEAS-2B cells were infected with recombinant hRSV expressing the firefly Luciferase (Luc-rRSV), which displays the same hRSV sequence and kinetics of replication compared to mCherry-rRSV [32], at MOI 0.2. Cells were incubated for 36 h in the absence or presence of 2  $\mu$ M Retro-2.2, a concentration close to the  $IC_{50}$ . The N and F proteins were then immunolabelled in order to observe IBs and virions assembly, respectively. Within syncytia observed in untreated cell cultures, the F protein was concentrated in granules around the nuclei, but also at the plasma membrane where a large number of short filaments were observed (Figure 2A). Similar localization of F was observed in cells treated with Retro-2.2, although the distribution of F appeared less homogenous along the plasma membrane and longer filaments were detected compared to the control condition. However, N labelling allowed to validate the presence of IBs of similar shape in both conditions, suggesting that Retro-2.2 did not impair viral genome transcription or replication, nor viral proteins expression (Figure 2A).

Although the formation of syncytia in untreated cell culture makes the comparison difficult, these results suggest that Retro-2.2 could affect the activity of the F protein. We therefore studied the effect of Retro-2.2 in the context of cells expressing only the F protein. BEAS-2B cells were transfected with an hRSV F encoding plasmid before labelling with anti-F and anti-GM130, a marker of the cis-Golgi network, 24 h post-transfection. As shown in Figure 2B, the expression of F alone induced the formation of syncytia similar to those observed during infection and the treatment with Retro-2.2 at 2  $\mu$ M impaired the formation of large syncytia. F was also observed in perinuclear granules distinct from the cis-Golgi, and at the plasma membrane in both conditions. It is noteworthy that no modification of the GM130 labelling was observed in the presence of Retro-2.2.

These data suggest that the inhibition of large syncytia formation upon Retro-2.2 treatment during hRSV infection is mainly dependent on its effect on the F protein.



**Figure 2.** Effect of Retro-2.2 treatment on the localization of N and F viral proteins. **(A)** BEAS-2B cells were infected for 2 h with Luc-rRSV, and the medium was changed to incubate cells in the absence (DMSO) or the presence of Retro-2.2 at 2  $\mu$ M. Thirty-six hours post infection cells were fixed, immunostained with anti-F (red) and anti-N (green) antibodies, and nuclei were colored with Hoechst. Insets correspond to the zoom of the regions indicated by dashed squares. **(B)** BEAS-2B cells were transfected with a plasmid encoding the F protein. Twenty-four hours post-transfection cells were fixed and immunostained with anti-F (red) and anti-GM130 (green) antibodies, and nuclei were colored with Hoechst. Images are from confocal microscopy sections. Scale bars, 50  $\mu$ m.


### 2.3. Retro-2.2 affects the trafficking of the F protein to the plasma membrane

The F protein is a class I fusion protein expressed through the endoplasmic reticulum and Golgi network as a precursor F0. It is glycosylated, cleaved by furine to generate the F1 (50 kDa) and F2 (15 kDa) subunits linked by two disulphide bonds, and forms trimers folded in a prefusion conformation

[35] (Figure 3A). We therefore investigated if the Retro-2.2 treatment could impair the maturation of F and/or its trafficking to the plasma membrane. We first analyzed the migration profile on SDS-PAGE of the F protein from infected cells incubated in the absence or presence of Retro-2.2 for 48 hours. Cells were lysed in reducing and non-reducing condition to observe the F1 subunit only or F1 linked to F2, respectively. Surprisingly, Western blot analysis revealed similar migration profiles in all the conditions, with only a band of approximately 50 kDa corresponding to the F1 subunit (Figure 3B). Although we were not able to detect the full-length form of F in these conditions, these data show that Retro-2.2 did not abrogate the cleavage by furine. However, the treatment with Retro-2.2 led to a lower quantity of F protein, which correlates with the inhibition of viral spreading. Nevertheless, we cannot exclude a partial defect of F maturation or the induction of F degradation upon Retro-2.2 treatment. In order to assess the effect of Retro-2.2 on F trafficking, we quantified then the presence of F at the plasma membrane. To overcome the potential bias linked to cell fusion and of immunostaining, we generated a F surrogate which consisted in the insertion in place of the F ectodomain of the sequences of the mCherry and of the HiBit peptide, a 11 amino-acids long optimized peptide of the split Nano Luciferase (Figure 3A) [36]. The plasmid encoding this chimeric protein was transfected in HEp-2 cells which were incubated for 24 h in the absence or presence of 2  $\mu$ M Retro-2.2. Analysis of mCherry by fluorescence revealed that the chimeric protein mainly concentrated within intracellular granules upon Retro-2.2 treatment compared to the control condition (Figure 3C, upper panel). Western blot analysis of the chimeric protein expression revealed with an anti-mCherry antibody allowed to confirm that Retro-2.2 did not affect cell transfection nor protein expression or stability (calculated expected size, 42 kDa) (Figure 3C, lower panel). In order to quantify F expression in cells and its specific localization at the plasma membrane, the luminescence activity in total cell lysates or in the cell culture medium was then measured. Our data revealed similar levels of total luminescence in both conditions, but a clear decrease of extracellular luminescence induced by Retro-2.2 treatment (Figure 3D). Finally, we performed a similar experiment to quantify extracellular luminescence in the presence of increased concentrations of Retro-2.2. Our results show a dose dependent inhibition of the presence of the chimeric protein at the plasma membrane, with a calculated  $IC_{50}$  close to 1.3  $\mu$ M (Figure 3E), that correlated with the antiviral  $IC_{50}$  of Retro-2.2 (Figure 1A).

Altogether these results show that Retro-2.2 impairs the trafficking of F to the plasma membrane.






**Figure 3.** Impact of Retro-2.2 on the trafficking of the F protein to the plasma membrane. **(A)** Schematic representation of the F primary sequence and of the derived chimeric HiBit-mCherry-F protein. The F0 form and the F1 and F2 subunits of F are indicated. Arrows indicate the cleavage sites on F0. The points and lines indicate the cysteines and disulfide bonds. SP, signal peptide; FP, fusion peptide; TM, transmembrane domain; Ctail, intracellular C-terminal domain. The amino acid sequence of the chimeric protein is indicated with domains as colored boxes, and the mCherry sequence written in red. **(B)** Western blot analysis of the migration profile of the F protein from infected HEp-2 cells lysates incubated in the presence of DMSO or Retro-2.2 (2 μM) during 48 h. The presence or absence of β-Mercaptoethanol (β-ME) is indicated. Membranes were sequentially incubated with anti-F, anti-N and anti-tubulin antibodies. Mass weights are indicated on the left. **(C)** HEp-2 cells were transfected with the plasmid encoding the chimeric HiBit-mCherry-F protein. Six hours post-transfection, the medium was replaced to incubate the cells in the presence of Retro-2.2 at 2 μM. Twenty-four hours post-transfection cells were fixed and nuclei were stained with Hoechst, before analysis of the mCherry fluorescence by confocal microscopy (upper panel). Scale bars, 20 μm. The expression of the

chimeric protein was also validated by Western blot using an anti-mCherry antibody (lower panel) and anti-tubulin as a control. **(D)** In parallel, the presence of the chimeric protein in whole cell lysates or at the cell surface was quantified by luminescence measurement. Data are means SEM from quadruplicates. \*\*,  $P < 0.01$ , ns, not significant. Data are representative of three experiments. **(E)** Similar experiment was performed in the presence of serial dilutions of Retro-2.2 and the presence of the chimeric protein at the cell surface was quantified by extracellular luminescence measurement. Error bars are standard deviations from duplicates. Data are representative of two experiments. The curves were fitted in Graph Pad software using a four parameters logistic (4PL) regression.  $IC_{50}$  is indicated.

#### 2.4. Impact of Retro-2.2 on the G glycoprotein trafficking to the plasma membrane

We then studied whether the antiviral activity of Retro-2.2 against hRSV was restricted to F or could also modify the trafficking of the G glycoprotein to the plasma membrane. The G glycoprotein is expressed as a type II membrane protein (Figure 4A), but can also be secreted as a truncated form [37,38]. Although the transmembrane G protein is involved in the initial attachment to the host cell membrane during entry, the G protein was shown to be dispensable for hRSV replication in cell culture [39]. Therefore, the potential effect of Retro-2.2 on this protein could not have been observed using our previous assays. Using a similar strategy described above for F quantification, we generated a chimeric protein by replacing the extracellular domain of G by the HiBit sequence in fusion with EGFP (Figure 4A), and performed the assay described above. Observation of HEp-2 cells transfected with this construct for 24 h in the absence or presence of Retro-2.2 revealed the presence of chimeric protein within intracellular compartments, but no chimeric protein was clearly detected at the plasma membrane (Figure 4B). However, quantification by luminescence of the chimeric protein expression allowed to validate its presence at the plasma membrane, and revealed that Retro-2.2 treatment induced a decrease of the quantity of the chimeric protein at the plasma membrane (Figure 4C). Of note, the expression of the chimeric protein was similar in cells treated with Retro-2.2 compared to the control condition, as confirmed by total luminescence quantification and Western blot analysis of the chimeric protein expression revealed with anti-GFP antibody (calculated expected size, 38 kDa) (Figure 4C). Our results also showed a dose dependent inhibition of the presence of this chimeric protein at the plasma membrane upon Retro-2.2 treatment, with a calculated  $IC_{50}$  close to 1.1  $\mu$ M (Figure 4D).

These results suggest that Retro-2.2 could affect not only F but also G trafficking to the plasma membrane.



**Figure 4.** Impact of Retro-2.2 on the addressing of the hRSV G protein to the plasma membrane. **(A)** Schematic representation of the G protein and of the derived chimeric HiBit-EGFP-G protein. The points and line indicate the cysteines and disulfide bonds. Ntail, intracellular N-terminal domain; TM, transmembrane domain; CX<sub>3</sub>C, binding domain on the CX<sub>3</sub>C receptor. The amino acid sequence of the chimeric protein is indicated with domains represented as colored boxes, and the EGFP sequence


written in green. **(B)** HEp-2 cells were transfected with the plasmid encoding the chimeric HiBit-EGFP-G protein. Six hours post-transfection, the medium was replaced to incubate the cells in the presence of Retro-2.2 at 2  $\mu$ M. Twenty-four hours post-transfection cells were fixed and nuclei were stained with Hoechst, before analysis of the EGFP fluorescence by confocal microscopy. Scale bars, 20  $\mu$ m. **(C)** In parallel, the luminescence in whole cell lysates or in the extracellular environment was quantified (upper panel), and expression of the chimeric protein was validated by Western blot using an anti-GFP antibody. Data are means SEM from quadruplicats. \*,  $P < 0.05$ ; ns, not significant. Data are representative of two experiments. **(D)** Similar experiment was performed in the presence of serial dilutions of Retro-2.2 and the presence of the chimeric protein at the cell surface was quantified by extracellular luminescence measurement. Error bars are standard deviations from duplicates. Data are representative of two experiments. The curves were fitted in Graph Pad software using a four parameters logistic (4PL) regression.  $IC_{50}$  is indicated.

### 2.5. Down-expression of syntaxin-5 and Sec16A impair hRSV replication and syncytia formation

We then considered whether Retro-2.2 antiviral activity could be related to the mechanism of action already described for the Retro-2 compounds. Syntaxin-5 (Syn5) is a SNARE protein that is involved in cellular retrograde transport pathways. Syn5 was previously described to be relocated in the presence of Retro-2 [23]. Sec16A is a protein located at the reticulum endoplasmic exit sites that regulates the formation of COPII vesicles. Sec16A is involved in the endoplasmic reticulum secretion pathway. It was previously identified as a target of the Retro-2 compound [40]. We therefore studied the impact of the down-regulation of Syn5 or Sec16A expression on hRSV replication. Human epithelial A549 cells were transfected with siRNA for 24 h, then infected for 48 h with Luc-rRSV with a MOI of 0.2. Quantification of viral replication by luminescence measurement on cells lysates revealed that down-regulation of Syn5 or Sec16A both induced a 30-40% decrease of the infection (Figure 5A). Western blot analysis of cell lysates confirmed the down-expression of the two isoforms of Syn5 at this final time point (Figure 5A) [41]. Unfortunately, we did not manage to validate Sec16A extinction as we did not succeed to reveal Sec16A expression by Western blot. In order to observe the impact of siRNA on syncytia formation during infection, the same experiment was performed on cells infected with mCherry-rRSV. As shown on Figure 5C, down-regulation of Syn5 and Sec16A expression impaired the formation of large syncytia compared to the control condition.

Therefore, our data reveal that, similarly to the treatment by Retro-2.2, down-regulation of Syn5 or Sec16A expression affects the capacity of hRSV to form large syncytia in cell culture. These observations suggest that Retro-2.2 viral inhibition could be linked to its inhibitory activity on the retrograde pathway.





**Figure 5.** Downregulation of Syn5 and Sec16A expression impair hRSV replication and syncytia formation. HEp-2 cells were transfected with control siRNA (siCT) or siRNA targeting syntaxin-5 (siSyn5) or Sec16A (siSec16A) during 24 h, before infection with Luc-rRSV at MOI 0.2 during 48 h. **(A)** Viral replication was quantified by measuring the luminescence in cell lysates. Data are means SEM from quadruplicates. \*,  $P < 0.05$ . Data are representative of three experiments. **(B)** Western blot analysis of Syn5 expression in total cells lysates. **(C)** HEp-2 cells transfected with siRNA as described above were infected with mCherry-rRSV at MOI 0.2 during 48 h. Representative images of mCherry fluorescence of infected HEp-2 cell cultures. Scale bars, 200  $\mu\text{m}$ .

### 3. Discussion

Although vaccines against hRSV have been recently commercialized, there is still a need for antiviral treatments to treat hRSV infections. Many attempts have been made by pharmaceutical companies to develop compounds specifically targeting the viral F and L proteins. However, the emergence of escape mutants upon treatment highlights the necessity to search for alternatives and/or complementary antiviral approaches. Among those, strategies aiming at interfering with intracellular trafficking pathways required for pathogen infections could be promising.

In the present study, we evaluated the potential antiviral activity of the Retro-2.2 compound. This compound was derived from the structure-activity relationship study performed to increase the efficiency of the initial Retro-2 molecule and its optimized Retro-2.1 analogue (EP3886859). Retro-2.2 was found twice more effective against Shiga toxins than Retro-2.1, Shiga toxins being reference bacterial toxins using the retrograde transport in our structure-activity relationship study. Retro-2 and Retro-2.1 were shown to be active against a large panel of viruses [42]. Here, we first showed that Retro-2.2 presents an antiviral activity against hRSV in cell cultures, in the micromolar range. However, we also detected a non-expected cytotoxic effect of the compound in our conditions, with a  $CC_{50}$  close to 15  $\mu\text{M}$ . This low selectivity index value compared to the one obtained for the optimized Retro-2.1 molecule in previous studies [43,44] will need to be investigated.

Our data also show that Retro-2.2 did not impact the viral particle or the initial infection nor the formation of viral factories, but induced a defect of viral particles release associated to a decrease of syncytia formation in cell culture. Altogether, these results suggest that Retro-2.2 targets a late stage of the viral cycle, and more specifically the F protein function. By engineering chimeric transmembrane fluorescent proteins which display the HiBit peptide fused with the transmembrane

sequences of the F or G proteins, we were able to reveal that Retro-2.2 could impair the traffic of the F but also of the G protein to the plasma membrane. However, despite several attempts of co-immunofluorescence labelling of F and proteins of either the endoplasmic reticulum, the Golgi apparatus or the endosomes (not shown), we were not able to characterize a specific site of retention of the F and G proteins inside the cells. Noteworthy, in order to decipher the specificity of action of Retro-2.2 against hRSV, and to gain information on the cytotoxicity of the treatment, it would be necessary to evaluate the impact of Retro-2.2 treatment on the trafficking of cellular proteins known to be addressed to the plasma membrane such as TGN46 or Cl-MPR. In parallel, we showed that down-regulation of Syn5 and Sec16A, two proteins that were described as indirect and direct targets of Retro-2 respectively, also inhibited hRSV replication in cell culture, and the formation of large syncytia. Taken together, our results strongly suggest that, by modifying the cellular retrograde pathway, Retro-2.2 affects the trafficking of hRSV membrane proteins to the plasma membrane. However, we cannot exclude that Retro-2.2 treatment could also alter the full maturation of the F and G proteins.

As mentioned above, Retro-2.2 displays only low antiviral activity in cell culture. Nevertheless, given the dual effect of the treatment on F and G proteins, and based on the fact that G protein is not critical for hRSV replication in cell culture but favours *in vivo* replication, it would be interesting to test the antiviral activity of this compound *in vivo* in a mouse model. For this purpose, one would expect that, as recently shown for Retro-2.1, the formulation in a thermosensitive hydrogel could foster the delivery of Retro-2.2 *in vivo* [45].

In conclusion, our data highlight that molecules that interfere with the trafficking of intracellular vesicles could be used as antivirals against hRSV. Optimisation of such molecules to improve their selectivity index, their solubility and their delivery *in vivo*, still remains necessary. Finally, our results allowed to extend the list of the pathogens sensitive to Retro-2 derivatives, and reinforce the potential interest of these molecules as broad-spectrum antivirals.

## 4. Materials and Methods

### 4.1. Cells

HEp-2 cells (ATCC, CCL-23), A549 (ATCC, CCL-185) and 293T cells were maintained in Dulbecco's modified Eagle's medium (DMEM; Eurobio) supplemented with 10% heat-inactivated foetal calf serum (FCS; Eurobio), 1% L-glutamine, and 1% penicillin-streptomycin. The transformed human bronchial epithelial cell line (BEAS-2B) (ATCC CRL-9609) was maintained in RPMI 1640 medium (Eurobio) supplemented with 10% FCS, 1% L-glutamine, and 1% penicillin-streptomycin. The cells were grown at 37°C in 5% CO<sub>2</sub> and transfected using Lipofectamine 2000 (Invitrogen) as described by the manufacturer.

### 4.2. Retro-2.2 compound synthesis

A mixture containing 6-fluoro-1-methyl-2*H*-benzo[*d*][1,3]oxazine-2,4(1*H*)-dione (500 mg, 2.56 mmol), 2-aminophenol (280 mg, 2.56 mmol) and 5-(2-methyl-1,3-thiazol-4-yl)-2-thiophenecarbaldehyde (536 mg, 2.56 mmol) in acetic acid (3 mL), was heated at 130 °C in a microwave synthesizer (CEM) for 4 h. The crude mixture was diluted in ethyl acetate and a saturated NaHCO<sub>3</sub> aqueous solution was added. The mixture was stirred vigorously for 1 h. The aqueous layer was extracted two more times with EtOAc. The combined organic phase was dried over magnesium sulfate, filtered off and the solvent was evaporated under vacuum. Purification by column chromatography (CombiFlash Serlabo; column 40G) on silica gel using cyclohexane/ethyl acetate (60/40) afforded **1** as a pale brown solid (985 mg, 85% yield). At the end of the synthesis the compound was analyzed by <sup>1</sup>H and <sup>13</sup>C nuclear magnetic resonance (NMR), and mass spectrometry (LC-MS). The *m/z* calculated for C<sub>23</sub>H<sub>18</sub>FN<sub>3</sub>O<sub>2</sub>S<sub>2</sub>H [M+H]<sup>+</sup> (Retro-2.2) was 452.4Da, with a purity of 99.5%.

#### 4.3. Viruses and titration by plaque formation assay

The recombinant mCherry-rRSV and Luc-rRSV corresponding to the RSV Long strain expressing either the mCherry or the firefly luciferase protein were amplified in HEp-2 cells and titrated using a plaque assay procedure as previously described [32]. Briefly, for titration, cells were infected with serial 10-fold dilutions of the viral supernatant in complete minimum essential medium (MEM). The overlay was prepared with Avicel RC581 microcrystalline cellulose (FMC Biopolymer) at a final concentration of 0.6% in complete MEM containing 1% fetal calf serum. After 6 days at 37°C with 5% CO<sub>2</sub>, plaques were revealed by staining of the cell layers with a solution containing 0.5% crystal violet and 20% ethanol, and the number of PFU per well was counted.

#### 4.4. Plasmids

pcDNA3.1 codon-optimized plasmid for mammalian expression encoding the hRSV F was a gift from Marty Moore, Emory University [46]. The sequences encoding the chimeric proteins (primary sequences shown in Figures 3 and 4) were optimized for eukaryotic expression and commercially made (GeneCust). The full-length sequence of mCherry-HiBit-F was cloned into pCI vector (Genbank accession n° U47119). For HiBit-GFP-G expression, the optimized sequence encoding the N-terminal and transmembrane domains of G, in fusion with the HiBit sequence was clone in fusion with EGFP into the pEGFP-N2 vector (Clontech).

#### 4.5. Antibodies

The following primary antibodies were used for immunostaining and/or immunoblotting: mouse anti-F (BioRad, Hercules, USA), mouse anti-F 2F7 (Abcam, Cambridge, UK), mouse anti-G (Abcam), rabbit anti-N antiserum [47], rabbit anti-GM130 (Abcam), mouse anti-Syntaxin-5 (Santa Cruz, Dallas, USA), mouse monoclonal anti- $\beta$  tubulin antibody (Sigma-Aldrich, Saint-Louis, USA), rabbit anti-GFP (Invitrogen, Waltham, USA), and anti-mCherry (Rockland immunochemicals, Limerick, USA).

Secondary antibodies directed against mouse and rabbit IgG coupled to HRP (SeraCare, Milford, USA) were used for immunoblotting, and antibodies directed against mouse and rabbit IgG coupled to Alexa 594 or Alexa 488 (Invitrogen) respectively were used for immunostaining.

#### 4.6. Antiviral and cytotoxic activity assays

Cells were seeded at  $5 \times 10^4$  cells per well in 96-well plates. Twenty-four hours later, the cells were infected at a multiplicity of infection (MOI) of 0.2 for 2 h with mCherry-rRSV or Luc-rRSV diluted in MEM without phenol red and without FCS. In parallel, Retro-2.2 compound was serially diluted in dimethyl sulfoxide (DMSO) and then further diluted in MEM without phenol red medium containing 2% FCS. Two hours post-infection, the cell culture medium was replaced by the medium containing Retro-2.2, and plates were incubated at 37 °C for 24 to 48 h before analysis. The mCherry fluorescence was measured using a spectrophotometer (Tecan Infinite M200PRO) with excitation and emission wavelengths of 580 and 620 nm, respectively. For Luc-rRSV replication quantification, cells were lysed in luciferase lysis buffer (30 mM Tris [pH 7.9], 10 mM MgCl<sub>2</sub>, 1 mM DTT, 1% Triton X-100, and 15% glycerol). After the addition of luciferase assay reagent (Promega), luminescence was measured using a Tecan Infinite M200 Pro luminometer (Tecan, Männedorf, Switzerland). The values obtained for noninfected HEp-2 cells was used as a standard for the fluorescence and luminescence background levels respectively, and the value obtained for infected and untreated cells was used to normalize the data. Cytotoxicity assays were performed on non-infected cells incubated for 48 h with the serial dilutions of Retro-2.2, using the CellTiter-Glo luminescent cell viability kit (Promega). Each experiment was performed in duplicate and repeated at least twice. IC<sub>50</sub> and CC<sub>50</sub> values were calculated by fitting the data to a sigmoidal curve equation in GraphPad software (GraphPad Prism 10).

#### 4.7. siRNA experiments

The control siRNA and siRNAs against Syn-5 and Sec16A were purchased from Qiagen (Qiagen, Hilden, Germany). Each siRNA is a pool containing 4 different sequences (Catalog number: SI03075345, SI03052966, SI00048636 and SI00048615). A549 cells were transfected with the indicated siRNAs at a final concentration of 20 nM by reverse transfection into 96-well plates ( $7 \times 10^4$  cells per well) using Lipofectamine RNAiMAX (Thermo Fisher) according to the manufacturer's instructions. Briefly, a mixture containing Opti-MEM (Invitrogen), Lipofectamine RNAiMAX, and siRNA was incubated for 5 min at room temperature before being deposited at the bottom of the wells. The cells in MEM without antibiotics were then added dropwise before incubation at 37°C with 5% CO<sub>2</sub>. After 24 h of transfection in the presence of siRNA, the medium was removed, and the cells were infected following the protocol described above. Forty-eight hours post-infection, the mCherry fluorescence was measured using a spectrophotometer (Tecan Infinite M200PRO) with excitation and emission wavelengths of 580 and 620 nm, respectively. The values obtained for noninfected HEp-2 cells were used as a standard for the fluorescence background level, and the values obtained for infected cells transfected with the siCT were used to normalize the data. Each experiment was performed in quadruplicate and repeated three times. For each experiment, cells were finally lysed in Laemmli, and Syn5 and tubulin expression was analyzed by Western blotting.

#### 4.8. SDS-PAGE and Western blot analysis

Protein samples were separated by electrophoresis on 12% polyacrylamide gels in Tris-glycine buffer. All samples were boiled for 3 min prior to electrophoresis. Proteins were then transferred to a nitrocellulose membrane (BioRad). The blots were blocked with 5% non-fat milk in PBS Tween20 0.2%, followed by incubation with mouse anti-F antibody (1:1,000), rabbit anti-N antiserum (1:5,000), mouse anti-tubulin (1:1,000), or mouse anti-Syn5 (1:2,000) and horseradish peroxidase (HRP)-conjugated anti-mouse or anti-rabbit (1:10,000) secondary antibodies (SeraCare). Western blots were developed using Clarity™ Western ECL substrate (Bio-Rad) and exposed using Bio-Rad ChemiDoc Touch Imaging System.

#### 4.9. Immunofluorescence

Cells were seeded in 24-wells plates containing coverslip. Plates were infected or transfected depending on the assay. Cells were fixed using 4% paraformaldehyde for 30 minutes at room temperature, then permeabilized in PBS buffer containing 3% bovine albumin serum (BSA) and 0.1% Triton X-100 for 30 minutes at room temperature. Cells were then washed once in PBS-3% BSA-0.05% Tween20, and incubated with relevant mouse or rabbit primary antibody (anti-F 1:1,000, anti-N 1:20,000 ; anti-GM130 1:1,000) diluted in PBS-3% BSA-0.05% Tween20 for 1 h at room temperature. Cells were washed 3 times with PBS-3% BSA-0.05% Tween20, then incubated with goat anti-mouse or goat anti-rabbit secondary antibody coupled to Alexa 488 or Alexa 594. Cells were then washed 3 times with PBS-3% BSA-0.05% Tween20 and two times with PBS. For nucleus labeling, cells were exposed to Hoechst 33342 stain (Invitrogen) during incubation with secondary antibodies. Coverslips were mounted with ProLong Gold antifade reagent (Invitrogen). Images were obtained using an inverted SP8x confocal microscope (Leica) using a 63x oil immersion objective. Maximum projections of optical Z slices are shown. Images were processed with Leica imaging software LAS X and ImageJ/FIJI software.



#### 4.10. HiBit Luminescence assay

Hep-2 cells were seeded at  $5 \times 10^4$  cells per well in 96-well plates or  $2 \times 10^5$  cells 24-well plates containing coverslips. Cells were transfected in Opti-MEM (Gibco) with 0.2  $\mu\text{g}$  DNA per well (96-well plate) or 1  $\mu\text{g}$  DNA per well (24-well plate) using Lipofectamine 2000 for 5 h. The medium was then changed for Opti-MEM supplemented with Retro-2.2 at various concentrations. Twenty-four hours post-transfection, the expression of chimeric proteins in total cell lysates or at the cell surface was quantified by luminescence measurement using the Nano-Glo HiBiT Lytic Detection System (Promega, N3030) or the Nano-Glo HiBiT Extracellular Detection System (Promega, N2420), respectively, according to manufacturer's protocol. For each experiment, cells were finally lysed in Laemmli, and the expression of chimeric proteins was analyzed by Western blotting.

#### 4.11. Statistical analysis

A nonparametric Mann-Whitney test (comparison of two groups) ( $n = 4$  or  $6$ ) was used to compare unpaired values (GraphPad Prism software). Significance is represented in the figures (\*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ).

**Supplementary Materials:** The following supporting information can be downloaded at the website of this paper posted on Preprints.org. Figure S1: Antiviral activity of Retro-2.2 against hRSV infection of different cell lines.

**Author Contributions:** Conceptualization, J.B., D.G., M.K. and M.G.; methodology, A.L.-R., J.F.E., J.B., D.G., M.G.; validation, A.L.-R., M.G.; formal analysis, D.G., J.B., A.L.-R., and M.G.; investigation, A.L.-R., J.F., R.V., J.-C.C., M.G., J.B., S.K.-G., R.V., F.G.; writing—original draft preparation, D.G., and M.G.; writing—review and editing, J.B., D.G., and M.G.; supervision, D.G. and M.G.; project administration, D.G. and M.G.; funding acquisition, M.K., D.G., J.B. and M.G. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by the joint ministerial program of R&D against CBRNE risks, CEA and under Contract ANR20-CE18-0016 (SMERSEC), ANR-11-IDEX-0003-02 (PFI n° 20LU1000-131 ReCoVer and OTP n° 20LU1000\_138 PIMVir). A.L.-R. was funded by Université Paris-Saclay.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Acknowledgments:** We thank Dr Marie-Anne Rameix-Welti for the plasmid of pmCherry-F. The present work has benefited from Imagerie-Gif core facility supported by l'Agence Nationale de la Recherche (ANR-11-EQPX-0029/Morphoscope, ANR-10-INBS-04/FranceBioImaging; ANR-11-IDEX-0003-02/ Saclay Plant Sciences).

**Conflicts of Interest:** S.K.-G. and F.G. are employees and shareholders of NeoVirTech SAS. No other conflict of interest to declare.

## References

- Shi, T.; McAllister, D. A.; O'Brien, K. L.; Simoes, E. A. F.; Madhi, S. A.; Gessner, B. D.; Polack, F. P.; Balsells, E.; Acacio, S.; Aguayo, C.; Alassani, I.; Ali, A.; Antonio, M.; Awasthi, S.; Awori, J. O.; Azziz-Baumgartner, E.; Baggett, H. C.; Baillie, V. L.; Balmaseda, A.; Barahona, A.; Basnet, S.; Bassat, Q.; Basualdo, W.; Bigogo, G.; Bont, L.; Breiman, R. F.; Brooks, W. A.; Broor, S.; Bruce, N.; Bruden, D.; Buchy, P.; Campbell, S.; Carosone-Link, P.; Chadha, M.; Chipeta, J.; Chou, M.; Clara, W.; Cohen, C.; de Cuellar, E.; Dang, D. A.; Dash-Yandag, B.; Deloria-Knoll, M.; Dherani, M.; Eap, T.; Ebruke, B. E.; Echavarria, M.; de Freitas Lazaro Emediato, C. C.; Fasce, R. A.; Feikin, D. R.; Feng, L.; Gentile, A.; Gordon, A.; Goswami, D.; Goyet, S.; Groome, M.; Halasa, N.; Hirve, S.; Homaira, N.; Howie, S. R. C.; Jara, J.; Jroundi, I.; Kartasasmita, C. B.; Khuri-Bulos, N.; Kotloff, K. L.; Krishnan, A.; Libster, R.; Lopez, O.; Lucero, M. G.; Lucion, F.; Lupisan, S. P.; Marcone, D. N.; McCracken, J. P.; Mejia, M.; Moisi, J. C.; Montgomery, J. M.; Moore, D. P.; Moraleta, C.; Moyes, J.; Munywoki, P.; Mutyara, K.; Nicol, M. P.; Nokes, D. J.; Nymadawa, P.; da Costa Oliveira, M. T.; Oshitani, H.; Pandey, N.; Paranhos-Baccala, G.; Phillips, L. N.; Picot, V. S.; Rahman, M.; Rakoto-Andrianarivelo, M.; Rasmussen, Z. A.; Rath, B. A.; Robinson, A.; Romero, C.; Russomando, G.; Salimi, V.; Sawatwong, P.; Scheltema, N.; Schweiger, B.; Scott, J. A. G.; Seidenberg, P.; Shen, K.; Singleton, R.; Sotomayor, V.; Strand, T. A.; Sutanto, A.; Sylla, M.; Tapia, M. D.; Thamthitawat, S.; Thomas, E. D.; Tokarz, R.; Turner, C.; Venter, M.; Waicharoen, S.; Wang, J.; Watthanaworawit, W.; Yoshida, L. M.; Yu, H.; Zar, H. J.; Campbell, H.; Nair, H.; Network, R. S. V. G. E., Global, regional, and national disease burden estimates

- of acute lower respiratory infections due to respiratory syncytial virus in young children in 2015: a systematic review and modelling study. *Lancet* **2017**, 390, (10098), 946-958.
2. Pneumonia Etiology Research for Child Health Study, G., Causes of severe pneumonia requiring hospital admission in children without HIV infection from Africa and Asia: the PERCH multi-country case-control study. *Lancet* **2019**, 394, (10200), 757-779.
  3. Phillips, M.; Finelli, L.; Saiman, L.; Wang, C.; Choi, Y.; Patel, J., Respiratory Syncytial Virus-associated Acute Otitis Media in Infants and Children. *J Pediatric Infect Dis Soc* **2020**, 9, (5), 544-550.
  4. Shi, T.; Ooi, Y.; Zaw, E. M.; Utjesanovic, N.; Campbell, H.; Cunningham, S.; Bont, L.; Nair, H.; Investigators, R., Association Between Respiratory Syncytial Virus-Associated Acute Lower Respiratory Infection in Early Life and Recurrent Wheeze and Asthma in Later Childhood. *J Infect Dis* **2020**, 222, (Suppl 7), S628-S633.
  5. Busack, B.; Shorr, A. F., Going Viral-RSV as the Neglected Adult Respiratory Virus. *Pathogens* **2022**, 11, (11).
  6. Papi, A.; Ison, M. G.; Langley, J. M.; Lee, D. G.; Leroux-Roels, I.; Martinon-Torres, F.; Schwarz, T. F.; van Zyl-Smit, R. N.; Campora, L.; Dezutter, N.; de Schrevel, N.; Fissette, L.; David, M. P.; Van der Wielen, M.; Kostanyan, L.; Hulstrom, V.; Group, A. R.-S., Respiratory Syncytial Virus Prefusion F Protein Vaccine in Older Adults. *N Engl J Med* **2023**, 388, (7), 595-608.
  7. Walsh, E. E.; Perez Marc, G.; Zareba, A. M.; Falsey, A. R.; Jiang, Q.; Patton, M.; Polack, F. P.; Llapur, C.; Doreski, P. A.; Ilangovan, K.; Ramet, M.; Fukushima, Y.; Hussen, N.; Bont, L. J.; Cardona, J.; DeHaan, E.; Castillo Villa, G.; Ingilizova, M.; Eiras, D.; Mikati, T.; Shah, R. N.; Schneider, K.; Cooper, D.; Koury, K.; Lino, M. M.; Anderson, A. S.; Jansen, K. U.; Swanson, K. A.; Gurtman, A.; Gruber, W. C.; Schmoele-Thoma, B.; Group, R. C. T., Efficacy and Safety of a Bivalent RSV Prefusion F Vaccine in Older Adults. *N Engl J Med* **2023**, 388, (16), 1465-1477.
  8. Kampmann, B.; Madhi, S. A.; Munjal, I.; Simoes, E. A. F.; Pahud, B. A.; Llapur, C.; Baker, J.; Perez Marc, G.; Radley, D.; Shittu, E.; Glanternik, J.; Snaggs, H.; Baber, J.; Zachariah, P.; Barnabas, S. L.; Fausett, M.; Adam, T.; Perreras, N.; Van Houten, M. A.; Kantele, A.; Huang, L. M.; Bont, L. J.; Otsuki, T.; Vargas, S. L.; Gullam, J.; Tapiero, B.; Stein, R. T.; Polack, F. P.; Zar, H. J.; Staerke, N. B.; Duron Padilla, M.; Richmond, P. C.; Koury, K.; Schneider, K.; Kalinina, E. V.; Cooper, D.; Jansen, K. U.; Anderson, A. S.; Swanson, K. A.; Gruber, W. C.; Gurtman, A.; Group, M. S., Bivalent Prefusion F Vaccine in Pregnancy to Prevent RSV Illness in Infants. *N Engl J Med* **2023**, 388, (16), 1451-1464.
  9. Pinquier, D.; Crepey, P.; Tissieres, P.; Vabret, A.; Roze, J. C.; Dubos, F.; Cahn-Sellem, F.; Javouhey, E.; Cohen, R.; Weil-Olivier, C., Preventing Respiratory Syncytial Virus in Children in France: A Narrative Review of the Importance of a Reinforced Partnership Between Parents, Healthcare Professionals, and Public Health Authorities. *Infect Dis Ther* **2023**, 12, (2), 317-332.
  10. Griffin, M. P.; Yuan, Y.; Takas, T.; Domachowske, J. B.; Madhi, S. A.; Manzoni, P.; Simoes, E. A. F.; Esser, M. T.; Khan, A. A.; Dubovsky, F.; Villafana, T.; DeVincenzo, J. P.; Nirsevimab Study, G., Single-Dose Nirsevimab for Prevention of RSV in Preterm Infants. *N Engl J Med* **2020**, 383, (5), 415-425.
  11. Hammit, L. L.; Dagan, R.; Yuan, Y.; Baca-Cots, M.; Bosheva, M.; Madhi, S. A.; Muller, W. J.; Zar, H. J.; Brooks, D.; Grenham, A.; Wahlby-Hamren, U.; Mankad, V. S.; Ren, P.; Takas, T.; Abram, M. E.; Leach, A.; Griffin, M. P.; Villafana, T.; Group, M. S., Nirsevimab for Prevention of RSV in Healthy Late-Preterm and Term Infants. *N Engl J Med* **2022**, 386, (9), 837-846.
  12. Afonso, C. L.; Amarasinghe, G. K.; Banyai, K.; Bao, Y.; Basler, C. F.; Bavari, S.; Bejerman, N.; Blasdel, K. R.; Briand, F. X.; Briese, T.; Bukreyev, A.; Calisher, C. H.; Chandran, K.; Cheng, J.; Clawson, A. N.; Collins, P. L.; Dietzgen, R. G.; Dolnik, O.; Domier, L. L.; Durrwald, R.; Dye, J. M.; Easton, A. J.; Ebihara, H.; Farkas, S. L.; Freitas-Astua, J.; Formenty, P.; Fouchier, R. A.; Fu, Y.; Ghedin, E.; Goodin, M. M.; Hewson, R.; Horie, M.; Hyndman, T. H.; Jiang, D.; Kitajima, E. W.; Kobinger, G. P.; Kondo, H.; Kurath, G.; Lamb, R. A.; Lenardon, S.; Leroy, E. M.; Li, C. X.; Lin, X. D.; Liu, L.; Longdon, B.; Marton, S.; Maisner, A.; Muhlberger, E.; Netesov, S. V.; Nowotny, N.; Patterson, J. L.; Payne, S. L.; Paweska, J. T.; Randall, R. E.; Rima, B. K.; Rota, P.; Rubbenstroth, D.; Schwemmler, M.; Shi, M.; Smither, S. J.; Stenglein, M. D.; Stone, D. M.; Takada, A.; Terregino, C.; Tesh, R. B.; Tian, J. H.; Tomonaga, K.; Tordo, N.; Towner, J. S.; Vasilakis, N.; Verbeek, M.; Volchkov, V. E.; Wahl-Jensen, V.; Walsh, J. A.; Walker, P. J.; Wang, D.; Wang, L. F.; Wetzler, T.; Whitfield, A. E.; Xie, J. T.; Yuen, K. Y.; Zhang, Y. Z.; Kuhn, J. H., Taxonomy of the order Mononegavirales: update 2016. *Arch Virol* **2016**, 161, (8), 2351-60.
  13. Srinivasakumar, N.; Ogra, P. L.; Flanagan, T. D., Characteristics of fusion of respiratory syncytial virus with HEp-2 cells as measured by R18 fluorescence dequenching assay. *J Virol* **1991**, 65, (8), 4063-9.
  14. Krzyzaniak, M. A.; Zumstein, M. T.; Gerez, J. A.; Picotti, P.; Helenius, A., Host cell entry of respiratory syncytial virus involves macropinocytosis followed by proteolytic activation of the F protein. *PLoS Pathog* **2013**, 9, (4), e1003309.
  15. Kahn, J. S.; Schnell, M. J.; Buonocore, L.; Rose, J. K., Recombinant vesicular stomatitis virus expressing respiratory syncytial virus (RSV) glycoproteins: RSV fusion protein can mediate infection and cell fusion. *Virology* **1999**, 254, (1), 81-91.

16. Pierangeli, A.; Scagnolari, C.; Antonelli, G., Respiratory syncytial virus. *Minerva Pediatr* **2018**, *70*, (6), 553-565.
17. Garcia, J.; Garcia-Barreno, B.; Vivo, A.; Melero, J. A., Cytoplasmic inclusions of respiratory syncytial virus-infected cells: formation of inclusion bodies in transfected cells that coexpress the nucleoprotein, the phosphoprotein, and the 22K protein. *Virology* **1993**, *195*, (1), 243-7.
18. Rincheval, V.; Lelek, M.; Gault, E.; Bouillier, C.; Sitterlin, D.; Blouquit-Laye, S.; Galloux, M.; Zimmer, C.; Eleouet, J. F.; Rameix-Welti, M. A., Functional organization of cytoplasmic inclusion bodies in cells infected by respiratory syncytial virus. *Nat Commun* **2017**, *8*, (1), 563.
19. DeVincenzo, J. P.; Whitley, R. J.; Mackman, R. L.; Scaglioni-Weinlich, C.; Harrison, L.; Farrell, E.; McBride, S.; Lambkin-Williams, R.; Jordan, R.; Xin, Y.; Ramanathan, S.; O'Riordan, T.; Lewis, S. A.; Li, X.; Toback, S. L.; Lin, S. L.; Chien, J. W., Oral GS-5806 activity in a respiratory syncytial virus challenge study. *The New England journal of medicine* **2014**, *371*, (8), 711-22.
20. Detalle, L.; Stohr, T.; Palomo, C.; Piedra, P. A.; Gilbert, B. E.; Mas, V.; Millar, A.; Power, U. F.; Stortelers, C.; Allosery, K.; Melero, J. A.; Depla, E., Generation and Characterization of ALX-0171, a Potent Novel Therapeutic Nanobody for the Treatment of Respiratory Syncytial Virus Infection. *Antimicrob Agents Ch* **2015**, *60*, (1), 6-13.
21. Stevens, M.; Rusch, S.; DeVincenzo, J.; Kim, Y. I.; Harrison, L.; Meals, E. A.; Boyers, A.; Fok-Seang, J.; Huntjens, D.; Lounis, N.; Mari, N. K.; Remmerie, B.; Roymans, D.; Koul, A.; Verloes, R., Antiviral Activity of Oral JNJ-53718678 in Healthy Adult Volunteers Challenged With Respiratory Syncytial Virus: A Placebo-Controlled Study. *J Infect Dis* **2018**, *218*, (5), 748-756.
22. Cockerill, G. S., JNJ-5371678, Defining a Role for Fusion Inhibitors in the Treatment of Respiratory Syncytial Virus. *J Med Chem* **2020**, *63*, (15), 8043-8045.
23. Stechmann, B.; Bai, S. K.; Gobbo, E.; Lopez, R.; Merer, G.; Pinchard, S.; Panigai, L.; Tenza, D.; Raposo, G.; Beaumelle, B.; Sauvaire, D.; Gillet, D.; Johannes, L.; Barbier, J., Inhibition of retrograde transport protects mice from lethal ricin challenge. *Cell* **2010**, *141*, (2), 231-42.
24. Nelson, C. D.; Carney, D. W.; Derdowski, A.; Lipovsky, A.; Gee, G. V.; O'Hara, B.; Williard, P.; DiMaio, D.; Sello, J. K.; Atwood, W. J., A retrograde trafficking inhibitor of ricin and Shiga-like toxins inhibits infection of cells by human and monkey polyomaviruses. *mBio* **2013**, *4*, (6), e00729-13.
25. Maru, S.; Jin, G.; Desai, D.; Amin, S.; Shwetank; Lauver, M. D.; Lukacher, A. E., Inhibition of Retrograde Transport Limits Polyomavirus Infection In Vivo. *mSphere* **2017**, *2*, (6).
26. Treasure, T.; Nelson, C. D. S., Inhibition of JC polyomavirus infectivity by the retrograde transport inhibitor Retro-2.1. *Microbiol Immunol* **2020**, *64*, (12), 783-791.
27. Nonnenmacher, M. E.; Cintrat, J. C.; Gillet, D.; Weber, T., Syntaxin 5-dependent retrograde transport to the trans-Golgi network is required for adeno-associated virus transduction. *J Virol* **2015**, *89*, (3), 1673-87.
28. Sivan, G.; Weisberg, A. S.; Americo, J. L.; Moss, B., Retrograde Transport from Early Endosomes to the trans-Golgi Network Enables Membrane Wrapping and Egress of Vaccinia Virus Virions. *J Virol* **2016**, *90*, (19), 8891-905.
29. Harrison, K.; Haga, I. R.; Pechenick Jowers, T.; Jasim, S.; Cintrat, J. C.; Gillet, D.; Schmitt-John, T.; Digard, P.; Beard, P. M., Vaccinia Virus Uses Retromer-Independent Cellular Retrograde Transport Pathways To Facilitate the Wrapping of Intracellular Mature Virions during Virus Morphogenesis. *J Virol* **2016**, *90*, (22), 10120-10132.
30. Shtanko, O.; Sakurai, Y.; Reyes, A. N.; Noel, R.; Cintrat, J. C.; Gillet, D.; Barbier, J.; Davey, R. A., Retro-2 and its dihydroquinazolinone derivatives inhibit filovirus infection. *Antiviral Res* **2018**, *149*, 154-163.
31. Holwerda, M.; V'Kovski, P.; Wider, M.; Thiel, V.; Dijkman, R., Identification of an Antiviral Compound from the Pandemic Response Box that Efficiently Inhibits SARS-CoV-2 Infection In Vitro. *Microorganisms* **2020**, *8*, (12).
32. Rameix-Welti, M. A.; Le Goffic, R.; Herve, P. L.; Sourimant, J.; Remot, A.; Riffault, S.; Yu, Q.; Galloux, M.; Gault, E.; Eleouet, J. F., Visualizing the replication of respiratory syncytial virus in cells and in living mice. *Nature communications* **2014**, *5*, 5104.
33. Gaillard, V.; Galloux, M.; Garcin, D.; Eleouet, J. F.; Le Goffic, R.; Larcher, T.; Rameix-Welti, M. A.; Boukadiri, A.; Heritier, J.; Segura, J. M.; Baechler, E.; Arrell, M.; Mottet-Osman, G.; Nyanguile, O., A Short Double-Stapled Peptide Inhibits Respiratory Syncytial Virus Entry and Spreading. *Antimicrob Agents Chemother* **2017**, *61*, (4).
34. Galloux, M.; Gsponer, N.; Gaillard, V.; Fenner, B.; Larcher, T.; Vilotte, M.; Riviere, J.; Richard, C. A.; Eleouet, J. F.; Le Goffic, R.; Mettier, J.; Nyanguile, O., Targeting the Respiratory Syncytial Virus N-0-P Complex with Constrained alpha-Helical Peptides in Cells and Mice. *Antimicrob Agents Ch* **2020**, *64*, (10).
35. McLellan, J. S.; Ray, W. C.; Peeples, M. E., Structure and function of respiratory syncytial virus surface glycoproteins. *Curr Top Microbiol Immunol* **2013**, *372*, 83-104.
36. Oh-Hashi, K.; Furuta, E.; Fujimura, K.; Hirata, Y., Application of a novel HiBiT peptide tag for monitoring ATF4 protein expression in Neuro2a cells. *Biochem Biophys Rep* **2017**, *12*, 40-45.

37. Hendricks, D. A.; Baradaran, K.; McIntosh, K.; Patterson, J. L., Appearance of a soluble form of the G protein of respiratory syncytial virus in fluids of infected cells. *J Gen Virol* **1987**, 68 ( Pt 6), 1705-14.
38. Bukreyev, A.; Yang, L.; Fricke, J.; Cheng, L.; Ward, J. M.; Murphy, B. R.; Collins, P. L., The secreted form of respiratory syncytial virus G glycoprotein helps the virus evade antibody-mediated restriction of replication by acting as an antigen decoy and through effects on Fc receptor-bearing leukocytes. *J Virol* **2008**, 82, (24), 12191-204.
39. Karron, R. A.; Buonagurio, D. A.; Georgiu, A. F.; Whitehead, S. S.; Adamus, J. E.; Clements-Mann, M. L.; Harris, D. O.; Randolph, V. B.; Udem, S. A.; Murphy, B. R.; Sidhu, M. S., Respiratory syncytial virus (RSV) SH and G proteins are not essential for viral replication in vitro: clinical evaluation and molecular characterization of a cold-passaged, attenuated RSV subgroup B mutant. *Proc Natl Acad Sci U S A* **1997**, 94, (25), 13961-6.
40. Forrester, A.; Rathjen, S. J.; Daniela Garcia-Castillo, M.; Bachert, C.; Couhert, A.; Tepshi, L.; Pichard, S.; Martinez, J.; Munier, M.; Sierocki, R.; Renard, H. F.; Augusto Valades-Cruz, C.; Dingli, F.; Loew, D.; Lamaze, C.; Cintrat, J. C.; Linstedt, A. D.; Gillet, D.; Barbier, J.; Johannes, L., Functional dissection of the retrograde Shiga toxin trafficking inhibitor Retro-2. *Nat Chem Biol* **2020**, 16, (3), 327-336.
41. Miyazaki, K.; Wakana, Y.; Noda, C.; Arasaki, K.; Furuno, A.; Tagaya, M., Contribution of the long form of syntaxin 5 to the organization of the endoplasmic reticulum. *J Cell Sci* **2012**, 125, (Pt 23), 5658-66.
42. Gupta, N.; Noel, R.; Goudet, A.; Hinsinger, K.; Michau, A.; Pons, V.; Abdelkafi, H.; Secher, T.; Shima, A.; Shtanko, O.; Sakurai, Y.; Cojean, S.; Pomel, S.; Lievin-Le Moal, V.; Leignel, V.; Herweg, J. A.; Fischer, A.; Johannes, L.; Harrison, K.; Beard, P. M.; Clayette, P.; Le Grand, R.; Rayner, J. O.; Rudel, T.; Vacus, J.; Loiseau, P. M.; Davey, R. A.; Oswald, E.; Cintrat, J. C.; Barbier, J.; Gillet, D., Inhibitors of retrograde trafficking active against ricin and Shiga toxins also protect cells from several viruses, Leishmania and Chlamydiales. *Chem Biol Interact* **2017**, 267, 96-103.
43. Dai, W.; Wu, Y.; Bi, J.; Lu, X.; Hou, A.; Zhou, Y.; Sun, B.; Kong, W.; Barbier, J.; Cintrat, J. C.; Gao, F.; Gillet, D.; Su, W.; Jiang, C., Antiviral effects of Retro-2(cycl) and Retro-2.1 against Enterovirus 71 in vitro and in vivo. *Antiviral Res* **2017**, 144, 311-321.
44. Dai, W.; Wu, Y.; Bi, J.; Wang, J.; Wang, S.; Kong, W.; Barbier, J.; Cintrat, J. C.; Gao, F.; Jiang, Z.; Gillet, D.; Su, W.; Jiang, C., Antiviral Effect of Retro-2.1 against Herpes Simplex Virus Type 2 In Vitro. *J Microbiol Biotechnol* **2018**, 28, (6), 849-859.
45. Vinck, R.; Nguyen, L. A.; Munier, M.; Caramelle, L.; Karpman, D.; Barbier, J.; Pruvost, A.; Cintrat, J. C.; Gillet, D., In Vivo Sustained Release of the Retrograde Transport Inhibitor Retro-2.1 Formulated in a Thermosensitive Hydrogel. *Int J Mol Sci* **2022**, 23, (23).
46. Hotard, A. L.; Shaikh, F. Y.; Lee, S.; Yan, D.; Teng, M. N.; Plemper, R. K.; Crowe, J. E., Jr.; Moore, M. L., A stabilized respiratory syncytial virus reverse genetics system amenable to recombination-mediated mutagenesis. *Virology* **2012**, 434, (1), 129-36.
47. Galloux, M.; Tarus, B.; Blazevic, I.; Fix, J.; Duquerroy, S.; Eleouet, J. F., Characterization of a viral phosphoprotein binding site on the surface of the respiratory syncytial nucleoprotein. *Journal of virology* **2012**, 86, (16), 8375-87.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.