



HAL
open science

DNA-based studies and genetic diversity indicator assessments are complementary approaches to conserving evolutionary potential

Sean Hoban, Ivan Paz-Vinas, Robyn E Shaw, Luis Castillo-Reina, Jessica M D A Silva, J. Andrew Dewoody, Robert Ekblom, Ancuta Fedorca, Brenna R Forester, W. Chris Funk, et al.

► To cite this version:

Sean Hoban, Ivan Paz-Vinas, Robyn E Shaw, Luis Castillo-Reina, Jessica M D A Silva, et al.. DNA-based studies and genetic diversity indicator assessments are complementary approaches to conserving evolutionary potential. *Conservation Genetics*, 2024, 25, pp.1147-1153. 10.1007/s10592-024-01632-8 . hal-04684879v1

HAL Id: hal-04684879

<https://hal.inrae.fr/hal-04684879v1>

Submitted on 3 Sep 2024 (v1), last revised 6 Dec 2024 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



DNA-based studies and genetic diversity indicator assessments are complementary approaches to conserving evolutionary potential

Sean Hoban^{1,2,3,4,7} · Ivan Paz-Vinas^{2,4,8} · Robyn E. Shaw^{1,3,9} · Luis Castillo-Reina¹⁰ · Jessica M. D. A. Silva^{1,2,3,4,11,12} · J. Andrew DeWoody^{1,4,13} · Robert Ekblom^{1,2,4,14} · Ancuta Fedorca^{1,2,4,15,16} · Brenna R. Forester¹⁷ · W. Chris Funk^{1,3,4,5,18,19} · Julia C. Geue^{1,4,20} · Myriam Heuertz^{2,21} · Peter M. Hollingsworth²² · Alice C. Hughes²³ · Margaret E. Hunter^{1,3,4,24} · Christina Hvilson^{1,2,25} · Fumiko Ishihama²⁶ · Rebecca Jordan^{4,27} · Belma Kalamujić Stroil^{2,3,28} · Francine Kershaw^{3,4,29} · Colin K. Khoury³⁰ · Viktoria Köppä^{2,31} · Linda Laikre^{1,2,3,4,6,31} · Anna J. Macdonald^{1,3,4,32} · Alicia Mastretta-Yanes^{3,4,6,33,34} · Mariah H. Meek^{1,3,35,36} · Joachim Mergeay^{1,2,4,37,38} · Katie L. Millette^{4,6,39} · David O'Brien^{3,22,40} · Victor J. Rincón-Parra⁴¹ · M. Alejandra Rodríguez-Morales⁴² · Meredith C. Schuman^{6,43,44} · Gernot Segelbacher^{1,2,3,4,45} · Paul Sunnucks^{1,46} · Rebecca S. Taylor^{4,47} · Henrik Thurfjell^{2,48} · Cristiano Vernesi^{1,2,3,4,6,49,50} · Catherine E. Grueber^{1,3,4,51}

Received: 20 December 2023 / Accepted: 5 August 2024
© The Author(s) 2024

Abstract

Genetic diversity is essential for maintaining healthy populations and ecosystems. Several approaches have recently been developed to evaluate population genetic trends without necessarily collecting new genetic data. Such “genetic diversity indicators” enable rapid, large-scale evaluation across dozens to thousands of species. Empirical genetic studies, when available, provide detailed information that is important for management, such as estimates of gene flow, inbreeding, genetic erosion and adaptation. In this article, we argue that the development and advancement of genetic diversity indicators is a complementary approach to genetic studies in conservation biology, but not a substitute. Genetic diversity indicators and empirical genetic data can provide different information for conserving genetic diversity. Genetic diversity indicators enable affordable tracking, reporting, prioritization and communication, although, being proxies, do not provide comprehensive evaluation of the genetic status of a species. Conversely, genetic methods offer detailed analysis of the genetic status of a given species or population, although they remain challenging to implement for most species globally, given current capacity and resourcing. We conclude that indicators and genetic studies are both important for genetic conservation actions and recommend they be used in combination for conserving and monitoring genetic diversity.

Keywords Biodiversity monitoring · Genomics · Population genetics · Prioritisation

Introduction

Several approaches have recently been developed and used to efficiently and rapidly evaluate the genetic status and trends of populations and species, in the absence of collecting new genetic data (in this article we use “genetic data” and “genetic studies” to mean molecular techniques to examine the DNA or similar studies of inherited variation, e.g. DNA sequencing). Building on decades of prior work (Wright 1943; Frankham 1995; Manel et al. 2003; Hamann et al. 2005; Hanson et al. 2017), these approaches leverage population genetic theory and proxy-based geographic and

demographic data to approximate genetic status and trends. For example, two indicators were developed and adopted as part of the United Nations Convention on Biological Diversity (CBD) Global Biodiversity Framework (GBF): 1) the proportion of populations large enough to maintain genetic diversity (i.e., effective population size, N_e , > 500), and 2) the proportion of populations maintained (Laikre et al. 2020; Hoban et al. 2020, 2023a). These indicators are both valuable because they reflect different aspects of genetic diversity, and one or both may be derived from existing ecological monitoring data, if available (Mastretta-Yanes et al. 2024a). Other simple and policy-relevant genetic indicators have been developed, such as a multi-dimensional genetic health index (Kriesner et al. 2020), genetic scorecards for

Extended author information available on the last page of the article

high-profile species that document possible genetic threats and ongoing actions (Hollingsworth et al. 2020; O'Brien et al. 2022), and indicators of the sufficiency of protection (i.e., safeguarding) of genetic diversity across species' ranges (Khoury et al. 2019, 2020). The latter two are also indicators under the GBF. All of these indicators can use existing non-genetic data to varying degrees, along with summaries of previous genetic datasets when available, to assess the genetic status, trends, and/or effectiveness of conservation actions for species and populations, without the collection of new genetic (e.g. DNA-based) information. The use of indicators therefore allows rapid evaluation and comparison among dozens to thousands of species using simple, repeatable and accessible metrics with available data.

Here, we aim to emphasize that the development and advancement of such indicators complement genetic studies, that generate and analyse genetic information (including "genomic" approaches e.g. high throughput DNA sequence data; Fig. 1), and do not reduce the pressing need for studies that collect genetic data. When genetic data are feasible to obtain, they can be informative for specific cases, while also contributing to the growing evidence-base that underlies

pragmatic indicators. Genetic diversity indicators and genetic studies each provide different types of information for different uses, and *both are vital for informing the conservation of genetic diversity*. Indicators are especially useful for rapid, large-scale assessments when genetic data are not comprehensively available, and support policymakers and other stakeholders in prioritizing management among many species and populations using a common metric. In turn, these data can help direct resources towards species or populations in greatest need of more in-depth investigation, and support communicating to the public about genetic concepts and the urgency of genetic conservation (see Table 1).

Genetic diversity indicators

The genetic indicators included in the CBD GBF were originally designed to enable tracking of genetic status across many species and monitoring change over time, especially for the many species without genetic data, within the context of policy and management. One of the weaknesses of the previous CBD Aichi Biodiversity Targets was a lack of

Fig. 1 Genetic studies (typically using DNA data) and genetic indicator assessments (based on new and existing diverse datasets, such as ecological surveys) provide complementary and distinct information for guiding the genetic management of biodiversity




Table 1 Multiple assessments of genetic diversity indicators across moderate to large numbers of species have been conducted and have generated policy or management guidance, quickly, by using existing data. Genetic data and proxies of genetic diversity were integrated to varying degrees in each of these analyses

Number of species assessed	Conclusions and use for policy and management	Citation
909	Many species retain a majority of their populations, but most populations are below the threshold of $N_e > 500$ and may be in danger of genetic diversity loss. The indicator highlights populations and species in need of active intervention	Mastretta-Yanes, da Silva, et al. (2024a)
7,336 and 4,470	In a first-pass evaluation of genetic indicators in Sweden, 60% of 7,336 species potentially have populations with census size $N_c > 5,000$, while 91% of 4,470 species have lost some subpopulations or distribution during the last 100 years	Thurfjell et al. (2022)
594	Crop wild relative genetic diversity has not been well conserved ex situ or in situ, and most species require urgent action. The indicator provides a quantitative measure of progress on this action (geographic or ecological range conserved)	Khoury et al. (2020)
26	Assessment of several criteria show that 14 species are likely at negligible genetic risk, eight species at moderate risk (with effective mitigation in place for five of these), and four species are likely at risk of severe genetic problems	Hollingsworth et al. (2020)
1,148	Approximately 15–20% of assessed species are “likely to benefit from... genetic rescue or targeted gene flow/genetic augmentation. Over half... would benefit from re-establishing new populations”	Kriesner et al. (2020)
1,321	Documents that genetic diversity (GD) is reduced in threatened species and proposes a straightforward framework to rank species based on publicly available GD data and time to expected loss of GD due to small populations	Willoughby et al. (2015)
83	Mean individual zygosity (i.e., heterozygosity and autozygosity) estimated from publicly available resequencing data can be used as a key criterion in a novel analytical framework to help rank species by conservation category	Jeon et al. (2024)

quantitative, reliable indicators (Butchart et al. 2016; Xu et al. 2021), and so these were required in the more-recent CBD GBF. Although genetic diversity has been overlooked in conservation policy in the past (Laikre et al. 2010), genetic diversity indicators enabled this important component of biodiversity to be included in the GBF (Hoban et al. 2023a; Carroll et al. 2023). Genetic indicators have some notable limitations, such as focusing primarily on only two aspects of genetic health (amount of genetic drift and numbers of putatively locally adapted populations), utilizing thresholds (such as $N_e > 500$) rather than continuous metrics, and relying on assumptions such as population-level adaptation and a universal estimate of the N_e/N_c ratio. However, these indicators enable biodiversity managers to quantify likely genetic change for the thousands of species for which detailed genetic studies are not currently available or likely to be so in the near future, and manage many populations that may need intervention (Hoban et al. 2023b, 2024; Mastretta-Yanes et al. 2024a). Indicators can thus enable the following key outcomes: tracking change over hundreds of species and across the world; reporting such change under policy frameworks including the CBD and regional and national frameworks where hundreds or thousands of species (and often their distinct populations) are protected and assessed (e.g., U.S. Endangered Species Act, Canadian Species at Risk Act, European Habitats Directive, European Nature Restoration Law, Mexican SEMARNAT NOM-059, South African National Environmental Management:

Biodiversity Act); prioritizing—based on a common metric—species that have the most urgent conservation needs to remedy genetic diversity loss; and facilitating communication on genetic issues to non-geneticists, the public and policymakers. Further research on genetic diversity indicators, including complementary studies with genetic data, will help improve indicators into the future.

Genetic data, notably from DNA-based studies

Genetic data have been used for decades to study the genetic “health” of species in detail, and the processes impacting the conservation of genetic diversity (Frankham 2010). Genetic data can reveal diverse and detailed information, such as understanding hybridization, population distinctions or barriers to gene flow, ancient population size and size changes (and thus providing context for current size), assessing inbreeding at the genomic level, and quantifying genetic load, etc. (Allendorf et al. 2010, 2022; DeWoody et al. 2021). Genetic data can also detect gene flow between domesticated species, their wild relatives, and/or genetically modified organisms or improved varieties (Rojas-Barrera et al. 2019). The use of recently defined Genetic Essential Biodiversity Variables—standardized genetic summary metrics that focus on core attributes of genetic composition—can facilitate comparison among genetic monitoring

studies (Hoban et al. 2022). Genetic data can sometimes also provide information on the type and extent of local adaptation and maladaptation (Hoban et al. 2016; Capblancq et al. 2020). Combining genetic data with simulations can predict likely trajectories of populations and their fitness and diversity in response to management (Hoban 2014; Robinson et al. 2022) including time until a certain threshold of genetic erosion occurs (Hoban et al. 2014; Willoughby et al. 2015). This kind of information is often vital to on-the-ground management decisions as well as informing policy. As a few of the many examples, genetic data contributed to decisions about successful genetic rescue for helmeted honeyeaters in Australia (Harrisson et al. 2016; Pavlova et al. 2023), thresholds for hybrid management in bontebok and blesbok in South Africa (van Wyk et al. 2017), and red wolf management in the USA (Gese et al. 2015) (see also Bertola et al. 2024). Nonetheless, recent surveys show that a very small portion of species have any genetic data available (perhaps 1 to 5%) and an even smaller portion (less than 0.5%, even in wealthy nations) have temporal genetic monitoring (Torres-Florez et al. 2018; Posledovich et al. 2021; Thurffjell et al. 2022; Pearman et al. 2024; Paz-Vinas et al. 2023).

Synthesis

As discussed above, genetic indicators and genetic data collection methods both provide data necessary to monitor and protect genetic diversity, albeit in different ways. They are complementary strategies that should be deployed together for conserving and managing biodiversity. While genetic data can be used to generate detailed analyses of a given entity's genetic status, genetic data alone have not been sufficient to meet all conservation genetic needs, including prioritizing among many species, and supporting affordable and rapid decision-making. Conversely, while genetic indicators enable broad-scale evaluation of many species and facilitate affordable and accessible tracking, reporting, prioritization and communication, they are proxies and do not provide a full assessment of genetic composition and genetic change in a population or species, nor answer applied questions such as whether gene flow is occurring between locations. Genetic indicators (e.g., the proportion of large populations or the proportion of extant populations relative to extirpated populations; Hoban et al. 2023a, b) can be compiled for one species in just a few hours if field-based ecological or demographic data have already been generated, including via citizen scientists or local communities (Mastretta-Yanes et al. 2024a, b). In contrast, conducting an empirical genetic or genomic study commonly takes months, and substantial infrastructure, expertise, samples, and funding (Kriesner et al. 2020; Mastretta-Yanes et al. 2024a).

The complementary use of indicators and detailed genetic studies may be analogous to species threat assessments (e.g. IUCN Red List) and detailed population viability models (PVMs). Like genetic studies, PVMs offer a high level of precision and insight, although they typically require detailed information that is slow and expensive to gather, making their application feasible for relatively few species. In contrast, species threat assessments, like indicators, use pragmatic thresholds of a few simple criteria to characterize a species' status, enabling them to be done for many species, and indicating where more detailed studies are required. Like broad-scale threat assessments, genetic diversity indicators can point to potential issues, raising alarm in situations where concern is warranted, which can promote collection of more data, generate specific actions, and encourage monitoring.

Genetic indicators can be integrated with genetic data to provide a comprehensive genetic management approach (Fig. 1). For example, indicators may reveal which species or populations are most likely in need of genetic management, and genetic data can then be generated for those species to provide management-relevant information such as migration rates, temporal changes in N_e , and inbreeding. Meanwhile, DNA-based genetic studies of wild populations can provide better understanding of the biological processes underpinning genetic change, such as N_e dynamics, determinants of the N_e/N_c ratio, undesired effects of gene flow or artificial selection, and the needs of populations for long-term persistence. Such genetic knowledge can then be directly used to improve indicator calculation, such as by clarifying population genetic structure of a species or applying a tailored N_e/N_c ratio, leading to increasingly robust, empirically supported indicator estimates.

Conclusion

In summary, the development of simple indicator approaches enables tracking and reporting of genetic information at large scales in all countries. While genetic studies provide irreplaceable information often useful for on-the-ground management, especially for issues like hybridization, inbreeding, fitness, and adaptation, they are challenging to scale up and therefore should be complemented by less expensive and more scalable approaches (e.g., indicators). Both are important and useful for genetic conservation action and should be used to complement each other. We conclude that a holistic approach at a country-level through the development of both genetic indicators and genetic studies can rapidly improve the monitoring and conservation of genetic diversity.

Acknowledgements The findings and conclusions in this article are those of the authors and do not necessarily represent the views

of the US Fish and Wildlife Service. We acknowledge the efforts of the Coalition for Conservation Genetics and its component members, G-BIKE, the IUCN Conservation Genetics Specialist Group, the GEO BON Genetic Composition Working Group and the SCB Conservation Genetics Working Group, whose members have diligently worked to advance the science and practice of DNA-based studies and genetic diversity indicators, usually on a volunteer basis. We are grateful for this community. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Author contributions SH drafted the initial manuscript, CG helped revise key points, RS and I P–V drafted the Figure, and all authors contributed edits, comments, or dialogue on the manuscript topics. All authors approved the final manuscript.

Funding Open Access funding enabled and organized by CAUL and its Member Institutions. We acknowledge support of the Biodiversa+ project GINAMO and the COST Action GENOA (CA23121). LL acknowledges Formas [2020-01290], Swedish Research Council [2019-05503], Swedish Environmental Protection Agency, Swedish Agency for Marine and Water Management.

Data availability No datasets were generated or analysed during the current study.

Declarations

Competing interests Sean Hoban is an editor for Conservation Genetics; there are no other conflicts of interest to declare.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Allendorf FW, Hohenlohe PA, Luikart G (2010) Genomics and the future of conservation genetics. *Nat Rev Genet* 11:697–709
- Allendorf FW, Chris Funk W, Aitken SN et al (2022) Conservation and the Genomics of Populations. Oxford University Press
- Bertola LD, Brüniche-Olsen A, Kershaw F et al (2024) A pragmatic approach for integrating molecular tools into biodiversity conservation. *Conserv Sci Pract* 6:e13053
- Butchart SHM, Di Marco M, Watson JEM (2016) Formulating smart commitments on biodiversity: Lessons from the Aichi targets. *Conserv Lett* 9:457–468
- Capblancq T, Fitzpatrick MC, Bay RA et al (2020) Genomic prediction of (Mal)adaptation across current and future climatic landscapes. *Annu Rev Ecol Evol Syst* 51:245–269
- Carroll C, Hoban S, Ray JC (2023) Lessons from COP15 on effective scientific engagement in biodiversity policy processes. *Conserv Biol*. <https://doi.org/10.1111/cobi.14192>
- DeWoody JA, Harder AM, Mathur S, Willoughby JR (2021) The long-standing significance of genetic diversity in conservation. *Mol Ecol* 30:4147–4154
- Frankham R (1995) Effective population size/adult population size ratios in wildlife: a review. *Genet Res* 66:95–107
- Frankham R (2010) Where are we in conservation genetics and where do we need to go? *Conserv Genet* 11:661–663
- Gese EM, Knowlton FF, Adams JR et al (2015) Managing hybridization of a recovering endangered species: The red wolf *Canis rufus* as a case study. *Curr Zool* 61:191–205
- Hamann A, Smets P, Yanchuk AD, Aitken SN (2005) An ecogeographic framework for in situ conservation of forest trees in British Columbia. *Can J Res* 35:2553–2561
- Hanson JO, Rhodes JR, Riginos C, Fuller RA (2017) Environmental and geographic variables are effective surrogates for genetic variation in conservation planning. *Proc Natl Acad Sci U S A* 114:12755–12760
- Harrison KA, Pavlova A, da Silva AG et al (2016) Scope for genetic rescue of an endangered subspecies through re-establishing natural gene flow with another subspecies. *Mol Ecol* 25:1242–1258
- Hoban S (2014) An overview of the utility of population simulation software in molecular ecology. *Mol Ecol* 23:2383–2401
- Hoban S, Arntzen JA, Bruford MW et al (2014) Comparative evaluation of potential indicators and temporal sampling protocols for monitoring genetic erosion. *Evol Appl* 7:984–998
- Hoban S, Kelley JL, Lotterhos KE et al (2016) Finding the genomic basis of local adaptation: pitfalls, practical solutions, and future directions. *Am Nat* 188:379–397
- Hoban S, Bruford M, D'Urban Jackson J et al (2020) Genetic diversity targets and indicators in the CBD post-2020 Global Biodiversity Framework must be improved. *Biol Conserv* 248:108654
- Hoban S, Archer FI, Bertola LD et al (2022) Global genetic diversity status and trends: towards a suite of Essential Biodiversity Variables (EBVs) for genetic composition. *Biol Rev Camb Philos Soc* 97:1511–1538
- Hoban S, Bruford MW, da Silva JM et al (2023a) Genetic diversity goals and targets have improved, but remain insufficient for clear implementation of the post-2020 global biodiversity framework. *Conserv Genet* 24:181–191
- Hoban S, da Silva JM, Mastretta-Yanes A et al (2023b) Monitoring status and trends in genetic diversity for the Convention on Biological Diversity: An ongoing assessment of genetic indicators in nine countries. *Conserv Lett* 16:e12953
- Hoban S, da Silva J, Hughes A et al (2024) Too simple, too complex, or just right? Advantages, challenges and resolutions for indicators of genetic diversity. *Bioscience* 74:269–280
- Hollingsworth PM, O'Brien D, Ennos RA, et al. (2020) Scotland's biodiversity progress to 2020 aichi targets: Aichi target 13-Genetic Diversity Maintained—supplementary report 2020. Scottish Natural Heritage, Inverness
- Jeon JY, Black A, Heenkenda E, Mularo A, Lamka G, Janjua S, Brüniche-Olsen A, Bickham JW, Willoughby J, DeWoody JA (2024) Genomic diversity as a key conservation criterion: proof-of-concept from mammalian whole-genome resequencing data. *Evolutionary Applications*, in press.
- Khoury CK, Amariles D, Soto JS et al (2019) Comprehensiveness of conservation of useful wild plants: An operational indicator for biodiversity and sustainable development targets. *Ecol Indic* 98:420–429
- Khoury CK, Carver D, Greene SL et al (2020) Crop wild relatives of the United States require urgent conservation action. *Proc Natl Acad Sci U S A* 117:33351–33357
- Kriesner P, Weeks A, Razeng E, Sunnucks P (2020) Assessing genetic risks to Victorian flora and fauna. Department of Environment, Land, Water and Planning

- Laikre L, Allendorf FW, Aroner LC et al (2010) Neglect of genetic diversity in implementation of the convention of biological diversity. *Conserv Biol* 24:86–88
- Laikre L, Hoban S, Bruford MW et al (2020) Post-2020 goals overlook genetic diversity. *Science* 367:1083–1085
- Manel S, Schwartz MK, Luikart G, Taberlet P (2003) Landscape genetics: combining landscape ecology and population genetics. *Trends Ecol Evol* 18:189–197
- Mastretta-Yanes A, da Silva J, Grueber CE et al (2024a) Multinational evaluation of genetic diversity indicators for the Kunming-Montreal Global Biodiversity Monitoring Framework. *Ecol Lett* 27:e14461
- Mastretta-Yanes A, Suárez S, Jordan R, Hoban S, da Silva JM, Castillo-Reina L, Heuertz M, Ishihama F, Köppä V, Laikre L, MacDonald AJ, Mergeay J, Paz-Vinas I, Segelbacher G, Knapps A, Rakoczy H, Weiler A, Atsaves A, Cullmann K, Bagnato S, Forester BR (2024b) Guideline materials and documentation for the genetic diversity indicators of the Monitoring Framework for the Kunming-Montreal Global Biodiversity Framework. *Biodiversity Informatics* 18. <https://doi.org/10.17161/bi.v18i.22332>
- O'Brien D, Laikre L, Hoban S et al (2022) Bringing together approaches to reporting on within species genetic diversity. *J Appl Ecol* 59:2227–2233
- Pavlova A, Petrovic S, Harrison KA et al (2023) Benefits of genetic rescue of a critically endangered subspecies from another subspecies outweigh risks: Results of captive breeding trials. *Biol Conserv* 284:110203
- Paz-Vinas I, Vandergast A, Schmidt C, et al. (2023) Uneven genetic data limits biodiversity assessments in protected areas globally. *EcoEvoRxiv* <https://doi.org/10.32942/XZ2C84>
- Pearman PB, Broennimann O, Albayrak T et al (2024) Conserving genetic diversity during climate change: Niche marginality and discrepant monitoring capacity in Europe. *Nat Ecol Evol* 8:267–281
- Posledovich D, Ekblom R, Laikre L (2021) Mapping and monitoring genetic diversity in Sweden: A proposal for species, methods and costs. Swedish Environmental Protection Agency.
- Robinson JA, Kyriazis CC, Nigenda-Morales SF et al (2022) The critically endangered vaquita is not doomed to extinction by inbreeding depression. *Science* 376:635–639
- Rojas-Barrera IC, Wegier A, Sánchez González JDJ et al (2019) Contemporary evolution of maize landraces and their wild relatives influenced by gene flow with modern maize varieties. *Proc Natl Acad Sci U S A* 116:21302–21311
- Thurfjell H, Laikre L, Ekblom R et al (2022) Practical application of indicators for genetic diversity in CBD post-2020 Global Biodiversity Framework implementation. *Ecol Indic* 142:109167
- Torres-Florez JP, Johnson WE, Nery MF et al (2018) The coming of age of conservation genetics in Latin America: what has been achieved and what needs to be done. *Conserv Genet* 19:1–15
- van Wyk AM, Dalton DL, Hoban S et al (2017) Quantitative evaluation of hybridization and the impact on biodiversity conservation. *Ecol Evol* 7:320–330
- Willoughby JR, Sundaram M, Wijayawardena BK et al (2015) The reduction of genetic diversity in threatened vertebrates and new recommendations regarding IUCN conservation rankings. *Biol Conserv* 191:495–503
- Wright S (1943) Isolation by distance. *Genetics* 28:114–138
- Xu H, Cao Y, Yu D et al (2021) Ensuring effective implementation of the post-2020 global biodiversity targets. *Nat Ecol Evol* 5:411–418

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors and Affiliations

Sean Hoban^{1,2,3,4,7} · Ivan Paz-Vinas^{2,4,8} · Robyn E. Shaw^{1,3,9} · Luis Castillo-Reina¹⁰ · Jessica M. D. A. Silva^{1,2,3,4,11,12} · J. Andrew DeWoody^{1,4,13} · Robert Ekblom^{1,2,4,14} · Ancuta Fedorca^{1,2,4,15,16} · Brenna R. Forester¹⁷ · W. Chris Funk^{1,3,4,5,18,19} · Julia C. Geue^{1,4,20} · Myriam Heuertz^{2,21} · Peter M. Hollingsworth²² · Alice C. Hughes²³ · Margaret E. Hunter^{1,3,4,24} · Christina Hvilson^{1,2,25} · Fumiko Ishihama²⁶ · Rebecca Jordan^{4,27} · Belma Kalamujić Strojil^{2,3,28} · Francine Kershaw^{3,4,29} · Colin K. Khoury³⁰ · Viktoria Köppä^{2,31} · Linda Laikre^{1,2,3,4,6,31} · Anna J. Macdonald^{1,3,4,32} · Alicia Mastretta-Yanes^{3,4,6,33,34} · Mariah H. Meek^{1,3,35,36} · Joachim Mergeay^{1,2,4,37,38} · Katie L. Millette^{4,6,39} · David O'Brien^{3,22,40} · Victor J. Rincón-Parra⁴¹ · M. Alejandra Rodríguez-Morales⁴² · Meredith C. Schuman^{6,43,44} · Gernot Segelbacher^{1,2,3,4,45} · Paul Sunnucks^{1,46} · Rebecca S. Taylor^{4,47} · Henrik Thurfjell^{2,48} · Cristiano Vernesi^{1,2,3,4,6,49,50} · Catherine E. Grueber^{1,3,4,51}

✉ Catherine E. Grueber
catherine.grueber@sydney.edu.au

- ¹ International Union for Conservation of Nature (IUCN) Conservation Genetics Specialist Group (CGSG), Gland, Switzerland
- ² Genomic Biodiversity Knowledge for Resilient Ecosystems (G-BiKE), San Michele All'Adige, Italy
- ³ Coalition for Conservation Genetics (CCG), Gland, Switzerland
- ⁴ The Group On Earth Observations Biodiversity Observation Network (GEO BON), Montreal, Canada

- ⁵ The Society for Conservation Biology (SCB) Conservation Genetics Working Group (CGWG), Washington, USA
- ⁶ International Space Science Institute International Team "Genes From Space", Bern, Switzerland
- ⁷ The Center for Tree Science, The Morton Arboretum, Lisle 60187, USA
- ⁸ CNRS, ENTPE, UMR5023 LEHNA, Université Claude Bernard Lyon 1, F-69622 Villeurbanne, France
- ⁹ Centre for Conservation Ecology and Genomics, University of Canberra, Bruce, Canberra, ACT 2617, Australia

- ¹⁰ Department of Biology, Faculty of Science, KU Leuven, Louvain, Belgium
- ¹¹ South African National Biodiversity Institute, Kirstenbosch Research Centre, Private Bag X7, Claremont 7735, Cape Town, South Africa
- ¹² Centre for Ecological Genomics and Wildlife Conservation, Department of Zoology, University of Johannesburg, Johannesburg, Auckland Park 2006, South Africa
- ¹³ Departments of Forestry & Natural Resources and Biological Sciences, Purdue University, West Lafayette, IN, USA
- ¹⁴ Wildlife Analysis Unit, Swedish Environmental Protection Agency, 106 48 Stockholm, Sweden
- ¹⁵ Department of Wildlife, National Institute for Research and Development in Forestry “Marin Dracea”, Closca 13, Brasov, Romania
- ¹⁶ Department of Silviculture, Faculty of Silviculture and Forest Engineering, Transilvania University of Brasov, Beethoven Lane 1, Brasov, Romania
- ¹⁷ U.S. Fish and Wildlife Service, Fort Collins, CO 80525, USA
- ¹⁸ Graduate Degree Program in Ecology, Colorado State University, Fort Collins, CO, USA
- ¹⁹ Department of Biology, Colorado State University, Fort Collins, CO, USA
- ²⁰ Biology Department, Trent University, Peterborough, Canada
- ²¹ Biogeco, INRAE, Univ. Bordeaux, 33610 Cestas, France
- ²² Royal Botanic Garden Edinburgh, Edinburgh EH3 5LR, UK
- ²³ School of Biological Sciences, University of Hong Kong, Pok Fu Lam, China
- ²⁴ U.S. Geological Survey, Gainesville, FL 32605, USA
- ²⁵ Copenhagen Zoo, Frederiksberg 2000, Denmark
- ²⁶ Biodiversity Division, National Institute for Environmental Studies, Onogawa 16-2, Tsukuba, Ibaraki, Japan
- ²⁷ CSIRO Environment, Sandy Bay, TAS 7005, Australia
- ²⁸ University of Sarajevo-Institute for Genetic Engineering and Biotechnology, 71000 Sarajevo, Bosnia and Herzegovina
- ²⁹ Natural Resources Defense Council, New York, NY 10011, USA
- ³⁰ San Diego Botanic Garden, 300 Quail Gardens Dr, Encinitas, CA 92024-2707, USA
- ³¹ Department of Zoology, Stockholm University, 10691 Stockholm, Sweden
- ³² Australian Antarctic Division, Department of Climate Change, Energy, The Environment and Water, Kingston, TAS 7050, Australia
- ³³ Consejo Nacional de Humanidades Ciencias y Tecnología (CONAHCYT), Avenida Insurgentes Sur 1582, Crédito Constructor, Benito Juárez, Ciudad de México C.P. 03940, México
- ³⁴ Departamento de Ecología de la Biodiversidad, Instituto de Ecología, Universidad Nacional Autónoma de México, Ciudad Universitaria, Coyoacán, Ciudad de México 04500, México
- ³⁵ The Wilderness Society, Bozeman, MT, USA
- ³⁶ Department of Integrative Biology, Michigan State University, East Lansing, MI, USA
- ³⁷ Research Institute for Nature and Forest, Gaverstraat 4, 9500 Geraardsbergen, Belgium
- ³⁸ Ecology, Evolution and Biodiversity Conservation, KU Leuven, Charles Deberiotstraat 32, Box 2439, 3000 Leuven, Belgium
- ³⁹ GEO BON Secretariat, Department of Biology, McGill University, Montreal, QC H3A 1B1, Canada
- ⁴⁰ NatureScot, Inverness IV3 8NW, UK
- ⁴¹ Alexander Von Humboldt Biological Resources Research Institute, Bogotá, Colombia
- ⁴² Department of Biology, Faculty of Science, Pontificia Universidad Javeriana, Bogotá D.C. Colombia
- ⁴³ Remote Sensing Laboratories, Department of Geography, University of Zurich, Zurich, Switzerland
- ⁴⁴ Department of Chemistry, University of Zurich, Zurich, Switzerland
- ⁴⁵ Chair of Wildlife Ecology and Management, University Freiburg, Breisgau, Germany
- ⁴⁶ School of Biological Sciences, Monash University, Melbourne, Australia
- ⁴⁷ Landscape Science and Technology Division, Environment and Climate Change Canada, Ottawa, Canada
- ⁴⁸ Swedish Species Information Center, Swedish University of Agricultural Sciences, Box 7070, 75007 Uppsala, Sweden
- ⁴⁹ Research and Innovation Centre, Fondazione Edmund Mach, S. Michele all'Adige, Italy
- ⁵⁰ National Biodiversity Future Centre, 90133 Palermo, Italy
- ⁵¹ School of Life and Environmental Sciences, Faculty of Science, The University of Sydney, Camperdown NSW 2006, Australia