



HAL
open science

The phosphoproteome signature of *Listeria monocytogenes* dormancy

Alessandro Pagliuso, Lorie Pelmont, Lydie Oliveira Correia, Filipe Carvalho,
Céline Henry

► **To cite this version:**

Alessandro Pagliuso, Lorie Pelmont, Lydie Oliveira Correia, Filipe Carvalho, Céline Henry. The phosphoproteome signature of *Listeria monocytogenes* dormancy. PTMBact2024, May 2024, Rouen (FR), France. hal-04691873

HAL Id: hal-04691873

<https://hal.inrae.fr/hal-04691873v1>

Submitted on 9 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Background

Several bacteria can adapt to stress by entering a viable but non-culturable (VBNC) state (1), where they remain alive and metabolically active but cannot grow in standard culture media (Fig. 1).

Listeria monocytogenes (Lm), a deadly foodborne pathogen for immunocompromised or elderly individuals and newborns, can survive in various stressful environments by transitioning to a VBNC state (2, 3). However, the molecular mechanisms governing this transition are not well understood.

We have recently shown that during starvation in mineral water, Lm progressively transitions to a VBNC state (Fig. 2A-B) (4). Phase contrast microscopy observation revealed that Lm rods become progressively shorter to transform into coccoid forms after 2-3 weeks (Fig. 2C).

Cryo-electron tomography (Fig. 2D) and peptidoglycan profiling (not shown) indicated that this transformation is due to a shedding process of the peptidoglycan, resulting in cell wall-deficient bacteria.




Figure 1. During prolonged stressful conditions, a proportion of viable bacteria will enter into a VBNC state, under which they cannot develop into colonies on agar medium, but cellular metabolic activity is retained. When provided with favorable conditions, VBNC cells can resuscitate to the culturable state.





Figure 2. (A) Total, viable and culturable cell profiles of Lm in mineral water. (B) Fraction of bacterial population consisting of VBNC formed in mineral water. (C) Phase-contrast micrographs of Lm cells during transition into a VBNC state. Bacteria highlighted by white squares are shown enlarged in bottom panels. (D) Cryo-electron tomography of Lm cells sampled after 7 days of incubation in mineral water.



To understand the molecular events leading to the VBNC transition and cell wall loss, we analyzed the proteome of bacteria transitioning to a VBNC state (Fig 3).

Proteomic analysis identified up- and down-regulated proteins (red and blue, respectively) involved in cell wall metabolism and remodeling, cell metabolism (PTS systems), and cell signaling (two component systems). Since many of these proteins are known to be phosphorylated, we hypothesize that phosphorylation cascades regulate the VBNC transition.


Figure 3. Volcano plot showing the intensity fold change (in log₂) of proteins at day 14 (D14) and day 28 (D28) compared to day 0 (D0) on the x axis. The experiment was performed 4 times and a t-test was performed to calculate -log P values for each protein, indicated on the y axis

Objectives

Hypothesis:
Phosphorylation/dephosphorylation events regulate VBNC transition


- ✓ Characterize the phosphoproteome of Lm transitioning to the VBNC state
- ✓ Identify differentially-phosphorylated targets that might regulate VBNC transitions

Experimental pipeline



Preliminary results

To set up phosphopeptide enrichment we started from bacteria grown in BHI until the stationary phase or incubated in water for 1 or 2 days. We used the pipeline shown beside. Phosphopeptides were purified via TiO₂ beads.



Conclusion and perspectives

- ✓ These results show that our protocol is suitable to isolate phosphoproteins.
- ✓ We were able to identify 104 phosphoproteins and 130 phosphoislands.
- ✓ We are currently optimising our experimental pipeline with the aim to profile the Lm phosphoproteome during VBNC transition and identify molecular players and signalling pathways involved in this physiological state

References

- Pazos-Rojas LA, Cuellar-Sánchez A, Romero-Cerón AL, Rivera-Urbalejo A, Van Dillewijn P, Luna-Vital DA, Muñoz-Rojas J, Morales-García YE, Bustillos-Cristales MDR. The Viable but Non-Culturable (VBNC) State, a Poorly Explored Aspect of Beneficial Bacteria. *Microorganisms*. 2023 Dec 25;12(1):39. doi: 10.3390/microorganisms12010039. PMID: 38257865; PMCID: PMC10818521.
- Lotoux A, Milohanic E, Biernie H. The Viable But Non-Culturable State of *Listeria monocytogenes* in the One-Health Continuum. *Front Cell Infect Microbiol*. 2022 Mar 18;12:849915. doi: 10.3389/fcimb.2022.849915. PMID: 35372114; PMCID: PMC8974916.
- Kortebi M, Milohanic E, Mitchell G, Péchoux C, Prevost MC, Cossart P, Biernie H. *Listeria monocytogenes* switches from dissemination to persistence by adopting a vacuolar lifestyle in epithelial cells. *PLoS Pathog*. 2017 Nov 30;13(11):e1006734. doi: 10.1371/journal.ppat.1006734. PMID: 29190284; PMCID: PMC5708623.
- Filipe Carvalho, et al Diving into bacterial dormancy: emergence of osmotically stable wall-less forms in an aquatic environment, *Biorxiv*, doi: <https://doi.org/10.1101/2023.11.16.566987>