



**HAL**  
open science

## Multiperformance of slow-growing and dual-purpose strains in organic chicken production: learning from the PPILOW project

Claire Bonnefous, Cesare Castellini, Simona Mattioli, Sandrine Mignon-Grasteau, Julie M Collet, Laurence A. Guilloteau, Bertrand Méda, Théophane de Rauglaudre, Helen Pluschke, Petra Thobe, et al.

### ► To cite this version:

Claire Bonnefous, Cesare Castellini, Simona Mattioli, Sandrine Mignon-Grasteau, Julie M Collet, et al.. Multiperformance of slow-growing and dual-purpose strains in organic chicken production: learning from the PPILOW project. 75. Annual meeting of the european federation of animal science (EAAP), European Federation of Animal Science, Sep 2024, Florence, Italy. pp.471. hal-04692952

**HAL Id: hal-04692952**

**<https://hal.inrae.fr/hal-04692952v1>**

Submitted on 10 Sep 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Multiperformance of slow-growing and dual-purpose strains in organic chicken production: learning from the PPILOW project

*Bonnefous C., Castellini C., Mattioli S., Mignon-Grasteau S., Collet J., Guilloteau L.A., Méda B., de Rauglaudre T., Pluschke H., Thobe P., Werner D., Calandreau L., Guesdon V., Steinfeldt S., Germain K., Ravon L., Berri C., Lombard S., Reverchon M., Le Bihan-Duval E., **Collin A.***

INRAE



THÜNEN

JUNIA Grande école d'ingénieurs

AARHUS UNIVERSITY

acta LES INSTITUTS TECHNIQUES AGRICOLES

itab l'Institut de l'agriculture et de l'alimentation biologiques

SYSAAF

*The 75th EAAP Annual Meeting, 1/5 September 2024 - Florence, Italy*

# PPILOW What is multiperformance for broiler production?



## Economic performance

Zootechnical parameters  
Meat yield  
Costs and revenues  
...

## Animal Welfare performance

5 freedoms

Freedom from Hunger and Thirst  
Freedom from Discomfort  
Freedom from Pain, Injury or Disease  
Freedom to Express Normal Behavior  
Freedom from Fear and Distress

the Farm Animal Welfare Council, 1993

Satisfaction of physiological and behavioural needs,  
as well as expectations ANSES, 2018

## Animal health performance

Absence of disease  
Food safety  
Immunity and adaptive capacities  
...

## Environmental performance

Energy and water use  
Climate change  
Pollution  
...

## Societal performance

Work conditions  
Job perception  
Social involvement  
Meat quality and food security  
...

**MULTI-  
PERFORMANCE**

***Some indicators shared with those included in multicriteria sustainability assessments***

# Why do we consider multiperformance in organic broiler systems?



- Diversity of practices and breeds used with different performance and environmental impacts throughout Europe
- Still a need to improve animal welfare and limit mortality, in relation to the **outdoor access challenging the animals** and **ethical issues**

Objective of PPILOW: Identify, test and evaluate animal welfare-improving practices by taking into account environmental, economic and social impacts including human well-being  
*One Welfare concept (Garcia Pinillos et al., 2016) and Multiperformance*

# H2020 PPILOW partners and collaborators



**PPILOW**

Poultry and Pig Low-input and Organic production systems' Welfare

2019-2024




Coordination: **INRAE**

22 PPILOW Partners in 9 countries

9 National Practitioner Groups (NPG): 5 dedicated to poultry

[www.ppilow.eu](http://www.ppilow.eu)

- Research institute
- Technical Institute
- Citizen/Practitioner Association
- SME
- University/Higher education



# PPILOW Involvement of National Practitioner Groups

## Innovative breeding and rearing strategies



*Favouring positive behaviours, improving health and robustness*

*Avoiding feather pecking in non beak-trimmed hens and the elimination of layer male chicks*



## Broilers in organic free range systems (PPILOW WP6.1)

Free-range



Key request for European consumers : **Expression of species behaviours** (walking/running, foraging, social interactions...) - **welfare**

Interesting functions for the **agroecological transition** (nutrient inputs from plants and insects, closing nutrients cycles, biodiversity...)

Potential benefits are subject to the fact that **poultry use the outdoor space...** and are able to maintain **good performances, health and welfare** when exposed to biotic or abiotic stressors on the range

- **Variability of range use** between **individuals from different genetic lines** of broilers and **trade-offs between functions**

# Dual-purpose breeds in organic systems (PPILOW WP5)

National legislations and knowledge for practitioners and European policy makers:  
**Alternatives to the elimination of layer male chicks**

Layer strain

*Selection based on egg production, egg quality traits*

*Culling of layer male chicks*



FR: Article R214-17

- From 01/01/2023 : all hatcheries have to be equipped with operational material to avoid culling chick in coloured strains



DE: Article TierSchG Art. 1 § 4c

- From 01/01/2022 : makes it a punishable offence to kill a vertebrate animal "without reasonable cause" (unprofitability) or to cause it suffering and pain



## Two strategies developed in PPILOW


- Study of dual-purpose strains experimentally and on-farm

**Multiperformance data**, multicriteria analysis and business models

ACTA(ITAB)  
 Thuenen Institute  
 Aarhus University  
 INRAE  
 SYSAAF



Meat-type



Pure breed




Layer-type



Pluschke et al., EAAP 2024

© Photos / Pluschke

- Development of an on-ovo sexing tool:  
*Non invasive method*  
*As early as possible*





# PPILOW Multiperformance of organic broiler production in PPILOW?



## Economic performance

Zootechnical parameters /  
growth performance  
Meat yield  
Costs and revenues

## Animal Welfare

5 freedoms

Freedom from Hunger and Thirst  
Freedom from Discomfort  
Freedom from Pain, Injury or Disease  
Freedom to Express Normal Behavior  
Freedom from Fear and Distress

Satisfaction of physiological and behavioural needs,  
as well as expectations

Animal welfare assessments  
Range use

## Animal health

Absence of disease  
Food safety  
Immunity and adaptive capacities  
Bone health and redox status

## Environmental performance

Energy and water use  
Climate change  
Pollution

## Societal performance

Work conditions  
Job perception  
Social involvement  
Meat quality and food security  
Consumer preference


Multiperformance


# Multiperformance of broilers from different breeds and range use (France), WP6

3 strains: 1 per range; 750 animals per strain ; 50% males, 50% females



WP6




- 
- Welfare indicators
  - Performances and meat quality
  - Blood parameters

Bonnefous et al., 2023  
Collet et al., 2024

# Results - Variability of individual range use (WP6)

Evaluation of individual Range Use by the Distance Index from scan samplings (N=100 males per line)

Impact of range use variation within strains?



Selection :  
 25 animals with the lowest Final Distance Index = low-rangers  
 25 animals with the highest Final Distance Index = high-rangers

Age at slaughter (d) **86**  
 Cumulated time spent outdoor (hr) 108 419

**100**  
 296 741

**121**  
 212 699



Follow-up by RFID  
 Collet et al., 2024

Bonnefous et al., 2023




# Growth performance - Relationship between range use and **body weight**?

Lower BW predisposal factor to higher range use?  
HR birds already more active before range access?



Higher range use tends to impair final BW

WP6




No relationship between range use and body weight

Low-rangers  
High-rangers

# Results - Range use and meat production and quality? WP6

	JA757		S757N		Dual-purpose	
Range use	Low	High	Low	High	Low	High
Breast weight (g)	233	201	183	168	83	84

*In all strains but the dual-purpose, carcass/breast/thigh weights higher in Low Rangers than in High Rangers*




Foraging favors the intake of grass that contains coloring carotenoids

# Economic performance

## Comparison of production costs (EUR /100 kg live weight)

Calculated from  
2021 data



Thuenen Institute, Thobe et al., 2024

## PPILOW Comparison of the on-station fattening performances of dual-purpose breeds - males (WP5)

Meat-type Crossbreed	<b>Genotype A</b>	<b>Denmark</b>	<b>Germany</b>	<b>France Spring / summer</b>	<b>France Autumn / winter</b>
	Live weight wk 12, g	2019	2203	1977	1885
	FCR	3.1	3.4	3.3	3.4
Pure breed	<b>Genotype B</b>	<b>Denmark</b>	<b>Germany</b>	<b>France Spring / summer</b>	<b>France Autumn / winter</b>
	Live weight wk 12, g	1645	1763	1577	1466
	FCR	3.3	3.5	3.4	3.7
Layer-type Crossbreed	<b>Genotype C</b>	<b>Denmark</b>	<b>Germany</b>	<b>France Spring / summer</b>	<b>France Autumn / winter</b>
	Live weight wk 12, g	1732	1634	1393	1551
	FCR	3.1	3.7	3.2	3.6


Lombard et al.,  
2024

## Production costs of the use different dual-purpose breeds in organic experimental facilities in Germany in WP5 - males

Thuenen Institute

Calculated from  
2021 data

Production costs (€/100 kg meat)



Thobe et al., EAAP 2023



## PPILOW – Dual purpose females

## Results



## Evolution of the laying rate through time




Laying rate recorded every week from week 19 to week 72 in Germany and week 62 in Denmark

**Genotypes D** laying peaks higher than any of the dual purpose lines and sustained for a longer period of time

**Genotype A** always showed the lowest laying peak

Laying persistence: **genotype B < C**


genotype



Lombard et al., 2024

Pluschke et al., EAAP 2024

# PPILOW Genotypes & National Practitioner Group decision in WP5



Based on the laying data that represents the biggest part of revenues from dual-purpose genotypes, the NPG in each country selected the genotype to be tested on farm



13

	France		Germany	
	C	F (S757N)	C	D (JA757)
Mortality, %	4.6	1.4	11	1.2
FCR (13 wk)	3.7	2.6	3.7	2.7
Carcass weights at 13 wk, kg	1.38*	1,98*		2.4
Carcass weights at 15 wk, kg	1.72*	2.41*		
Carcass weights at 16 wk, kg			1.8	

\* Including neck



© Photos / Pluschke



© Hubbard Breeders

F



© Photos / Pluschke



© Hubbard Breeders

D

At week 13: Avg ± SE

	C	F (label)
Legs weight (g)	448 ± 9	668 ± 12
Wings weight (g)	180 ± 3	246 ± 4
Breast weight (g)	201 ± 5	354 ± 11

At week 15: Avg ± SE

	C	F (label)
Legs weight (g)	574 ± 12	838 ± 9
Wings weight (g)	219 ± 6	286 ± 3
Breast weight (g)	269 ± 4	462 ± 6

**C cuts from 40 to 30% lighter than F (label-type) cuts**

### Carcass conformation scores

	Genotype	Score 0	Score 1	Score 2
Wk 13	F (label)	<b>100%</b>	0	0
	C	0	0	<b>100%</b>
Wk 15	F (label)	<b>97%</b>	3%	0
	C	4%	<b>39%</b>	<b>58%</b>



### **Direct sale on-farm :**

- Consumers quite satisfied to buy **small carcasses**
- Interest for the approach
- But not sold at higher price


**Farmer enthusiastic with the ethical concept but technical adjustments and support needed**

Lombard et al., 2024

# Welfare and health indicators? Different breeds on the free range (WP6)

	JA757		S757N		Dual-purpose	
Pododermatitis (severe)	48%	40%	16%	12%	0%	0%
Hock burn % (severe)	8%	0%	0%	0%	0%	0%

Low Rangers
  High Rangers



Range use

**Bone health**  
 HR > LR in JA757  
**Higher locomotor activity**  
 of high-rangers



Bonnefous et al., 2023<sup>20</sup>

# PPILOW On-farm trials results – Welfare indicators in Germany (WP5)

**Control D**  
n=200

**JA757**

**Dual-purpose Genotype C**  
n=200




Pluschke et al., EAAP 2023



<sup>2</sup>  
<sup>2</sup>  
PPILOW On-farm trials results – Behaviour Observations in Germany in WP5

Proportions of behaviours during continuous observation in week before slaughter






Pluschke et al., EAAP 2023

# Results - Relationship between range use and **bird physiology (WP6)?**

Redox status

Antimicrobial defense



Low Rangers High Rangers


**In medium-growing birds, higher physical activity (in High ranging group) may limit muscle antimicrobial potential and increase oxidative stress**





## Results – Nutrient excretion (WP6)

Measure	S757N	JA757	Dual-purpose
Total feed intake (kg)	6207	6072	5332
Average weight per animal at commercial slaughter (kg)	2.6	3.2	1.9
<b>Average feed conversion ratio (g/g)</b>	<b>3.2</b>	<b>2.7</b>	<b>3.8</b>




Lower nutrient excretion with medium-growing JA757

*The quantity and distribution of outdoor excretion depends on the breed and range design*

## Working conditions, job perception and meat sensory quality compared to control label-type animals

- **JA757**: shorter rearing period, easier catching despite additional cleaning and straw bedding, lower stress and healthy working environment for the farmer
- Additional care needed for new **dual-purpose genotype C** :
  - smaller birds (feeders to adjust, feed spillage),
  - longer time indoors before range access (thermoregulatory needs),
  - much enrichment needed from start to avoid feather pecking,
  - quite stressful for the farmer/caretaker despite ethical interest
- The **meat of all strains was judged to be good** considering sensory quality



# Conclusions (1)

Interest of a **multi-trait approach** to evaluate the multiple consequences of range use and search for well-adapted breeds or birds

- **Best growth performance** and **lowest nutrient excretion** with **medium-growing JA757 strain** with the best (lowest) feed conversion ratio.

- Good meat quality parameters
- **Good perception by farmers/caretakers**

*Reasons why it has spread in organic farming throughout Northern Europe*

But...

**Animal welfare impaired** in our conditions (FPD), and redox status/health parameters affected in high-ranging animals...

... and **what about the breeders?**

Fast-growing  
male parental  
line



+



=



Feed restriction:  
*Sustainability of the  
whole production chain?*

## Conclusions (2)

- **Regulation (EU) 2018/848** of the European Parliament and of the Council **on organic production and labelling of organic products** and repealing Council Regulation (EC) No 834/2007.



**Organic chickens born as organic chick:** from organic breeders having access to the outdoor range  
*Questions about the genetics and breeding conditions – Exemption until 2036*

- **Label-type (slow-growing) chickens:**
  - Selection on a combination of range use (assessed by RFID) and growth?
  - Outdoor rearing of parental flocks and possibility of using a veranda during sanitary events?
- **Consider slow-growing breeds (among which local ones) in interaction with the free range design (agroforestry)?**  
*Presentation of Castellini, EAAP 2024*



## Conclusions (3)

### - Development of dual-purpose breeds in organic systems?



- For some organic farmers: the only ethical way to avoid the culling of layer male chicks
- Sustainability gains to find by nutrition (using by-products) for lower costs and management?
- Marketing actions towards consumers and public support?
- Consider laying performance together with meat production: cost of eggs and meat!

*Slow development in the absence of EU regulation on the culling of male chicks but only national ones*



*Towards sustainability assessments taking into account **all the dimensions** considered in “**multiperformance**”,  
e.g. including **Animal Welfare (and health)***

All stages of the production chain to be considered?

**Network of the European Partnership on Animal Health and Welfare (EUPAHW\*) to work on this topic**



28



# PPILOW PARTNERS

*Thanks to all the collaborators involved to to the members of the National Practitioner Groups!*



Fondazione Slow Food  
per la Biodiversità  
ONLUS



Harper Adams  
University



Instituut voor Landbouw-  
en Visserijonderzoek



Utrecht University



*Thanks for your attention!*

[www.ppilow.eu](http://www.ppilow.eu)