



HAL
open science

The *Dicorynia guianensis* population genomic structure mirrors the distribution of environmental variables in French Guiana

Julien Bonnier, Enrique Sáez-Laguna, Thomas Francisco, Olivier Brunaux, Sylvain Schmitt, Stephane Traissac, Niklas Tysklind, Myriam Heuertz

► **To cite this version:**

Julien Bonnier, Enrique Sáez-Laguna, Thomas Francisco, Olivier Brunaux, Sylvain Schmitt, et al.. The *Dicorynia guianensis* population genomic structure mirrors the distribution of environmental variables in French Guiana. 6. European Conservation Genetics Meeting 2024, Aug 2024, Lausanne, Switzerland. hal-04693713

HAL Id: hal-04693713

<https://hal.inrae.fr/hal-04693713v1>

Submitted on 10 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The *Dicorynia guianensis* population genomic structure mirrors the distribution of environmental variables in French Guiana

Julien Bonnier, Enrique Sáez Laguna, Thomas Francisco, Olivier Brunaux, Sylvain Schmitt, Stéphane Traissac, Niklas Tysklind & Myriam Heuertz

bonnier.ju@gmail.com


Background

Effective forest management can protect forests and enhance adaptation to climate change. In French Guiana, timber exploitation and climate change are threatening *Dicorynia guianensis*, a drought-sensitive tree that plays a crucial role in timber harvests¹. Understanding its genetic adaptation to environmental variations through landscape genomic studies² is essential to designing better management programs.

Objective: identify significant outlier loci associated with environmental variation in *Dicorynia guianensis* in French Guiana

Material: whole genome resequencing of 87 individuals distributed over 11 sites with varying environmental conditions

Methods: outlier analysis methods, based on genetic structure and genotype-environment association methods³



Figures. On top, geographic distribution of sampling sites of *Dicorynia guianensis* in French Guiana along the rainfall gradient with the number of individuals (N) in each site. On the left, picture of *Dicorynia guianensis* in sampling area.

Genetic structure and environmental variation

- **3 clusters** : west, inland, and east, improving the previously described population structure^{4,5}
- Regions characterized by **varying environmental conditions**, mainly influenced by rainfall during the rainy season

First signs of local adaptation

- **5 different methods** to detect evidence of selection.
- 459 SNPs were retained by at least four methods, highlighting **robust signals of selection**.

Precipitation and evapotranspiration during rainy season are the most significant variables modulating local adaptation and in shaping the genetic





Figure. Principal component analysis (PCA) based on 599,959 SNP markers and conducted on the sampling sites using the first two components (PC1 and PC2).




Figure : Biplot of redundancy analysis (RDA) illustrating the impact of environmental variables on genetic variation at 11 study sites for *Dicorynia guianensis* in French Guiana.

- Mean temperature of the coldest quarter (MTWQ)
- Potential evapotranspiration of the wettest quarter (PETWQ)
- Topographic wettest index (TW)
- Aridity index (AI)
- Precipitation of the coldest quarter (PCQ)

Forest management perspective

- Our results showcase the critical role of specific environmental factors, specially **heavy rainfall** during the rainy season, as strong drivers of local adaptation.
- Using genomic offset calculations, we **observed spatial patterns of maladaptation** risk in *D. guianensis* in French Guiana.

Our results advocate for the integration of patterns of genetic structure and local adaptation into management plans by (1) protecting **genetically diverse populations**; (2) developing **region-specific strategies** to cope with environmental change




Figure. Spatial projection of adaptive genetic turnover in the permanent forest domain (DFP) in French Guiana.

References

1. Flora, C. Origine et prédiction de la variabilité de la durabilité naturelle chez *Dicorynia guianensis* Amsh. (Université de Guyane, 2018). doi:10/document.
2. Guariguata, M. R., Cornelius, J. P., Locatelli, B., Forner, C. & Sánchez-Azofeifa, G. A. Mitigation needs adaptation: Tropical forestry and climate change. *Mitig Adapt Strateg Glob Change* 13, 793–808 (2008).
3. Rellstab, C., Gugerli, F., Eckert, A. J., Hancock, A. M. & Holderegger, R. A practical guide to environmental association analysis in landscape genomics. *Molecular Ecology* 24, 4348–4370 (2015).
4. Bonnier, J. et al. Population genetic structure and demographic history of the timber tree *Dicorynia guianensis* in French Guiana. *Tree Genetics & Genomes* 20, 2 (2024).
5. Caron, H. et al. Spatial and temporal distribution of chloroplast DNA polymorphism in a tropical tree species. *Molecular Ecology* 9, 1089–1098 (2000).