



**HAL**  
open science

## The very-high resolution configuration of the EC-Earth global model for HighResMIP

Eduardo Moreno-Chamarro, Thomas Arsouze, Mario Acosta, Pierre-Antoine Bretonnière, Miguel Castrillo, Eric Ferrer, Amanda Frigola, Daria Kuznetsova, Eneko Martin-Martinez, Pablo Ortega, et al.

### ► To cite this version:

Eduardo Moreno-Chamarro, Thomas Arsouze, Mario Acosta, Pierre-Antoine Bretonnière, Miguel Castrillo, et al.. The very-high resolution configuration of the EC-Earth global model for HighResMIP. *Geoscientific Model Development*, 2025, 18 (2), pp.461-482. 10.5194/gmd-2024-119 . hal-04701037

**HAL Id: hal-04701037**

**<https://hal.inrae.fr/hal-04701037v1>**

Submitted on 18 Sep 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



1                   **The very-high resolution configuration of the**  
2                   **EC-Earth global model for HighResMIP**

3  
4

5 Eduardo Moreno-Chamarro<sup>1,2\*</sup>, Thomas Arsouze<sup>1,3</sup>, Mario Acosta<sup>1</sup>, Pierre-Antoine Bretonnière<sup>1</sup>,  
6 Miguel Castrillo<sup>1</sup>, Eric Ferrer<sup>1</sup>, Amanda Frigola<sup>1</sup>, Daria Kuznetsova<sup>1</sup>, Eneko Martin-Martinez<sup>1</sup>,  
7 Pablo Ortega<sup>1</sup>, Sergi Palomas<sup>1</sup>

8  
9

10 1. Barcelona Supercomputing Center (BSC), Barcelona, Spain  
11 2. Now at: Max Planck Institute for Meteorology, Hamburg, Germany  
12 3. Now at: CIRAD, UMR AMAP, F-34398 Montpellier, France  
13 \* Corresponding author: eduardo.chamarro@mpimet.mpg.de

14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30



## 31 Abstract

32 We here present the very-high resolution version of the EC-Earth global climate model,  
33 EC-Earth3P-VHR, developed for HighResMIP. The model features an atmospheric resolution of  
34 ~16 km and an oceanic resolution of  $1/12^\circ$  (~8 km), which makes it one of the finest combined  
35 resolutions ever used to complete historical and scenario-like CMIP6 simulations. To evaluate  
36 the influence of numerical resolution on the simulated climate, EC-Earth3P-VHR is compared  
37 with two configurations of the same model at lower resolution: the ~100-km-grid  
38 EC-Earth3P-LR, and the ~25-km-grid EC-Earth3P-HR. The models' biases are evaluated against  
39 observations over the period 1980–2014. Compared to LR and HR, VHR shows a reduced  
40 equatorial Pacific cold tongue bias, an improved Gulf Stream representation with a reduced  
41 coastal warm bias and a reduced subpolar North Atlantic cold bias, and more realistic orographic  
42 precipitation over mountain ranges. By contrast, VHR shows a larger warm bias and overly low  
43 sea ice extent over the Southern Ocean. Such biases in surface temperature have an impact on the  
44 atmospheric circulation aloft, with improved stormtrack over the North Atlantic, yet worsened  
45 stormtrack over the Southern Ocean compared to the lower resolution model versions. Other  
46 biases persist with increased resolution from LR to VHR, such as the warm bias over the tropical  
47 upwelling region and the associated cloud cover underestimation, and the precipitation excess  
48 over the tropical South Atlantic and North Pacific. VHR shows improved air–sea coupling over  
49 the tropical region, although it tends to overestimate the oceanic influence on the atmospheric  
50 variability at mid-latitudes compared to observations and LR and HR. Together, these results  
51 highlight the potential for improved simulated climate in key regions, such as the Gulf Stream  
52 and the Equator, when the atmospheric and oceanic resolutions are finer than 25 km in both the  
53 ocean and atmosphere. Thanks to its unprecedented resolution, EC-Earth3P-VHR offers a new  
54 opportunity to study climate variability and change of such areas on regional/local spatial scales,  
55 in line with regional climate models.

56

## 57 1. Introduction

58 Interest in high-resolution modeling has soared in the past years, specially thanks to large  
59 European research projects and initiatives such as [PRIMAVERA](#), [nextGEMS](#), [EERIE](#), and  
60 [Destination Earth](#) (last access: 20 June 2024). Broadly, these projects seek to build the next  
61 generation of high-resolution global climate (or Earth system) models capable of representing



62 climate phenomena with unprecedented accuracy, to simulate and predict regional climate, guide  
63 policymaking, and provide relevant climate information to end users. Thanks to these efforts,  
64 high-resolution models at resolutions of 25–50 km or even finer have been proved to lead to  
65 reduced biases in the simulated climate (see Introduction in Moreno-Chamarro et al., 2022 for a  
66 review), and to a better representation of, for example, tropical cyclones (Roberts et al., 2020a;  
67 Vidale et al., 2021; Zhang et al., 2021), storm-tracks (e.g., Hodges et al., 2011), the intertropical  
68 convergence zone (ITCZ; e.g., Doi et al., 2012; Tian et al., 2020), or the Gulf Stream and  
69 associated air–sea interactions (e.g., Kirtman et al., 2012; Bellucci et al., 2021) compared to  
70 standard resolution models (hereafter, ~100-km grid). An extensive review of the benefit of  
71 high-resolution modeling can be found in Haarsma et al. (2016), Hewitt et al. (2017), Roberts  
72 M.J. et al. (2018), and Czaja et al. (2019). However, increased model resolution alone is not  
73 always the answer: for example, persistent, well-known biases in clouds and radiation can be  
74 insensitive to an increase in atmospheric resolution from a ~100-km grid to a 25–50-km grid  
75 (Moreno-Chamarro et al., 2022). Inadequate model physics or insufficient tuning can thus mask  
76 or negate the benefits of increased resolution.

77 High-resolution modeling faces additional challenges. One is the large computational cost  
78 needed to complete the simulations, which also limits the model throughput. Both issues have  
79 gradually improved thanks to steady increases in supercomputing power and parallel  
80 enhancements in model efficiency to leverage that power. The community trusts in High  
81 Performance Computing (HPC) to increase the performance of climate models, developing  
82 different approaches to speed models up. These approaches can go from improving the  
83 traditional parallelization algorithms (Tintó Prims et al., 2019a) or reducing the accuracy of the  
84 variables from double to single precision (Tintó Prims et al., 2019b) to increasing the  
85 Input/Output throughput of complex model configurations (Xepes-Arbós et al., 2022). Faster  
86 models are also needed to complete, in a reasonable time, the tuning and the spin-up phases,  
87 which for a high-resolution model, can be extremely costly. The demand for high efficiency in  
88 high-resolution modeling has therefore accelerated the development and implementation of new  
89 modeling strategies to ensure an optimal use of the computing resources.

90 High-resolution models also need to find a fair compromise between the resolutions of the  
91 different climate components, which, sometimes, can be very disparate—for example, an  
92 eddy-rich ocean model (~10 km grid) coupled to a 25 km, 50 km, or even coarser-grid





93 atmosphere model (e.g., Gutjahr et al., 2019). Tsartsali et al. (2022), for example, reported  
94 increased ocean–atmosphere coupling strength and better agreement with reanalysis and  
95 observations over the Gulf Stream, when both the ocean and atmosphere resolutions are  
96 increased to comparable ~25-km grid at least. Moreton et al. (2021) showed a degraded  
97 representation of the air–sea interaction at increased oceanic resolution but a constant  
98 atmospheric resolution. Similarly, Ma et al. (2016) found that the mesoscale ocean temperature  
99 affects the storm track over the Pacific only when the atmospheric model resolution is enough to  
100 resolve the small-scale diabatic heating. Finally, Rai et al. (2023) described a disproportionate  
101 eddy killing when a coarse 200-km wind forcing is used to force a finer (~10–25-km) ocean,  
102 compared to the case with similar grid sizes. These results of these studies thus advocate for a  
103 similar resolution in both the atmosphere and ocean.

104 High-resolution modeling usually relies on single-model component, either atmospheric-only  
105 (Baker et al., 2019) or ocean-only configurations (e.g., Biastoch et al., 2021), or on regional  
106 models (e.g., Woollings et al., 2010; Ma et al., 2017) as in CORDEX (Jacob et al., 2014) for  
107 hypothesis testing and downscaling climate projections. Such configurations, however, lack  
108 global energy constraints, remote influences, and, potentially, key feedbacks rectifying the mean  
109 state. These models are also limited by the boundary conditions, which often are derived from  
110 coarser (~100 km) global models and can present biases in their mean climate that might be  
111 absent or much reduced at a higher resolution; these biases might then be passed onto the single  
112 model configurations. For example, an overly smooth Gulf Stream temperature gradient, an  
113 incorrect separation, or the lack of mesoscale in ocean temperatures can impact the response of  
114 the atmospheric circulation aloft (e.g., Ma et al., 2017; Lee et al., 2018). Low-resolution and  
115 high-resolution global models can also respond differently to climate change: for example, the  
116 northward shift and strong surface warming of the Gulf Stream projected by the eddy-rich  
117 configuration of the HadGEM3-GC3.1 model for the 21st century is absent at the  
118 lower-resolution model versions (Moreno-Chamarro et al., 2021). Associated with this, the  
119 increase in winter precipitation is similarly much larger over Europe at the highest resolution  
120 than at any lower one, which reinforces the idea that the response of the atmosphere is strongly  
121 sensitive to the boundary conditions. These findings put a limit to our confidence in single-model  
122 configurations and regional models, since they lack a global dynamical response.



123 As a response to the listed challenges, we here present the eddy-rich version of the EC-Earth  
124 climate model for PRIMAVERA/HighResMIP. This is likely one of the finest combined  
125 horizontal resolution global models ever used to complete CMIP-like simulations, with a  
126 nominal resolution of about 10–15 km; it also has the additional advantage that the resolution is  
127 comparable in both the atmosphere and ocean/sea-ice, which allows the atmosphere to “see” the  
128 fine-scale forcing from the ocean with minimal information lost from interpolation. In this paper,  
129 we describe the model configuration and the developments in model efficiency (Section 2), as  
130 well as the main characteristics of its climate for the period 1980–2014 compared to observations  
131 (Section 3).

132

## 133 **2. Model Description and Experimental Setup**

### 134 **2.1 Model description**

135 All HighResMIP contributions with the EC-Earth global coupled climate model have been  
136 performed with its version 3.2.2, developed within the PRIMAVERA project (EC-Earth3P). The  
137 model consists of the atmosphere, ocean, and sea ice components. The atmosphere model is  
138 based on the ECMWF Integrated Forecasting System (IFS), in the 36r4 cycle. A detailed account  
139 of the changes introduced in this cycle can be found on the ECMWF website  
140 (<https://confluence.ecmwf.int/display/FCST/Implementation+of+IFS+Cycle+36r4>, last access:  
141 20 June 2024). The very-high resolution version of the model, EC-Earth3P-VHR, features a  
142 triangular truncation at wave number 1279 (hence known as T1279) in spectral space, with a  
143 linear N640 reduced Gaussian grid. This corresponds to a spacing of ~16 km. However, because  
144 of the complexity of numerical solutions and parametrizations, the effective resolution (this is the  
145 smallest scale IFS T1279 can fully resolve) is of ~120 km (Abdalla et al., 2013). Vertically, the  
146 model features 91 levels, resolving the middle atmosphere up to 0.01 hPa. The model time step  
147 during the simulation was 360 s. IFS integrates the revised land surface hydrology Tiled  
148 ECMWF Scheme for Surface Exchanges over Land (H-Tessel) model (Balsamo et al., 2009;  
149 Hazeleger et al., 2012).

150 The ocean model is the Nucleus for European Modelling of the Ocean in its version 3.6  
151 (NEMO3.6; Madec, 2008, Madec and the NEMO team, 2016). This is a hydrostatic,  
152 finite-difference, free-surface, primitive equation general circulation model. EC-Earth3P-VHR  
153 uses the ORCA12 tripolar grid, with the horizontal resolution increasing from the Equator to the



154 poles: ~9 km at the Equator, ~7 km at mid-latitudes, and ~2 km near the poles. This corresponds  
155 to an effective resolution of ~45 km (roughly five times the ORCA grid spacing). The model  
156 uses a  $z^*$  coordinate system for the vertical grid and has 75 vertical levels, with the resolution  
157 decreasing from 1 m at the surface to 200 m in the deep ocean. The bottom topography is derived  
158 from the combination of ETOPO1 (Amante and Eakins, 2009) and GEBCO\_08 (Becker et al.,  
159 2009). The sea ice model is the Louvain-la-Neuve sea Ice Model in its version 3 (LIM3)  
160 (Vancoppenolle et al., 2012). This is a dynamic-thermodynamic sea ice model, with five ice  
161 thickness categories. The time steps are 240 s for NEMO3.6, and 720 s for LIM3 in the  
162 EC-Earth3P-VHR.

163 The atmosphere–land and ocean–sea-ice components are coupled through the OASIS  
164 (Ocean, Atmosphere, Sea Ice, Soil) coupler, version 3 (Valcke and Morel, 2006; Craig et al.,  
165 2017). The remapping of runoff from the atmospheric grid points to runoff areas on the ocean  
166 grid was re-implemented to be independent of the grid resolution. This was done by introducing  
167 an auxiliary model component and relying on the interpolation routines provided by the OASIS  
168 coupler.

169 EC-Earth3P-VHR (hereafter, VHR) is compared with two lower-resolution global model  
170 versions, also run within the PRIMAVERA/HighResMIP project: EC-Earth3P (hereafter, LR;  
171 EC-Earth Consortium, 2019), and EC-Earth3P-HR (hereafter, HR; EC-Earth Consortium, 2018).  
172 In the atmosphere, they use the T255 (~107 km) and T511 (~54.2 km) spectral resolution of the  
173 IFS model respectively (equivalent to an effective resolution of ~600 km and ~280 km  
174 respectively; Abdalla et al., 2013), both with 91 vertical levels. In the ocean, LR and HR use the  
175 ORCA1 (~100 km) and ORCA025 (~25 km) tripolar grid respectively (equivalent to an effective  
176 resolution of ~500 km and ~125 km respectively), both with 75 vertical levels. They both use the  
177 LIM3 sea ice model and the OASIS coupler as well. LR and HR's time steps are respectively  
178 2700 s and 900 s in all the atmosphere, ocean, and sea ice. More details of these two other model  
179 versions can be found in Haarsma et al. (2020).

180 Following the CMIP6 HighResMIP protocol, no additional tuning is applied across  
181 resolutions but for a short list of parameters that explicitly change with resolution, particularly  
182 for oceanic diffusion and viscosity. The higher resolution in the atmosphere results in a better  
183 representation of features such as tropical storms, land/sea transitions, heavy rainfall, and fronts  
184 (see Fig. 1 as an example), while in the ocean the increase in resolution allows mesoscale




185 processes to be resolved at a much larger range of latitudes and the representation of finer  
186 resolution bathymetric features and coastlines.

### 187 **2.2 Configuration and workflow setup and performance optimization**


188 The development and maintenance of the EC-Earth model is supported by the EC-Earth  
189 Consortium, which shares model code, configurations, and minimal software infrastructure to  
190 operate it. While the LR and HR configurations of EC-Earth-3P were developed in a broad  
191 collaboration of all the consortium members participating in PRIMAVERA, VHR's development  
192 was primarily completed at the Barcelona Supercomputing Center, in collaboration with the  
193 Swedish Meteorological and Hydrological Institute (SMHI) within the [ESiWACE2](#) H2020  
194 project (last access: 20 June 2024). The development was conducted on two different  
195 supercomputing machines: [MareNostrum3](#), and [MareNostrum4](#) (last access: 20 June 2024).  
196 VHR's configuration, at the time of the project, represented one of the most cutting-edge  
197 versions of a climate model to run over long time scales. Obtaining a production version of the  
198 model, however, entailed the development of novel source code and execution scripts, the  
199 generation of all requisite files for initializing the simulations, and the adaptation of the model  
200 workflow software. This presented a significant challenge for both the operations department and  
201 the workflow developers, which were required to fine-tune the system to achieve stable runs and  
202 minimize the loss of computing hours. For example, generating the interpolation weight files to  
203 couple the new model grids for the OASIS coupler was particularly challenging. This process  
204 could not readily be parallelized at that time and required collaborating with the OASIS  
205 development group. For the workflow, a significant proportion of the effort was devoted to  
206 integrating the dedicated data transfer nodes available in the MareNostrum4 cluster into the  
207 workflow. Additionally, the automatic algorithm that enables the suppression of land grid  
208 subdomains in the NEMO ocean model was incorporated, resulting in a reduction of about 12%  
209 in the required HPC resources. Finally, the MareNostrum4 new network, despite its fast and  
210 responsive nature, proved to be quite unstable when subjected to high workloads involving  
211 multiple concurrent communications, as was the case of the VHR configuration. At the end of  
212 the ESiWACE2 project (December 2022), all the code was versioned and shared with the other  
213 partners within the EC-Earth Consortium.




a. ERA5





b. VHR



c. HR



d. LR



214  
215

216 **Figure 1.** Snapshot of an extratropical storm over the North Atlantic in the winter 1999–2000 in  
217 a) ERA5, and in the b) VHR, c) HR, and d) LR models on their original grids. Shown are daily  
218 precipitation rate ( $\text{mmd}^{-1}$ ; blue shading), cloud cover (% of area; gray shading), and sea-level  
219 pressure (hPa; contours).




220 Once deployed, the workflow needed to be made more efficient to be put into operation.  
221 Emerging advancements in global climate modeling demand heightened focus on HPC,  
222 particularly to accommodate the increasing need for enhanced model resolution (Acosta et al.,  
223 2024). An example of such demanding requirements is the VHR configuration, underscoring the  
224 need for efficient resource use. In order to address this issue, we conducted a two-fold HPC  
225 performance exercise, which involved both a pure computational performance analysis and a  
226 scalability study for each model component (IFS and NEMO), complemented with a load  
227 balance optimization for the coupling. This analysis concluded that the coupling and output  
228 process could be a bottleneck. An optimization was included to package different coupling fields  
229 to be sent in the same MPI (Message Passing Interface) communications, reducing the latency  
230 and taking advantage of the bandwidth. Additionally, the I/O (Input/Output) setup was optimized  
231 to ensure minimal time was needed to produce the outputs. While the primary objective of the  
232 scalability and load-balance study was to assess the model's efficiency and determine an optimal  
233 resource utilization, findings by Acosta et al. (2023) also indicate that enhancing the  
234 performance of one component, such as reducing the execution time of IFS, may not necessarily  
235 decrease the overall execution time of the coupled model. This discrepancy could stem from a  
236 synchronization point at the end of each coupled time step, where both components exchange  
237 fields. In cases where other non-optimized components lag behind, a load rebalance becomes  
238 necessary.

239 We ran a series of scalability tests to balance the resources (computing cores) of the VHR's  
240 IFS and NEMO models (Fig. 2). To find the most balanced configuration for a given amount of  
241 resources, we followed two different but complementary approaches. The first and most costly  
242 one tried to find the optimal distribution by assigning the same number of processors to IFS and  
243 NEMO first, and moving resources between them alternately; this allowed identifying the  
244 intervals for which the model performance increases by using variations of half-interval search  
245 algorithm. The second approach to balance the configuration started from one separate  
246 scalability test for each model component that was later used to determine the optimal  
247 configuration.

248

249

250



251

252 **Figure 2.** Results of the scalability test of the VHR configuration (T1279 IFS and ORCA12  
253 NEMO) at MareNostrum4 (blue line) in simulated years per day (SYPD) for a given amount of  
254 processors. The orange line shows the ideal case with no loss in computing performance.

255

256 The simulations described here were configured and run with the workflow manager  
257 Autosubmit (Manubens-Gil et al., 2016). This Python toolbox facilitates the production of  
258 numerical experiments, like the EC-Earth ones. It creates an oriented graph, taking into account  
259 every step of the workflow, including data pre- and post-processing, the transfer to storage  
260 spaces, or the conversion of the output data to CMOR standard, with details on computing  
261 resources needed for each step. Autosubmit also allows easily handling experiments with  
262 different members, start dates, and initial conditions.

263

### 264 2.3 Simulations

265 The VHR simulations follow the HighResMIP experimental protocol (Haarsma et al., 2016) and  
266 consist of: i) a 50-year spin-up run (spin-up-1950), with initial conditions of temperature and  
267 salinity from an ocean state representative of the 1950s (Good et al., 2013, EN4 data set) and  
268 forcing consisting of well-mixed greenhouse gases, including O<sub>3</sub> and aerosol loading for a 1950s  
269 (~10-year mean) climatology; ii) a 105-year control run (control-1950), starting from the end of





270 spin-up-1950 and keeping the same fixed forcing; iii) the historical run (hist-1950), starting from  
271 the same initial state as the control, but with time-varying external forcing for the period  
272 1950–2014; iv) and the future scenario run (highres-future), as a continuation of the historical  
273 simulation under the CMIP6 SSP5-8.5 scenario (Kriegler et al., 2017) for the period 2015–2050.  
274 In this work, VHR's hist-1950 simulation is compared with corresponding hist-1950 runs from  
275 LR and HR (Haarsma et al., 2020).

276 During the model setup, we erroneously applied the EN4 initial conditions at the beginning  
277 of all the spin-up runs. While EN4 uses practical salinity and potential temperature, the NEMO  
278 model, which uses the TEOS-10 equation of state, requires absolute salinity and conservative  
279 temperature. Nonetheless, the differences between the two temperature and salinity types is  
280 indeed small (Pawlowicz, 2013; McDougall et al., 2021), and we expect the error to minimize  
281 throughout the spin-up.

282

#### 283 **2.4 Observations and reanalysis**

284 As we mainly aim to evaluate the performance of EC-Earth3P-VHR configuration and describe  
285 the main model biases and characteristics, we focus on the best-observed part of the historical  
286 period of the historical simulations, between 1980 and 2014. The three model configurations are  
287 compared with the following observational and reanalysis data: near-surface (2 m) air  
288 temperature (SAT), zonal winds, sea-level pressure, and turbulent fluxes from the ERA5  
289 reanalysis (Hersbach et al., 2020); precipitation rate from the version-2 GPCP dataset (Adler et  
290 al., 2003); cloud cover from the version-3 ESA Cloud\_cci dataset (ESA CCI-CLOUD; Stengel et  
291 al., 2020); potential temperature and salinity of the ocean from the Hadley Center EN4 (version  
292 4.2.2; Good et al., 2013); sea ice concentration from OSI SAF (OSI-409/OSI-409-a;  
293 EUMETSAT Ocean and Sea Ice Satellite Application Facility, 2015); and sea ice volume from  
294 GIOMAS (Global Ice-Ocean Modeling and Assimilation System; Zhang and Rothrock, 2003).  
295 The period of comparison maximizes data availability and is therefore 1980–2014 for all the  
296 cases but for the GPCP dataset (1983–2014) and the ESA CCI-CLOUD dataset (1982–2014).  
297 Biases in sea-surface temperature (SST) are very similar to those in SAT and are therefore not  
298 shown.

299

300






### 301 3. Results

#### 302 3.1 Spin-up phase

303 Across all three model resolutions, the length of the spin-up (50 years) appears to be insufficient  
304 to equilibrate the full ocean (Fig. 3b); in fact; the ocean temperature is still drifting about  
305 0.001–0.002 °C/yr (computed over the last 50 years) towards warmer conditions at the end of the  
306 control simulation in the three configurations. In the upper ocean, however, VHR shows the  
307 smallest warming drift of the three configurations: about 0.00005 °C/yr compared to 0.0025  
308 °C/yr and 0.0062 °C/yr in HR and LR, respectively (computed over the last 50 years; Fig. 3a). It  
309 is therefore safe to say that an analysis focused on the upper ocean and on the air–sea interface  
310 will enjoy a relatively stable climate in the control simulations. In the historical simulations, the  
311 warming of the ocean accelerates due to the CO<sub>2</sub> forcing; after 64 years (year 114 in Fig. 3), the  
312 whole ocean warming reaches similar values to those at the end of the control simulations after  
313 100 years in the three model resolutions. Near the surface, the warming trend is much larger. Of  
314 the three configurations, VHR is the one with the smallest drift in the control run and the  
315 smallest ocean warming in the historical period. Although the three runs start from similar initial  
316 conditions derived from an EN4 climatology (Section 2.3), VHR is ~0.4 °C warmer near the  
317 surface than LR and HR, especially over the spin-up period. This is likely related to the  
318 development of a widespread warm bias over the Southern Ocean (Fig. 4), which we discuss in  
319 detail in Section 3.6.

320 In the following Sections, we describe the main characteristics of the VHR compared to LR  
321 and HR by focusing on particular regions and biases. This approach should help us highlight the  
322 benefits, or lack thereof, due to increased resolution. The main biases in the three model  
323 configurations are compared with the observational data set listed in Section 2.4.

324



325

326 **Figure 3.** Mean oceanic temperature in the LR (yellow), HR (red), and VHR (blue) models in the  
327 spin-up runs (0–50-year period), control runs (50–150-year period; solid lines), and historical  
328 runs (50–114-year period; dashed lines) in a) the upper 100 m, and b) the whole ocean. The  
329 vertical dashed line marks the end of the spin-up period

330

### 331 3.2 Tropics

332 A warm bias of 1–2 K is present over the subtropical upwelling regions along the South  
333 American and African coasts in the three configurations and shows small variations across them  
334 (Fig. 4). The increase in resolution in VHR has thus no clear benefit to reduce it. Past studies



335 have related this bias to an underestimation of the stratocumulus cloud deck (Richter, 2015). This  
336 also seems to be the case in the three models, which all show negative cloud biases by about 20  
337 % over all the subtropical upwelling areas, specially along the subtropical Pacific and Atlantic  
338 western coasts (Fig. 5). A better resolved orography near the region does not contribute to  
339 reducing the bias either, as suggested in previous studies (Milinski et al., 2016): for example,  
340 although VHR shows reduced temperature biases along the Andes compared to HR and LR, it  
341 has no effect on the biases over the eastern subtropical Pacific upwelling.

342 Overall, VHR shows reduced tropical precipitation biases compared to HR and LR (Fig. 6).  
343 This is the case, for example, for the double ITCZ bias: this bias is usually characterized by a  
344 precipitation excess over the central tropical North Pacific and the western tropical South Pacific  
345 and a precipitation deficit over the equatorial Pacific, as LR clearly shows. The dry area over the  
346 Equator is reduced with resolution, and the anomaly is even non-significant in VHR. This is a  
347 clear improvement from increased resolution, and it can be related to a reduced cold bias over  
348 the Equator (Fig. 4). In contrast, the precipitation excess over the tropical North Pacific and the  
349 Maritime Continent persists into VHR, with only minor reductions of 1–2  $\text{mmd}^{-1}$  compared to  
350 HR and LR (Fig. 6). The precipitation excess over the tropical North Pacific suggests a seasonal  
351 cycle reaching too far north, while the excess over the Maritime Continent, together with that  
352 over the western tropical Atlantic and Indian oceans, suggests an excess in convective  
353 precipitation over very warm waters.

354 Over the tropical Atlantic, the precipitation bias pattern points to an ITCZ anchored to the  
355 south-western part and not reaching the Sahel area. This bias is somewhat reduced in VHR  
356 compared to HR and LR, although not entirely removed. Over land, the dry bias over North  
357 Brazil and the wet bias along the Andes are not reduced with resolution, either. These positive  
358 and negative precipitation biases appear together with positive and negative biases in cloud  
359 cover, respectively, related to an overestimation or underestimation in convective clouds (Fig. 5).

360

361

362


363

364


365




a. VHR



b. HR



c. LR




366

367 **Figure 4.** Bias in SAT (in K) with respect to ERA5in the a) VHR, b) HR, and c) LR models for  
368 the period 1980–2014. Stippling masks anomalies that are not significant at the 5 % level.


369





a. VHR



b. HR



c. LR




370

371 **Figure 5.** Bias in cloud cover (in %) with respect to ESA CCI-CLOUD (contours in all the  
372 panels; in %) in the a) VHR, b) HR, and c) LR models for the period 1982–2014. Stippling  
373 masks anomalies that are not significant at the 5 % level.








a. VHR



b. HR



c. LR



374

375 **Figure 6.** Bias in precipitation rate (in  $\text{mmd}^{-1}$ ) with respect to GPCP (contours in all the panels;  
376 in  $\text{mmd}^{-1}$ ) in the a) VHR, b) HR, and c) LR models for the period 1983–2014. Stippling masks  
377 anomalies that are not significant at the 5 % level.



### 378 **3.3 Northern Hemisphere mid- and high-latitudes**


379 The largest improvement in the simulated climate from LR to VHR is over the North Atlantic.  
380 From south to north, the Gulf Stream representation is much improved in VHR compared to HR  
381 and LR, with sharper gradients in temperature and sea-surface height (not shown). The position  
382 of the Gulf Stream separation is also improved, which leads to a reduction of the warm bias  
383 along the US East Coast from LR to VHR (Fig. 4). A paper on a dedicated analysis of the biases  
384 over the North Atlantic along the Gulf Stream is currently in preparation.

385 Farther north, the widespread cold bias up to about 6 K in LR is strongly reduced in HR, and  
386 even further in VHR, which is the configuration closest to observations (Fig. 4). The cold bias in  
387 LR is related to an unrealistically large sea ice extent, which covers the entire Labrador Sea and  
388 the western part of the subpolar North Atlantic (Figs. 7 and 8). The reduction of the cold bias  
389 between LR and VHR bias has a deep impact on the climate of the North Atlantic. In the  
390 atmosphere aloft, it improves the representation of the boreal winter (DJF) stormtrack (Fig. 9)  
391 and jet (Fig. 10). The boreal winter stormtrack is overestimated over the subpolar North Atlantic,  
392 particularly over the eastern part, in LR, likely related to an excessively strong meridional  
393 temperature gradient; by contrast, VHR stormtrack is much closer to ERA5 over the North  
394 Atlantic. In the ocean, excessive sea ice leads to a negative salinity bias above 2 psu in the  
395 subpolar North Atlantic in LR, which is much reduced in VHR (Fig. 11). Two mechanisms can  
396 explain this fresh bias in LR: on the one hand, a reduced oceanic salinity transport from  
397 subtropical latitudes by a weakened subpolar gyre (not shown); on the other, errors in the  
398 seasonal cycle of the sea ice, during which ice melting would cause an anomalous freshwater  
399 input in regions where it is not observed. The negative bias in surface salinity propagates into  
400 deeper levels, especially between 300 m and 1000 m in the Arctic (Fig. 12). Similarly, the warm  
401 subsurface bias at around 40 °N might also be related to the sea ice excess in the subpolar North  
402 Atlantic in LR (Fig. 11). Expanded sea ice in LR causes weaker subpolar gyre strength and  
403 associated northward heat transport (not shown), leading to heat accumulation in the intergyre  
404 region. However, although this bias is reduced at higher resolutions in HR and VHR, it is still  
405 present, suggesting other deficiencies in the formation of intermediate waters in the North  
406 Atlantic. The overly large sea ice cover also hampers oceanic deep mixing in the Labrador Sea in  
407 LR, whose main region of deep water formations are in the Nordic Seas instead (Fig. 13).  
408 Oceanic deep mixing takes larger values above 1000 m in VHR and HR in the Labrador Sea. A



409 detailed analysis of the characteristics and driving mechanisms of the deep water formation in  
410 the Labrador Sea across the three resolutions and compared to observations is currently in  
411 preparation.

412




413

414 **Figure 7.** Sea ice concentration (in % of area) in the VHR model (gray shading) for the period  
415 1980–2014. Contours are the 15-% value in the LR (orange), HR (red), and VHR (blue) models,  
416 as well as in OSI SAF (black) for the period 1980–2014. Top/bottom panels are for the  
417 Arctic/Antarctic in March (left) and September (right).

418





419

420 **Figure 8.** Monthly climatology in the sea ice extent (in  $10^6 \text{ km}^2$ ; top) and volume (in  $10^3 \text{ km}^3$ ;  
421 bottom) in the Arctic (left) and Antarctica (right) in the LR (yellow), HR (red), and VHR (blue)  
422 models, as well as in OSI SAF, for sea ice extent, and GIOMAS, for the volume, for the period  
423 1980–2014.

424

425


426

427

428


429

430



431

432 **Figure 9.** Bias in winter stormtrack, computed as the standard deviation of the 2–6 d band-pass  
433 filtered daily sea-level pressure (in Pa) with respect to ERA5 (contours in all the panels; in Pa) in  
434 the a) VHR, b) HR, and c) LR models for the period 1980–2014. Each panel show anomalies in  
435 the boreal winter (DJF; top) and austral winter (JJA; bottom)




436

437 **Figure 10.** Bias in winter zonal wind at 250 hPa (in  $\text{ms}^{-1}$ ) with respect to ERA5 (contours in all  
438 the panels; in  $\text{ms}^{-1}$ ) in the a) VHR, b) HR, and c) LR models for the period 1980–2014. Stippling  
439 masks anomalies that are not significant at the 5 % level. Each panel show anomalies in the  
440 boreal winter (DJF; top) and austral winter (JJA; bottom)








a. VHR



b. HR




c. LR



441


442 **Figure 11.** Sea-surface salinity bias (in psu) with respect to EN4 (contours in all the panels; in  
443 psu) in the a) VHR, b) HR, and c) LR models for the period 1980–2014. Stippling masks  
444 anomalies that are not significant at the 5 % level.



445

446 **Figure 12.** Bias in ocean potential temperature (in K; top) and in salinity (in psu; bottom) with  
447 respect to EN4 (contours in all the panels; in K, top, and psu, bottom) in the a) VHR, b) HR, and  
448 c) LR models for the period 1980–2014. Stippling masks anomalies that are not significant at the  
449 5 % level. Each panel is separated into the upper and lower 500 m.

450




451

452 **Figure 13.** Mixed layer depth (in m) in the a) VHR, b) HR, and c) LR models for the period  
453 1980–2014. Northern Hemisphere and Southern Hemisphere values are for March and  
454 September, respectively.



455 Weak deep mixing results in a relatively weak Atlantic Meridional Overturning Circulation  
456 (AMOC; Fig. 14) in LR. The AMOC strength increases with resolution, related to the reduction  
457 of the cold bias and sea ice extent bias over the subpolar North Atlantic. The strength of the  
458 AMOC in VHR is thus the closest to the observed RAPID strength at 26 °N ( $17 \pm 3$  Sv;  
459 Frajka-Williams et al., 2019) among the three models:  $14 \pm 3$  Sv in VHR,  $12 \pm 4$  Sv in HR,  $11 \pm$   
460  $2$  Sv in LR (computed from monthly streamfunction at 26 °N for the period 2004–2014). The  
461 structure of the AMOC cell is similar in the three model configurations, with a main positive cell  
462 in the upper 3000 m up to 60 °N and with a maximum at around 30 °N, and a negative deeper  
463 one below with a strength of 2–4 Sv.



464

465 **Figure 14.** Atlantic overturning streamfunction (in Sv) in the a) VHR, b) HR, and c) LR models  
466 for the period 1980–2014.



467 In HR, and even more in VHR, the cold bias over the Labrador Sea is replaced by a warm  
468 bias (Fig. 4), up to 3–4 K in VHR. This bias also appears in other eddy-rich climate models,  
469 related to a stronger Atlantic ocean heat transport than at lower resolutions (Roberts et al.,  
470 2020b). Over the Nordic Seas, by contrast, a cold bias is present in the three models, although it  
471 is somewhat reduced at VHR by 1–2 K compared to LR and HR (Fig. 4). In the three cases, this  
472 bias is related to an excessively large sea ice cover in the region (Fig. 7). The warm bias over the  
473 Labrador Sea and cold bias over the Nordic Seas in VHR might suggest a misrepresentation of  
474 the distribution of oceanic heat transport between the two basins, favoring the westward transport  
475 over the northward across-Ridge heat transport. It might also or instead be related to a  
476 misrepresentation of the sea ice drift across the Denmark Strait (Gutjahr et al., 2022). Relatively  
477 weak transport across the Strait would lead to ice deficit in the Labrador Sea, and hence  
478 warming, and to ice accumulation in the Nordic Seas, hence cooling.

479 On a hemispheric scale, the three models simulate a slightly low Northern Hemisphere sea  
480 ice extent, mainly due to the underestimation of the sea ice cover in the Sea of Okhotsk, Baltic  
481 Sea, and Labrador Sea in HR and VHR (Fig. 8). By contrast, the three models show an overly  
482 large sea ice volume by about  $10^4$  km<sup>3</sup> compared to GIOMAS (Fig. 9), as they all simulate very  
483 thick sea ice in the central Arctic (not shown). Anomalously thick ice in the models leads to an  
484 excess of brine rejection (not shown), which can explain the positive salinity bias above 2 psu in  
485 the upper 100–200 m of the Arctic Ocean (Figs. 11 and 12). In VHR, the associated increase in  
486 upper-ocean density leads to deeper oceanic mixing than in LR or HR, with a mixed layer depth  
487 in the central Arctic that can reach up to 1000 m (Fig. 13).

488 Over the Pacific, biases tend to be weaker than over the Atlantic. A warm bias of about 1 K  
489 develops over the subpolar North Pacific from LR to VHR (Fig. 4), which could explain the  
490 negative bias in boreal winter (DJF) stormtrack aloft (Fig. 9) and the weaker jet stream over the  
491 central Pacific in VHR (Fig. 10).

492 Over land, the cold bias over the Sahara is reduced with increased resolution (Fig. 4).  
493 Similarly, the cold biases over large mountain ranges, such as the Rockies, the Andes, and the  
494 Himalaya, up to about several degrees in LR are much reduced in VHR, related to better resolved  
495 orography.

496

497





### 498 3.4. Southern Ocean

499 The Southern Ocean is the region where VHR performs the worst compared to HR and LR. The  
500 warm bias over the Southern Ocean increases with resolution, up to 4–5 K in VHR, compared to  
501 1–2 K and 2–3 K for HR and LR respectively (Fig. 4). It tends to be largest over the Atlantic and  
502 Indian sectors of the Southern Ocean and close to the Antarctic coast. Although the warm bias  
503 remains generally confined to the upper 100–200 m at around 60 °S, it might also be connected  
504 to the warm bias at depth between 2000 m and 4000 m (Fig. 12).

505 Two main mechanisms could explain the Southern Ocean warm bias: VHR has the largest  
506 cloud cover underestimation of the three models, especially over the Atlantic and Indian sectors,  
507 up to 15 % in VHR compared to 5–10 % in LR and HR (Fig. 5). Previous studies have related  
508 the Southern Ocean warm biases to misrepresentation and underestimation of the mixed-phase  
509 clouds, which lead to an excess of shortwave radiation reaching the surface, thereby warming it  
510 (e.g., Hwang, and Frierson, 2013; Hyder et al., 2018). Connected to the warm bias, VHR also  
511 shows the lowest sea ice extent of the three resolutions all year round (Figs. 7 and 8). Although  
512 the three models underestimate the Antarctic sea ice extent, in VHR this is nearly half as in  
513 observations for the same period (OSI SAF, 1980–2014). In terms of sea ice volume (Fig. 8),  
514 however, LR shows larger values by about  $2 \cdot 10^3 \text{ km}^3$  than GIOMAS between November and  
515 April, pointing to overly thick sea ice. As for the extent, VHR also shows the lowest sea ice  
516 volume, nearly half of the values in GIOMAS. The three models show the maximum volume one  
517 month later than in GIOMAS, in October rather than in September. This contrasts with the  
518 Arctic, where the three models capture the general shape of the seasonal cycle.

519 The surface warming over the Southern Ocean leads to a widespread underestimation of the  
520 stormtracks (Fig. 9) and jet stream (Fig. 10) in the austral winter (JJA) in HR and, especially, in  
521 VHR, compared to LR, which is much closer to ERA5. Although precipitation is also  
522 underestimated over the Southern Ocean, specially in VHR, this is not a particularly strong bias,  
523 at least compared to those over the tropical regions (Fig. 6).

524 Late austral summer (September) deep mixing tends to increase by about 200 m from LR to  
525 HR and VHR, especially in the Pacific sector. These two latter resolutions show similar deep  
526 mixing mean state, with variations only due to resolution and the better representation of the  
527 mesoscale in VHR (Fig. 13). The underestimation of the stormtrack over the Southern Ocean  
528 therefore does not seem to have an impact on the oceanic mixing below in VHR.



### 529 **3.5 Air–sea coupling**

530 We compare the change in the intensity of air–sea coupling from LR to VHR via the computation  
531 of cross-correlation coefficients of the deseasonalized monthly SST and net surface energy flux  
532 (Fig. 15). This analysis has extensively been used to study regions in which the ocean tends to  
533 drive atmospheric variability (correlation coefficient values approaching one) or vice versa  
534 (correlation coefficient values close to zero; e.g., Bishop et al., 2017; Small et al., 2019). The  
535 three model configurations are compared with the ERA5 reanalysis, as done in the previous  
536 Sections for the biases. To complement the analysis with a non-model based product, we also  
537 include satellite observations of radiative fluxes from J-OFURO3 (Tomita et al., 2019). The two  
538 products show an overall good agreement, with areas of large correlation coefficient values at the  
539 Equator, along the western boundary currents, and over the Southern Ocean (Fig. 15a,b). These  
540 areas, nonetheless, tend to be broader in J-OFURO3 than in ERA5.

541 Over the tropics, the three configurations tend to underestimate the coupling around the  
542 Equator, although they all reproduce well the band of correlation coefficients of high values  
543 along the equatorial Pacific and Atlantic. However, this band is narrower in LR and HR over the  
544 subtropics than it is in ERA5 and J-OFURO3. VHR is thus the closest configuration to the two  
545 reference observational products in the region. This result highlights the need for a model  
546 resolution finer than 25 km in both the ocean and atmosphere to represent realistic tropical  
547 climate interactions, in agreement with conclusions in Section 3.2.

548 At mid-latitudes, the coupling is greatly improved in HR and VHR compared to LR,  
549 particularly over the subpolar regions compared to ERA5 and J-OFURO3. LR shows a rather  
550 smooth pattern, with very low values in key regions over the Gulf Stream, Kuroshio Current, and  
551 Southern Ocean, which suggests a standard 1° resolution is insufficient to represent a realistic  
552 air–sea coupling. VHR and HR show, by contrast, sharper gradients in the correlation coefficient  
553 values close to 1 over those regions. This result is consistent with previous studies, which also  
554 found a degradation of the air–sea coupling in coarse grids, especially above 1° (e.g., Small et  
555 al., 2019). However, VHR shows unrealistic broader areas of higher correlation coefficient  
556 values than ERA5 and J-OFURO3 at mid-latitudes, degrading results from HR. One hypothesis  
557 for this discrepancy might result from the difference of IFS grid resolution between VHR  
558 (T1279) and ERA5 (T639), since the relationship between SST and turbulent fluxes shows  
559 certain scale dependency (e.g., Small et al., 2019; Sun and Wu, 2022). However, results do not




560 improve even when regriding VHR onto ERA5 grid before computing the correlation  
561 coefficients (not shown). A second hypothesis is the lack of the ocean current feedback in VHR,  
562 hence the lack of eddy-killing, which can control the simulated Gulf Stream's dynamics and  
563 energy pathways (Renault et al., 2023). However, the pattern of correlation coefficient values  
564 remains relatively unchanged when it is computed with a VHR configuration that includes a  
565 parameterization that considers the wind adjustment to the ocean current feedback (not shown)  
566 (Renault et al., 2019). The results suggest that the VHR's ocean exerts a stronger and more  
567 widespread influence on the atmosphere variability than in HR and LR.

568 Further north, air–sea coupling is overestimated in all the models over the Nordic Seas, likely  
569 related to the excess in sea ice in the region and its changes over the seasonal cycle. Together, the  
570 results suggest that a realistic air–sea coupling requires grids finer than  $1/4^\circ$  at least, with  
571 potential local improvements on a  $1/12^\circ$  grid, especially over the Tropics.

572

#### 573 **4. Discussion and Conclusions**

574 This paper presents the eddy-rich configuration of the EC-Earth3P-VHR global model for  
575 HighResMIP. We describe both the necessary technical developments to run the model  
576 efficiently, and the main features of the simulated climate compared to recent observations  
577 (1980–2014 period) and to two lower-resolution model configurations (the eddy-present,  
578  $\sim 25$ -km-grid EC-Earth3P-HR; and the non-eddy,  $\sim 100$ -km-grid EC-Earth3P-LR). The  
579 EC-Earth3P-VHR (or VHR) uses a comparable atmospheric and oceanic resolution of 10–15 km  
580 in a global fully coupled setup, which is, to our knowledge, one of the finest combined grids ever  
581 used to date to perform long climate integrations for CMIP. Our focus here is on the  
582 HighResMIP historical simulation (HighResMIP's hist-1950). This run is part of a larger set of  
583 runs, which includes a spin-up and control runs (HighResMIP's control-1950), a future extension  
584 under the ssp8.5 scenario (HighResMIP's highres-future), three hosing simulations forced by  
585 idealized Greenland melting, and AMIP sensitivity simulations, all performed within the  
586 European PRIMAVERA project and the Spanish STREAM project. Those additional simulations  
587 will be described in their corresponding publications, which are currently in preparation.



588

589 **Figure 15.** Cross-correlation coefficients between monthly SST and net surface energy flux for  
590 the period 1980–2014 in a) ERA5, b) J-OFURO3, and in the c) VHR, d) HR, and e) LR models.  
591 The seasonal cycle and linear trends are removed from the monthly SSTs and energy fluxes  
592 before the correlation coefficients are computed. This is done on the original grid in all the cases.



593 The comparison across the three resolutions (this is, VHR, HR, and LR), all with the same  
594 physics and no additional tuning, allows identifying regions where increased resolution improves  
595 the model performance with respect to observations. One of those regions is the Tropics, and  
596 specially the equatorial Pacific, where the cold tongue bias and the dry bias above are both  
597 reduced in VHR compared to HR and LR. Wengel et al. (2021) also reports a similar bias  
598 reduction in an eddy-resolving configuration of the CESM (0.25° resolution in the atmosphere,  
599 0.1° resolution in the ocean), which they link to better represented mesoscale features, such as  
600 tropical instability waves. Similarly, the HadGEM3-GC3.1 global model shows a reduced dry  
601 bias over the equatorial Pacific in its configuration with a 1/12° ocean and a 50-km atmosphere  
602 (Roberts et al., 2019). By contrast, the eddy-rich MPI-ESM1.2-ER global model (1/12° ocean as  
603 well) shows no evident changes in equatorial precipitation when coupled to a 100-km  
604 atmosphere (Gutjahr et al., 2019). Combined, these results suggest that resolutions finer than  
605 25–50 km might be needed in both the atmosphere and ocean to improve surface coupling and  
606 reduce biases. However, minimizing equatorial precipitation biases might actually be much more  
607 complex than simply increasing model resolution, as found for the ICON global  
608 atmosphere–ocean model with a uniform grid spacing of 5 km. Despite its high atmosphere and  
609 ocean resolutions, this model still exhibits a strong dry bias over the equatorial Pacific driven by  
610 a surface cold bias underneath (Hohenegger et al., 2022; Segura et al., 2022). This model,  
611 however, is not directly comparable to those other HighResMIP models, as it includes a  
612 minimum set of parametrization. Thus, while convection is directly resolved in ICON, it is  
613 parametrized in VHR and the listed models. The incorrect representation of the equatorial SST  
614 structure in ICON might instead be related to unresolved sub-grid processes (Segura et al.,  
615 2022).

616 The Gulf Stream is another region in which increased model resolution is beneficial, with a  
617 reduced temperature biases over the separation region and the central North Atlantic in VHR  
618 compared to HR and LR. Such improvements have been related to the resolving of the first  
619 baroclinic Rossby radius of deformation over most of the region and/or the exceeding of a  
620 critical Reynolds number (e.g., Chassignet and Marshall, 2008). Similar results have also been  
621 reported for the HadGEM3-GC3.1 (Roberts et al., 2019) and MPI-ESM1.2-ER (Gutjahr et al.,  
622 2019) global models, both with a 1/12° oceanic grid but coarser atmospheric grids (~50 km and  
623 ~100 km, respectively). This suggests that oceanic resolution is a critical factor for the Gulf



624 Stream representation. Nonetheless, other model features might also be relevant to simulate a  
625 realistic Gulf Stream, as no improvement is found in the CESM1.3 model between a 1°- and a  
626 0.1°- oceanic grid, for which the Gulf Stream separation occurs too far north (Chang et al.,  
627 2020). One of the many potential reasons behind the discrepancy might be the obvious difference  
628 in the number of atmospheric vertical levels: 91 in VHR, 85 in HadGEM3-GC3.1 (Roberts et al.,  
629 2019), 95 in MPI-ESM1.2-ER (Gutjahr et al., 2019), but only 30 in CESM1.3 (Meehl et al.,  
630 2019), which is expected to degrade the representation of key stratosphere–troposphere  
631 interactions affecting North Atlantic variability, and, by extension, the wind field, which is  
632 critical for the Gulf Stream separation. As nicely summarized in Chassignet and Marshall (2008),  
633 however: “The Gulf Stream separation, indeed, turns out to be quite sensitive to a variety of  
634 other factors such as subgrid scale parametrization, subpolar gyre strength and water mass  
635 properties, [deep western boundary current] strength, representation of topography, and the  
636 choice of model grid”. A realistic representation of the Gulf Stream is crucial for the North  
637 Atlantic and European climate. SST biases in the Gulf Stream can drive not only local changes  
638 over the North Atlantic, but a large-scale dynamic response over remote regions of the Northern  
639 Hemisphere through a quasi-zonal planetary barotropic Rossby wave response (Lee et al., 2018).  
640 Similarly, a more realistic, farther-south Gulf Stream has been shown to shift north in  
641 simulations with increased CO<sub>2</sub> in models at eddy-rich resolutions (Saba et al., 2016;  
642 Moreno-Chamarro et al., 2021). This shift would lead to amplified warming of the US East  
643 coastal region, which might be consistent with the anomalous warming observed in the Gulf  
644 Stream area in recent decades (Pershing et al., 2015; Todd and Ren, 2023). Reducing biases in  
645 the Gulf Stream area is therefore key to reproducing a realistic atmospheric circulation and to the  
646 sensitivity of the response to an external forcing.

647 Mainly related to increased atmospheric resolution, VHR also shows reduced precipitation  
648 biases over mountain ranges all over the world. This suggests VHR might provide more realistic  
649 regional information of precipitation variability and future changes than lower resolution models  
650 can. Giorgi et al. (2016), in fact, showed that increased model resolution leads to stronger  
651 summer precipitation changes over the Alpine region, using climate change projections with a  
652 regional atmospheric model of ~12-km grid. VHR uses a similar resolution but on a global scale,  
653 without the need to be constrained by lower resolution models.



654 On the negative side, we find that increased model resolution alone can be insufficient to  
655 reduce important and well-known biases in the climate or even cause model degradation in VHR.  
656 The warm bias over the coastal tropical upwelling areas, the Southern Ocean warm bias, and the  
657 rainfall excess bias over warm tropical waters all persist or even increase in VHR compared to  
658 HR and LR. These biases point to deficiencies in the model physics, specially in the atmosphere,  
659 and more particularly, in the cloud parameterizations. In VHR, both the warm bias over eastern  
660 tropical upwelling areas and the Southern Ocean are connected to negative biases in cloud cover.  
661 This reinforces the established idea that insufficient stratocumulus decks over the upwelling  
662 areas (e.g., Richter, 2015) and mixed-phase clouds over the Southern Ocean (e.g., Hyder et al.,  
663 2018) play key roles in setting up those biases. Cloud biases can be particularly insensitive to  
664 increases in model resolution, both in the ocean and atmosphere, from ~100-km grids to  
665 25–50-km grids (Moreno-Chamarro et al., 2022). Yet, for example, improved cloud  
666 microphysics closer to observations have been shown to help reduce shortwave radiation biases  
667 over the Southern Ocean in the Met Office's Unified Model (Varma et al., 2020). Reducing these  
668 biases as much as possible is critical, since they can have wider, global impacts on the climate,  
669 driving, for example, additional biases in tropical precipitation through the effect on the global  
670 energy budget (e.g., Hwang et al., 2013; Hawcroft et al., 2017).

671 It is interesting to note, nonetheless, that although LR, HR, and VHR all share the same cloud  
672 scheme, it is VHR that develops the strongest Southern Ocean bias. This might be related to the  
673 lack of additional model tuning from LR to HR and VHR. Rackow et al. (2024) showed that  
674 tuning the top-of-the-atmosphere radiation contributed to reducing the warming excess over the  
675 Southern Ocean in the IFS-FESOM global model at ~5-km resolution. The HighResMIP  
676 protocol suggests that no tuning is performed across resolutions to ensure any changes in the  
677 simulated climate can solely be attributed to changes in resolution (Haarsma et al., 2016). This  
678 approach can lead to undesired model degradation: for example, the untuned, low-resolution  
679 ECMWF model for HighResMIP shows an overly weak AMOC and a large cold bias over the  
680 North Atlantic compared to its well-tuned, high-resolution counterpart (Roberts C.D. et al.,  
681 2018). This can hinder model comparison and a clean understanding of the effect of model  
682 resolution, as biases can have large-scale climatic impacts (e.g., Hwang et al., 2013; Hawcroft et  
683 al., 2017; Lee et al., 2018) and affect the response sensitivity to forcing (e.g., McGee et al.,  
684 2018).





685 With respect to the spin-up, the HighResMIP protocol suggests a 50-year period (Haarsma et  
686 al., 2016). For all the configurations, this period is insufficient to equilibrate the full ocean,  
687 although the upper 1000 m equilibrates faster than the lower-part, and VHR does it faster and  
688 appears more stable after 100 years than HR and LR. The eddy-rich HadGEM3-GC3.1 also  
689 shows smaller drifts at the end of the 50-year period than its lowest resolution versions (Roberts  
690 et al., 2019). By contrast, for the CESM1.3 model, the low and high-resolution configurations  
691 only show a more stable climate after 150 years, related to a strong top-of-the-atmosphere energy  
692 imbalance (Chang et al., 2020). This led the authors to propose “150 to 200 years of model  
693 spin-up as a future strategy for initializing HR climate model simulations” (Chang et al., 2020).  
694 However, considering how computationally expensive these simulations are, new techniques  
695 might need to be introduced to tune and spin these models up faster and for longer. As much as  
696 tuning can still be “artisanal in character” at many research centers (Mauritsen et al., 2012), new  
697 and faster methods are being implemented to speed up the exploration of the space of parameters  
698 to find the best fit with observations. These methods include for example machine learning  
699 (Hourdin et al., 2021), simplified configurations (Wan et al., 2014), adjoints (Lyu et al., 2018),  
700 or model emulators (Williamson et al., 2013). Additional techniques have also been proposed to  
701 spin models up faster at much less computational costs; these include using for example  
702 Newton-Krylov methods (Bernsen et al., 2008; Merlis and Khatiwala, 2008), or replacing the  
703 atmosphere model by model data (Lofverstrom et al., 2020). Implementing similar techniques in  
704 future HR and VHR simulations would help accelerate both the spin-up and tuning phases.

705 To summarize, we here present the eddy-rich version of the EC-Earth global climate model,  
706 EC-Earth3P-VHR, with atmospheric and oceanic resolutions of 10–15 km. The analysis of its  
707 main climate features reveals improvements with respect to two lower resolution versions, such  
708 as a reduced dry equatorial bias over the Pacific, a more realistic Gulf Stream representation, and  
709 more accurate rainfall over mountain areas. Other biases persist or degrade, such as the warm  
710 biases over the subtropical upwelling regions and Southern Ocean, or the tropical precipitation  
711 excess. VHR's global resolution is at a similar level of many regional models, such as those  
712 participating in CORDEX, and it is much finer than most of the standard CMIP models. This  
713 opens a window of opportunity for model comparison and evaluation, as well as process  
714 understanding of much more realistic present-day and future climate and on a more regional  
715 scale.





716

### 717 Code and Data Availability

718 The data of the EC-Earth3P-LR and -HR models are available from ESGF  
719 (<https://esgf-index1.ceda.ac.uk/search/cmip6-ceda/>, last access: 20 June 2024) via the references  
720 provided in Section 2.3: EC-Earth3P (<https://doi.org/10.22033/ESGF/CMIP6.4683>, EC-Earth,  
721 2018; <https://doi.org/10.22033/ESGF/CMIP6.4682>, EC-Earth, 2019). Data of ERA-5 are freely  
722 available at <https://www.ecmwf.int/en/forecasts/dataset/ecmwf-reanalysis-v5> (Hersbach et al.,  
723 2020; <https://doi.org/10.24381/cds.6860a573>, Hersbach et al., 2019), while GPCP data are at  
724 <https://psl.noaa.gov/data/gridded/data.gpcp.html> (Adler et al., 2003), ESA cloud cover data are at  
725 <https://climate.esa.int/en/projects/cloud/data/> (Stengel et al., 2020), EN4 data version 4.2.2 are at  
726 <https://www.metoffice.gov.uk/hadobs/en4/> (Good et al., 2013), OSI SAF (OSI-409/OSI-409-a)  
727 sea ice concentration data are at <https://osi-saf.eumetsat.int/products/sea-ice-products>  
728 (EUMETSAT Ocean and Sea Ice Satellite Application Facility, 2015), GIOMAS sea ice volume  
729 data are at [https://psc.apl.washington.edu/zhang/Global\\_seaice/data.html](https://psc.apl.washington.edu/zhang/Global_seaice/data.html) (Zhang and Rothrock,  
730 2003), and J-OFURO3 flux data are at <https://www.j-ofuro.com/en/dataset/> (Tomita et al., 2019).  
731 The model data and plot scripts to reproduce the figures can be obtained from  
732 <https://zenodo.org/records/12078052> (Moreno-Chamarro, 2024). The model code developed at  
733 ECMWF, including IFS and the Finite Volume Module (FVM), is intellectual property of  
734 ECMWF and its member states. Permission to access the EC-Earth source code can be requested  
735 from the EC-Earth community via the EC-Earth website (<http://www.ec-earth.org/>, last access:  
736 July 2024) and may be granted, if a corresponding software license agreement is signed with  
737 ECMWF. The repository tag for the version of IFS and EC-Earth3P-VHR used in this work is  
738 3.2.2 (see Section 2.1) and is available through r8643. The EC-Earth workflow software used to  
739 run the simulations at the BSC is stored and version controlled in the BSC Earth Sciences GitLab  
740 repository (<https://earth.bsc.es/gitlab/es/auto-ecearth3>, last access: July 2024). Permission to  
741 access the repository can be requested from the Earth Sciences Department at the BSC and may  
742 be granted if the applicant has access to the EC-Earth code and the BSC HPC infrastructure. The  
743 workflow management system for running the simulations is distributed under Apache License  
744 2.0 as a public project (<https://earth.bsc.es/gitlab/es/autosubmit>, last access: July 2024) in the  
745 BSC GitLab repository.

746



#### 747 **Author Contributions**

748 TA, MA, MC, EF, and SP developed the model setup. EMC and TA ran the simulations. PAB  
749 and DK post-processed and cmorized the model data. EMC analyzed the data and wrote the  
750 manuscript with input from all the authors.

751

#### 752 **Competing interests**

753 The authors declare that they have no conflict of interest.

754

#### 755 **Acknowledgements**

756 This research has been supported by the Horizon2020 PRIMAVERA project (H2020 GA  
757 641727). EMC acknowledges funding from the Spanish Science and Innovation Ministry  
758 (Ministerio de Ciencia e Innovación) via the STREAM project (PID2020-114746GB-I00). MA  
759 has received funding from the National Research Agency through OEMES  
760 (PID2020-116324RA-I00). This work has received funding from the European High  
761 Performance Computing Joint Undertaking (JU) under the ESIWACE CoE, grant agreement No  
762 101093054.

763

#### 764 **References**

- 765 Abdalla, S., Isaksen, L., Janssen, P. A. E. M., and Nils, W.: Effective spectral resolution of  
766 ECMWF atmospheric forecast models, ECMWF Newsletter No. 137, 19–22,  
767 <https://doi.org/10.21957/rue4o7ac>, 2013.
- 768 Acosta, M.C., Palomas, S. and Tourigny, E.: Balancing EC-Earth3 Improving the  
769 Performance of EC-Earth CMIP6 Configurations by Minimizing the Coupling Cost. Earth  
770 and Space Science, 10(8), p.e2023EA002912, <https://doi.org/10.1029/2023EA002912>,  
771 2023.
- 772 Acosta, M. C., Palomas, S., Paronuzzi Ticco, S. V., Utrera, G., Biercamp, J., Bretonniere,  
773 P.-A., Budich, R., Castrillo, M., Caubel, A., Doblas-Reyes, F., Epicoco, I., Fladrich, U.,  
774 Joussaume, S., Kumar Gupta, A., Lawrence, B., Le Sager, P., Lister, G., Moine, M.-P.,  
775 Rioual, J.-C., Valcke, S., Zadeh, N., and Balaji, V.: The computational and energy cost of  
776 simulation and storage for climate science: lessons from CMIP6, Geosci. Model Dev., 17,  
777 3081–3098, <https://doi.org/10.5194/gmd-17-3081-2024>, 2024.



- 778 Adler, R. F., Huffman, G. J., Chang, A., Ferraro, R., Xie, P. P., Janowiak, J., Rudolf, B.,  
779 Schneider, U., Curtis, S., Bolvin, D., and Gruber, A.: The version-2 global precipitation  
780 climatology project (GPCP) monthly precipitation analysis (1979–present), *J.*  
781 *Hydrometeorol.*, 4, 1147–1167,  
782 [https://doi.org/10.1175/1525-7541\(2003\)004<1147:TVGPCP>2.0.CO;2](https://doi.org/10.1175/1525-7541(2003)004<1147:TVGPCP>2.0.CO;2), 2003 (data  
783 available at: <https://psl.noaa.gov/data/gridded/data.gpcp.html>, last access: 30 March 2023).
- 784 Amante, C. and Eakins, B.W.: ETOPO1 arc-minute global relief model: procedures, data  
785 sources and analysis, 2009.
- 786 Baker, A.J., Schiemann, R., Hodges, K.I., Demory, M. E., Mizielinski, M. S., Roberts, M. J.,  
787 Shaffrey, L. C., Strachan, J. and Vidale, P. L.: Enhanced climate change response of  
788 wintertime North Atlantic circulation, cyclonic activity, and precipitation in a  
789 25-km-resolution global atmospheric model. *Journal of Climate*, 32(22), 7763–7781,  
790 <https://doi.org/10.1175/JCLI-D-19-0054.1>, 2019.
- 791 Balsamo, G., Beljaars, A., Scipal, K., Viterbo, P., van den Hurk, B., Hirschi, M., and Betts, A.  
792 K.: A revised hydrology for the ECMWF model: Verification from field site to terrestrial  
793 waterstorage and impact in the Integrated Forecast System, *J. Hydrometeorol.*, 10, 623–643,  
794 2009.
- 795 Becker, J.J., Sandwell, D.T., Smith, W.H.F., Braud, J., Binder, B., Depner, J.L., Fabre, D.,  
796 Factor, J., Ingalls, S., Kim, S.H. and Ladner, R.: Global bathymetry and elevation data at 30  
797 arc seconds resolution: SRTM30\_PLUS. *Marine Geodesy*, 32(4), 355–371,  
798 <https://doi.org/10.1080/01490410903297766>, 2009.
- 799 Bellucci, A., Athanasiadis, P. J., Scoccimarro, E., Ruggieri, P., Gualdi, S., Fedele, G.,  
800 Haarsma, R. J., Garcia-Serrano, J., Castrillo, M., Putrahasan, D., and Sanchez-Gomez, E.:  
801 Air-Sea interaction over the Gulf Stream in an ensemble of HighResMIP present climate  
802 simulations, *Clim. Dynam.*, 56, 2093–2111, <https://doi.org/10.1007/s00382-020-05573-z>,  
803 2021.
- 804 Bernsen, E., Dijkstra, H.A., Thies, J. and Wubs, F.W.: The application of Jacobian-free  
805 Newton–Krylov methods to reduce the spin-up time of ocean general circulation models.  
806 *Journal of Computational Physics*, 229(21), 8167–8179,  
807 <https://doi.org/10.1016/j.jcp.2010.07.015>, 2010.



- 808 Biastoch, A., Schwarzkopf, F. U., Getzlaff, K., Rühs, S., Martin, T., Scheinert, M., Schulzki,  
809 T., Handmann, P., Hummels, R. and Böning, C. W.: Regional imprints of changes in the  
810 Atlantic Meridional Overturning Circulation in the eddy-rich ocean model VIKING20X.  
811 *Ocean Science*, 17(5), 1177–1211, <https://doi.org/10.5194/os-17-1177-2021>, 2021.
- 812 Bishop, S. P., Small, R. J., Bryan, F. O. and Tomas, R. A.: Scale dependence of midlatitude  
813 air–sea interaction. *Journal of Climate*, 30(20), 8207–8221,  
814 <https://doi.org/10.1175/JCLI-D-17-0159.1>, 2017.
- 815 Chassignet, E. and Marshall, D.: Gulf Stream separation in numerical ocean models.  
816 *Geophysical Monograph Series*, 177, <https://doi.org/10.1029/177GM05>, 2008.
- 817 Craig, A., Valcke, S., and Coquart, L.: Development and performance of a new version of the  
818 OASIS coupler, OASIS3-MCT\_3.0, *Geosci. Model Dev.*, 10, 3297–3308,  
819 <https://doi.org/10.5194/gmd-10-3297-2017>, 2017.
- 820 Czaja, A., Frankignoul, C., Minobe, S. and Vannière, B.: Simulating the midlatitude  
821 atmospheric circulation: what might we gain from high-resolution modeling of air-sea  
822 interactions?. *Current climate change reports*, 5, 390–406,  
823 <https://doi.org/10.1007/s40641-019-00148-5>, 2019
- 824 Doi, T., Vecchi, G. A., Rosati, A. J., and Delworth, T. L.: Biases in the Atlantic ITCZ in  
825 seasonal–interannual variations for a coarse-and a high-resolution coupled climate model, *J.*  
826 *Climate*, 25, 5494–5511, <https://doi.org/10.1175/JCLI-D-11-00360.1>, 2012.
- 827 EC-Earth Consortium (EC-Earth): EC-Earth-Consortium EC-Earth3P-HR model output  
828 prepared for CMIP6 HighResMIP hist-1950, Earth System Grid Federation [data set; last  
829 access: 18 May 2023], <https://doi.org/10.22033/ESGF/CMIP6.4683>, 2018.
- 830 EC-Earth Consortium (EC-Earth): EC-Earth-Consortium EC-Earth3P model output prepared  
831 for CMIP6 HighResMIP hist-1950, Earth System Grid Federation [data set; last access: 18  
832 May 2023], <https://doi.org/10.22033/ESGF/CMIP6.4682>, 2019.
- 833 EUMETSAT Ocean and Sea Ice Satellite Application Facility: Global sea ice concentration  
834 reprocessing dataset 1978–2015 (v1.2), Norwegian and Danish Meteorological Institutes,  
835 available at: <https://catalogue.ceda.ac.uk/uuid/8bbde1a8a0ce4a86904a3d7b2b917955> (last  
836 access: 8 February 2019), 2015.
- 837 Frajka-Williams, E., Ansorge, I.J., Baehr, J., Bryden, H.L., Chidichimo, M.P., Cunningham,  
838 S.A., Danabasoglu, G., Dong, S., Donohue, K.A., Elipot, S. Heimbach, P., Holliday, N.P.,



- 839 Hummels, R., Jackson, L.C., Karstensen, J., Lankhorst, M., Le Bras, I.A., Lozier, M. S.,  
840 McDonagh, E.L., Meinen, C.S., Mercier, H., Moat, B.I., Perez, R.C., Piecuch, C.G., Rhein,  
841 M., Srokosz, M.A., Trenberth, K.E., Bacon, S., Forget, G., Goni, G., Kieke, D., Koelling, J.,  
842 Lamont, T., McCarthy, G.D., Mertens, C., Send, U., Smeed, D.A., Speich, S., van den Berg,  
843 M., Volkov, D., Wilson, C.: Atlantic Meridional Overturning Circulation: Observed Transport  
844 and Variability, *Frontiers in Marine Science*, 6, <https://doi.org/10.3389/fmars.2019.00260>,  
845 2019.
- 846 Giorgi, F., Torma, C., Coppola, E., Ban, N., Schär, C. and Somot, S.: Enhanced summer  
847 convective rainfall at Alpine high elevations in response to climate warming. *Nature*  
848 *Geoscience*, 9(8), 584-589, <https://doi.org/10.1038/ngeo2761>, 2016.
- 849 Good, S. A., M. J. Martin, M. J., and Rayner, N. A.: EN4: quality controlled ocean  
850 temperature and salinity profiles and monthly objective analyses with uncertainty estimates,  
851 *Journal of Geophysical Research: Oceans*, 118, 6704-6716,  
852 <https://doi.org/10.1002/2013JC009067>, 2013 (data available at:  
853 <https://www.metoffice.gov.uk/hadobs/en4/>, last access: 12 November 2021).
- 854 Gutjahr, O., Jungclaus, J. H., Brüggemann, N., Haak, H. and Marotzke, J.: Air-sea interactions  
855 and water mass transformation during a katabatic storm in the Irminger Sea. *Journal of*  
856 *Geophysical Research: Oceans*, 127(5),  
857 e2021JC018075, <https://doi.org/10.1029/2021JC018075>, 2022.
- 858 Gutjahr, O., Putrasahan, D., Lohmann, K., Jungclaus, J.H., von Storch, J.S., Brüggemann, N.,  
859 Haak, H. and Stössel, A.: Max Planck Institute earth system model (MPI-ESM1.2) for the  
860 high-resolution model intercomparison project (HighResMIP). *Geoscientific Model*  
861 *Development*, 12(7), 3241-3281, <https://doi.org/10.5194/gmd-12-3241-2019>, 2019.
- 862 Haarsma, R., Acosta, M., Bakhshi, R., Bretonnière, P. A., Caron, L. P., Castrillo, M., Corti, S.,  
863 Davini, P., Exarchou, E., Fabiano, F. and Fladrich, U.: HighResMIP versions of EC-Earth:  
864 EC-Earth3P and EC-Earth3P-HR—description, model computational performance and basic  
865 validation. *Geoscientific Model Development*, 13(8), 3507-3527,  
866 <https://doi.org/10.5194/gmd-13-3507-2020>, 2020.
- 867 Haarsma, R. J., Roberts, M. J., Vidale, P. L., Senior, C. A., Bellucci, A., Bao, Q., Chang, P.,  
868 Corti, S., Fučkar, N. S., Guemas, V., von Hardenberg, J., Hazeleger, W., Kodama, C.,  
869 Koenigk, T., Leung, L. R., Lu, J., Luo, J.-J., Mao, J., Mizielinski, M. S., Mizuta, R., Nobre,



- 870 P., Satoh, M., Scoccimarro, E., Semmler, T., Small, J., and von Storch, J.-S.: High  
871 Resolution Model Intercomparison Project (HighResMIP v1.0) for CMIP6, *Geosci. Model*  
872 *Dev.*, 9, 4185–4208, <https://doi.org/10.5194/gmd-9-4185-2016>, 2016.
- 873 Haarsma, R., Acosta, M., Bakhshi, R., Bretonnière, P.-A., Caron, L.-P., Castrillo, M., Corti,  
874 S., Davini, P., Exarchou, E., Fabiano, F., Fladrich, U., Fuentes Franco, R., García-Serrano,  
875 J., von Hardenberg, J., Koenigk, T., Levine, X., Meccia, V. L., van Noije, T., van den Oord,  
876 G., Palmeiro, F. M., Rodrigo, M., Ruprich-Robert, Y., Le Sager, P., Tourigny, E., Wang, S.,  
877 van Weele, M., and Wyser, K.: HighResMIP versions of EC-Earth: EC-Earth3P and  
878 EC-Earth3P-HR – description, model computational performance and basic validation,  
879 *Geosci. Model Dev.*, 13, 3507–3527, <https://doi.org/10.5194/gmd-13-3507-2020>, 2020.
- 880 Hawcroft, M., Haywood, J.M., Collins, M., Jones, A., Jones, A.C. and Stephens, G., 2017.  
881 Southern Ocean albedo, inter-hemispheric energy transports and the double ITCZ: Global  
882 impacts of biases in a coupled model. *Climate Dynamics*, 48, 2279–2295,  
883 <https://doi.org/10.1007/s00382-016-3205-5>, 2017.
- 884 Hazeleger, W., Wang, X., Severijns, C., Ştefănescu, S., Bintanja, R., Sterl, A., Wyser, K.,  
885 Semmler, T., Yang, S., van den Hurk, B., van Noije, T., van der Linden, E., and van der  
886 Wiel, K.: EC-Earth V2.2: description and validation of a new seamless earth system  
887 prediction model, *Clim. Dynam.*, 39, 2611–2629, 2012.
- 888 Hewitt, H. T., Bell, M. J., Chassignet, E. P., Czaja, A., Ferreira, D., Griffies, S. M., Hyder, P.,  
889 McClean, J. L., New, A. L., and Roberts, M. J.: Will high-resolution global ocean models  
890 benefit coupled predictions on short-range to climate timescales?, *Ocean Model.*, 120,  
891 120–136, <https://doi.org/10.1016/j.ocemod.2017.11.002>, 2017.
- 892 Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J.,  
893 Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and  
894 Thépaut, J.-N.: ERA5 monthly averaged data on pressure levels from 1979 to present,  
895 Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [data set],  
896 <https://doi.org/10.24381/cds.6860a573>, 2019.
- 897 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas,  
898 J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X.,  
899 Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., Chiara, G. D., Dahlgren, P.,  
900 Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A.,



- 901 Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P.,  
902 Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and  
903 Thépaut, J.: The ERA5 global reanalysis, *Q. J. Roy. Meteor. Soc.*, 146, 1999–2049,  
904 <https://doi.org/10.1002/qj.3803>, 2020 (data available at:  
905 <https://www.ecmwf.int/en/forecasts/dataset/ecmwf-reanalysis-v5>, last access: 23 January  
906 2020).
- 907 Hodges, K. I., Lee, R. W., and Bengtsson, L.: A comparison of extratropical cyclones in  
908 recent reanalyses ERA-Interim, NASA MERRA, NCEP CFSR, and JRA-25, *J. Climate*, 24,  
909 4888–4906, <https://doi.org/10.1175/2011JCLI4097.1>, 2011.
- 910 Hohenegger, C., Korn, P., Linardakis, L., Redler, R., Schnur, R., Adamidis, P., Bao, J., Bastin,  
911 S., Behraves, M., Bergemann, M. and Biercamp, J.: ICON-Sapphire: simulating the  
912 components of the Earth system and their interactions at kilometer and subkilometer scales.  
913 *Geoscientific Model Development Discussions*, 1–42,  
914 <https://doi.org/10.5194/gmd-16-779-2023>, 2022.
- 915 Hourdin, F., Williamson, D., Rio, C., Couvreur, F., Roehrig, R., Villefranque, N., Musat, I.,  
916 Fairhead, L., Diallo, F. B. and Volodina, V.: Process-based climate model development  
917 harnessing machine learning: II. Model calibration from single column to global. *Journal of*  
918 *Advances in Modeling Earth Systems*, 13(6), e2020MS002225,  
919 <https://doi.org/10.1029/2020MS002225>, 2021.
- 920 Hwang, Y. T. and Frierson, D. M.: Link between the double-Intertropical Convergence Zone  
921 problem and cloud biases over the Southern Ocean, *P. Natl. Acad. Sci. USA*, 110,  
922 4935–4940, <https://doi.org/10.1073/pnas.1213302110>, 2013.
- 923 Hyder, P., Edwards, J. M., Allan, R. P., Hewitt, H. T., Bracegirdle, T. J., Gregory, J. M., Wood,  
924 R. A., Meijers, A. J., Mulcahy, J., Field, P., and Furtado, K.: Critical Southern Ocean  
925 climate model biases traced to atmospheric model cloud errors, *Nat. Commun.*, 9, 1–17,  
926 <https://doi.org/10.1038/s41467-018-05634-2>, 2018.
- 927 Jacob, D., Petersen, J., Eggert, B., Alias, A., Christensen, O. B., Bouwer, L. M., Braun, A.,  
928 Colette, A., Déqué, M., Georgievski, G. and Georgopoulou, E.: EURO-CORDEX: new  
929 high-resolution climate change projections for European impact research. *Regional*  
930 *Environmental Change*, 14, 563–578, <https://doi.org/10.1007/s10113-013-0499-2>, 2014.





- 931 Kirtman, B. P., Bitz, C., Bryan, F., Collins, W., Dennis, J., Hearn, N., Kinter, J. L., Loft, R.,  
932 Rousset, C., Siqueira, L., and Stan, C.: Impact of ocean model resolution on CCSM climate  
933 simulations, *Clim. Dynam.*, 39, 1303–1328, <https://doi.org/10.1007/s00382-012-1500-3>,  
934 2012.
- 935 Kriegler, E., Bauer, N., Popp, A., Humpenöder, F., Leimbach, M., Strefler, J., Baumstark, L.,  
936 Bodirsky, B.L., Hilaire, J., Klein, D., and Mouratiadou, I.: Fossil-fueled development  
937 (SSP5): An energy and resource intensive scenario for the 21st century. *Global*  
938 *Environmental Change*, 42, 297–315, <https://doi.org/10.1016/j.gloenvcha.2016.05.015>,  
939 2017.
- 940 Lee, R.W., Woollings, T. J., Hoskins, B. J., Williams, K. D., O'Reilly, C. H. and Masato, G.:  
941 Impact of Gulf Stream SST biases on the global atmospheric circulation. *Climate Dynamics*,  
942 51, 3369–3387, <https://doi.org/10.1007/s00382-018-4083-9>, 2018.
- 943 Lofverstrom, M., Fyke, J. G., Thayer-Calder, K., Muntjewerf, L., Vizcaino, M., Sacks, W. J.,  
944 Lipscomb, W. H., Otto-Bliesner, B. L. and Bradley, S. L.: An efficient ice sheet/Earth  
945 system model spin-up procedure for CESM2-CISM2: Description, evaluation, and broader  
946 applicability. *Journal of Advances in Modeling Earth Systems*, 12(8), e2019MS001984,  
947 <https://doi.org/10.1029/2019MS001984>, 2020.
- 948 Lyu, G., Köhl, A., Matei, I. and Stammer, D.: Adjoint-based climate model tuning:  
949 Application to the planet simulator. *Journal of Advances in Modeling Earth Systems*, 10(1),  
950 207–222, <https://doi.org/10.1002/2017MS001194>, 2018.
- 951 Ma, X., Chang, P., Saravanan, R., Montuoro, R., Nakamura, H., Wu, D., Lin, X. and Wu, L.:  
952 Importance of resolving Kuroshio front and eddy influence in simulating the North Pacific  
953 storm track. *Journal of Climate*, 30(5), 1861–1880,  
954 <https://doi.org/10.1175/JCLI-D-16-0154.1>, 2017.
- 955 Madec, G.: NEMO reference manual, ocean dynamic component: NEMO-OPA, Note du Pôle  
956 modélisation, Inst. Pierre Simon Laplace, France, 2008.
- 957 Madec, G. and the NEMO team: NEMO ocean engine version 3.6 stable, Note du Pôle de  
958 modélisation de l'Institut Pierre-Simon Laplace No. 27, ISSN: 1288–1619, 2016.
- 959 Manubens-Gil, D., Vegas-Regidor, J., Prodhomme, C., Mula-Valls, O. and Doblas-Reyes, F.  
960 J.: Seamless management of ensemble climate prediction experiments on HPC platforms. In



- 961 2016 International Conference on High Performance Computing & Simulation (HPCS),  
962 895–900, 2016, <https://doi.org/10.1109/HPCSim.2016.7568429>, 2016.
- 963 Mauritsen, T., Stevens, B., Roeckner, E., Crueger, T., Esch, M., Giorgetta, M., Haak, H.,  
964 Jungclaus, J., Klocke, D., Matei, D. and Mikolajewicz, U.: Tuning the climate of a global  
965 model. *Journal of advances in modeling Earth systems*, 4(3),  
966 <https://doi.org/10.1029/2012MS000154>, 2012.
- 967 McDougall, T. J., Barker, P. M., Holmes, R. M., Pawlowicz, R., Griffies, S. M. and Durack, P.  
968 J.: The interpretation of temperature and salinity variables in numerical ocean model output  
969 and the calculation of heat fluxes and heat content. *Geoscientific Model Development*,  
970 14(10), 6445–6466, <https://doi.org/10.5194/gmd-14-6445-2021>, 2021.
- 971 McGee, D., Moreno-Chamarro, E., Marshall, J. and Galbraith, E.D.: Western US lake  
972 expansions during Heinrich stadials linked to Pacific Hadley circulation. *Science advances*,  
973 4(11), p.eaav0118, <https://doi.org/10.1126/sciadv.aav0118>, 2018.
- 974 Meehl, G.A., Yang, D., Arblaster, J.M., Bates, S.C., Rosenbloom, N., Neale, R., Bacmeister,  
975 J., Lauritzen, P.H., Bryan, F., Small, J. and Truesdale, J.: Effects of model resolution,  
976 physics, and coupling on Southern Hemisphere storm tracks in CESM1. 3. Geophysical  
977 Research Letters, 46(21), 12408–12416, <https://doi.org/10.1029/2019GL084057>, 2019.
- 978 Merlis, T. M. and Khatiwala, S.: Fast dynamical spin-up of ocean general circulation models  
979 using Newton–Krylov methods. *Ocean Modelling*, 21(3-4), 97–105,  
980 <https://doi.org/10.1016/j.ocemod.2007.12.001>, 2008.
- 981 Milinski, S., Bader, J., Haak, H., Siongco, A. C., and Jungclaus, J. H.: High atmospheric  
982 horizontal resolution eliminates the wind-driven coastal warm bias in the southeastern  
983 tropical Atlantic: *Geophys. Res. Lett.*, 43, 10455–10462,  
984 <https://doi.org/10.1002/2016GL070530>, 2016.
- 985 Moreno-Chamarro, E.: Data for “The very-high resolution configuration of the EC-Earth  
986 global model for HighResMIP”, Zenodo [data set, last access: 10 July 2024],  
987 <https://doi.org/10.5281/zenodo.12078052>, 2024.
- 988 Moreno-Chamarro, E., Caron, L. P., Ortega, P., Tomas, S. L. and Roberts, M. J.: Can we trust  
989 CMIP5/6 future projections of European winter precipitation?: *Environmental Research*  
990 *Letters*, 16(5), 054063, <https://doi.org/10.1088/1748-9326/abf28a>, 2021.



- 991 Moreno-Chamarro, E., Caron, L. P., Loosveldt Tomas, S., Vegas-Regidor, J., Gutjahr, O.,  
992 Moine, M. P., Putrasahan, D., Roberts, C. D., Roberts, M. J., Senan, R. and Terray, L.:  
993 Impact of increased resolution on long-standing biases in HighResMIP-PRIMAVERA  
994 climate models: *Geoscientific Model Development*, 15(1), 269–289,  
995 <https://doi.org/10.5194/gmd-15-269-2022>, 2022.
- 996 Moreton, S., Ferreira, D., Roberts, M. and Hewitt, H.: Air-Sea Turbulent Heat Flux Feedback  
997 Over Mesoscale Eddies. *Geophysical Research Letters*, 48(20), e2021GL095407, 2021.
- 998 Pawlowicz, R.: Key physical variables in the ocean: temperature, salinity, and density. *Nature*  
999 *Education Knowledge*, 4(4), 13, <https://doi.org/10.1029/2021GL095407>, 2013.
- 1000 Pershing, A.J., Alexander, M.A., Hernandez, C.M., Kerr, L.A., Le Bris, A., Mills, K.E., Nye,  
1001 J.A., Record, N.R., Scannell, H.A., Scott, J.D. and Sherwood, G.D.: Slow adaptation in the  
1002 face of rapid warming leads to collapse of the Gulf of Maine cod fishery. *Science*,  
1003 350(6262), 809–812, <https://doi.org/10.1126/science.aac9819>, 2015.
- 1004 Rai, S., Hecht, M.W., Maltrud, M.E. and Aluie, H.: Scale-dependent Air-Sea Mechanical  
1005 Coupling: Resolution Mismatch and Spurious Eddy-Killing,  
1006 <https://doi.org/10.22541/essoar.167525271.13326232/v1>, 2023.
- 1007 Rackow, T., Pedruzo-Bagazgoitia, X., Becker, T., Milinski, S., Sandu, I., Aguridan, R.,  
1008 Bechtold, P., Beyer, S., Bidlot, J., Boussetta, S., Diamantakis, M., Dueben, P., Dutra, E.,  
1009 Forbes, R., Goessling, H. F., Hadade, I., Hegewald, J., Keeley, S., Kluft, L., Koldunov, N.,  
1010 Koldunov, A., Kölling, T., Kousal, J., Mogensen, K., Quintino, T., Polichtchouk, I.,  
1011 Sármany, D., Sidorenko, D., Streffing, J., Sützl, B., Takasuka, D., Tietsche, S., Valentini,  
1012 M., Vannière, B., Wedi, N., Zampieri, L., and Ziemann, F.: Multi-year simulations at  
1013 kilometre scale with the Integrated Forecasting System coupled to FESOM2.5/NEMOv3.4,  
1014 *EGUsphere* [preprint], <https://doi.org/10.5194/egusphere-2024-913>, 2024.
- 1015 Renault, L., Lemarié, F. and Arsouze, T.: On the implementation and consequences of the  
1016 oceanic currents feedback in ocean–atmosphere coupled models. *Ocean Modelling*, 141,  
1017 101423, <https://doi.org/10.1016/j.ocemod.2019.101423>, 2019.
- 1018 Renault, L., Marchesiello, P., & Contreras, M.: Coaction of top and bottom drags in Gulf  
1019 Stream dynamics. *Journal of Geophysical Research: Oceans*, 128, e2022JC018939.  
1020 <https://doi.org/10.1029/2022JC018939>, 2023.



- 1021 Richter, I.: Climate model biases in the eastern tropical oceans: Causes, impacts and ways  
1022 forward, *Wires Clim. Change*, 6, 345–358, <https://doi.org/10.1002/wcc.338>, 2015.
- 1023 Roberts, C. D., Senan, R., Molteni, F., Boussetta, S., Mayer, M., and Keeley, S. P. E.: Climate  
1024 model configurations of the ECMWF Integrated Forecasting System (ECMWF-IFS cycle  
1025 43r1) for HighResMIP, *Geosci. Model Dev.*, 11, 3681–3712,  
1026 <https://doi.org/10.5194/gmd-11-3681-2018>, 2018.
- 1027 Roberts, M. J., Baker, A., Blockley, E. W., Calvert, D., Coward, A., Hewitt, H. T., Jackson, L.  
1028 C., Kuhlbrodt, T., Mathiot, P., Roberts, C. D. and Schiemann, R.: Description of the  
1029 resolution hierarchy of the global coupled HadGEM3-GC3. 1 model as used in CMIP6  
1030 HighResMIP experiments. *Geoscientific Model Development*, 12(12), 4999–5028,  
1031 <https://doi.org/10.5194/gmd-12-4999-2019>, 2019.
- 1032 Roberts, M. J., Camp, J., Seddon, J., Vidale, P. L., Hodges, K., Vanniere, B., Mecking, J.,  
1033 Haarsma, R., Bellucci, A., Scoccimarro, E., and Caron, L. P.: Impact of model resolution on  
1034 tropical cyclone simulation using the HighResMIP–PRIMAVERA multimodel ensemble, *J.*  
1035 *Climate*, 33, 2557–2583, <https://doi.org/10.1175/JCLI-D-19-0639.1>, 2020a.
- 1036 Roberts, M. J., Jackson, L. C., Roberts, C. D., Meccia, V., Docquier, D., Koenigk, T., Ortega,  
1037 P., Moreno-Chamarro, E., Bellucci, A., Coward, A., and Drijfhout, S.: Sensitivity of the  
1038 Atlantic meridional overturning circulation to model resolution in CMIP6 HighResMIP  
1039 simulations and implications for future changes, *J. Adv. Model. Earth Sy.*, 12,  
1040 e2019MS002014, <https://doi.org/10.1029/2019MS002014>, 2020b.
- 1041 Roberts, M. J., Vidale, P. L., Senior, C., Hewitt, H. T., Bates, C., Berthou, S., Chang, P.,  
1042 Christensen, H. M., Danilov, S., Demory, M. E., and Griffies, S. M.: The benefits of global  
1043 high resolution for climate simulation: process understanding and the enabling of  
1044 stakeholder decisions at the regional scale, *B. Am. Meteorol. Soc.*, 99, 2341–2359,  
1045 <https://doi.org/10.1175/BAMS-D-15-00320.1>, 2018.
- 1046 Saba, V.S., Griffies, S.M., Anderson, W.G., Winton, M., Alexander, M.A., Delworth, T.L.,  
1047 Hare, J.A., Harrison, M.J., Rosati, A., Vecchi, G.A. and Zhang, R.: Enhanced warming of  
1048 the Northwest Atlantic Ocean under climate change. *Journal of Geophysical Research:*  
1049 *Oceans*, 121(1), 118–132, <https://doi.org/10.1002/2015JC011346>, 2016.
- 1050 Segura, H., Hohenegger, C., Wengel, C. and Stevens, B.: Learning by doing: Seasonal and  
1051 diurnal features of tropical precipitation in a global-coupled storm-resolving model.



- 1052 Geophysical Research Letters, 49(24), p.e2022GL101796,  
1053 <https://doi.org/10.1029/2022GL101796>, 2022.
- 1054 Small, R. J., Bryan, F. O., Bishop, S. P. and Tomas, R. A.: Air–sea turbulent heat fluxes in  
1055 climate models and observational analyses: What drives their variability?. *Journal of*  
1056 *Climate*, 32(8), 2397–2421, <https://doi.org/10.1175/JCLI-D-18-0576.1>, 2019.
- 1057 Stengel, M., Stapelberg, S., Sus, O., Finkensieper, S., Würzler, B., Philipp, D., Hollmann, R.,  
1058 Poulsen, C., Christensen, M., and McGarragh, G.: Cloud\_cci Advanced Very High  
1059 Resolution Radiometer post meridiem (AVHRR-PM) dataset version 3: 35 year climatology  
1060 of global cloud and radiation properties, *Earth Syst. Sci. Data*, 12, 41–60,  
1061 <https://doi.org/10.5194/essd-12-41-2020>, 2020 (data available at:  
1062 <https://climate.esa.int/en/projects/cloud/data/>, last access: 10 March 2021).
- 1063 Sun, X. and Wu, R.: Spatial scale dependence of the relationship between turbulent surface  
1064 heat flux and SST. *Climate Dynamics*, 58(3), 1127–1145,  
1065 <https://doi.org/10.1007/s00382-021-05957-9>, 2022.
- 1066 Tian, B. and Dong, X.: The double-ITCZ bias in CMIP3, CMIP5, and CMIP6 models based  
1067 on annual mean precipitation, *Geophys. Res. Lett.*, 47, e2020GL087232,  
1068 <https://doi.org/10.1029/2020GL087232>, 2020.
- 1069 Tintó Prims, O., M. C. Acosta, A. M. Moore, M. Castrillo, K. Serradell, A. Cortés and F. J.  
1070 Doblas-Reyes: How to use mixed precision in ocean models: exploring a potential reduction  
1071 of numerical precision in NEMO 4.0 and ROMS 3.6. *Geoscientific Model Development*,  
1072 12, 3135–3148, <https://doi.org/10.5194/gmd-12-3135-2019>, 2019a.
- 1073 Tintó Prims O., M. Castrillo, M. C. Acosta, O. Mula-Valls, A. Sanchez Lorente, K. Serradell,  
1074 A. Cortés and F. J. Doblas-Reyes: Finding, analysing and solving MPI communication  
1075 bottlenecks in Earth System models. *Journal of Computational Sciences*, 36, 100864,  
1076 <https://doi.org/10.1016/j.jocs.2018.04.015>, 2019b.
- 1077 Todd, R. E. and Ren, A. S.: Warming and lateral shift of the Gulf Stream from in situ  
1078 observations since 2001. *Nature Climate Change*, 13(12), 1348–1352,  
1079 <https://doi.org/10.1038/s41558-023-01835-w>, 2023.
- 1080 Tomita, H., Hihara, T., Kako, S.I., Kubota, M. and Kutsuwada, K.: An introduction to  
1081 J-OFURO3, a third-generation Japanese ocean flux data set using remote-sensing  
1082 observations. *Journal of Oceanography*, 75(2), 171–194,



- 1083 <https://doi.org/10.1007/s10872-018-0493-x>, 2019 (data available at:  
1084 <https://www.j-ofuro.com/en/dataset/>, last access: 14 March 2024).
- 1085 Tsartsali, E. E., Haarsma, R. J., Athanasiadis, P. J., Bellucci, A., de Vries, H., Drijfhout, S., de  
1086 Vries, I. E., Putrahasan, D., Roberts, M. J., Sanchez-Gomez, E. and Roberts, C. D.: Impact  
1087 of resolution on the atmosphere–ocean coupling along the Gulf Stream in global high  
1088 resolution models. *Climate Dynamics*, 58(11–12), 3317–3333,  
1089 <https://doi.org/10.1007/s00382-021-06098-9>, 2022.
- 1090 Valcke, S. and Morel, T.: OASIS and PALM, the CERFACS couplers, Tech. rep., CERFACS,  
1091 2006.
- 1092 Vancoppenolle, M., Bouillon, S., Fichefet, T., Goosse, H., Lecomte, O., Morales Maqueda, M.  
1093 A., and Madec, G.: The Louvain-la-Neuve sea ice model, Notes du pole de modélisation,  
1094 Institut Pierre-Simon Laplace (IPSL), Paris, France, No. 31, 2012.
- 1095 Varma, V., Morgenstern, O., Field, P., Furtado, K., Williams, J., and Hyder, P.: Improving the  
1096 Southern Ocean cloud albedo biases in a general circulation model, *Atmos. Chem. Phys.*,  
1097 20, 7741–7751, <https://doi.org/10.5194/acp-20-7741-2020>, 2020.
- 1098 Vidale, P. L., Hodges, K., Vannière, B., Davini, P., Roberts, M. J., Strommen, K., Weisheimer,  
1099 A., Plesca, E., and Corti, S.: Impact of stochastic physics and model resolution on the  
1100 simulation of Tropical Cyclones in climate GCMs, *J. Climate*, 34, 4315–4341,  
1101 <https://doi.org/10.1175/JCLI-D-20-0507.1>, 2021.
- 1102 Wan, H., Rasch, P. J., Zhang, K., Qian, Y., Yan, H. and Zhao, C.: Short ensembles: An  
1103 efficient method for discerning climate-relevant sensitivities in atmospheric general  
1104 circulation models. *Geoscientific Model Development*, 7(5), 1961–1977,  
1105 <https://doi.org/10.5194/gmd-7-1961-2014>, 2014.
- 1106 Wengel, C., Lee, S. S., Stuecker, M. F., Timmermann, A., Chu, J. E., and Schloesser, F.:  
1107 Future high-resolution El Niño/Southern Oscillation dynamics, *Nat. Clim. Change*, 1–8,  
1108 <https://doi.org/10.1038/s41558-021-01132-4>, 2021.
- 1109 Williamson, D., Goldstein, M., Allison, L., Blaker, A., Challenor, P., Jackson, L. and  
1110 Yamazaki, K.: History matching for exploring and reducing climate model parameter space  
1111 using observations and a large perturbed physics ensemble. *Climate Dynamics*, 41,  
1112 1703–1729, <https://doi.org/10.1007/s00382-013-1896-4>, 2013.



- 1113 Woollings, T., Hoskins, B., Blackburn, M., Hassell, D. and Hodges, K.: Storm track  
1114 sensitivity to sea surface temperature resolution in a regional atmosphere model. *Climate*  
1115 *Dynamics*, 35, 341–353, <https://doi.org/10.1007/s00382-009-0554-3>, 2010.
- 1116 Xepes-Arbós, X., G. van den Oord, M. C. Acosta and G. D. Carver: Evaluation and  
1117 optimisation of the I/O scalability for the next generation of Earth system models: IFS  
1118 CY43R3 and XIOS 2.0 integration as a case study. *Geoscientific Model Development*, 15,  
1119 379–394, <https://doi.org/10.5194/gmd-15-379-2022>, 2022.
- 1120 Zhang, J. and Rothrock, D. A.: Modeling global sea ice with a thickness and enthalpy  
1121 distribution model in generalized curvilinear coordinates, *Mon. Weather Rev.*, 131,  
1122 845–861, [https://doi.org/10.1175/1520-0493\(2003\)131<0845:Mgsiwa>2.0.Co;2](https://doi.org/10.1175/1520-0493(2003)131<0845:Mgsiwa>2.0.Co;2), 2003 (data  
1123 available at: [https://psc.apl.washington.edu/zhang/Global\\_seaice/model.html](https://psc.apl.washington.edu/zhang/Global_seaice/model.html), last access: 6  
1124 March 2019).
- 1125 Zhang, W., Villarini, G., Scoccimarro, E., Roberts, M., Vidale, P. L., Vanniere, B., Caron, L.  
1126 P., Putrasahan, D., Roberts, C., Senan, R., and Moine, M. P.: Tropical cyclone precipitation  
1127 in the HighResMIP atmosphere-only experiments of the PRIMAVERA Project, *Clim.*  
1128 *Dynam.*, 57, 253–273, <https://doi.org/10.1007/s00382-021-05707-x>, 2021.