



HAL
open science

THE EXTRUSION ROLE IN PHENOLIC COMPOUND AND PROTEIN BIOACCESSIBILITY FOR SORGHUM BASED PRODUCTS: A INFOGEST PROTOCOL-BASED STUDY

M E a S Oliveira, M G O Carvalho, H Mameri, C W P Carvalho, Mariana
Simões Larraz Ferreira

► **To cite this version:**

M E a S Oliveira, M G O Carvalho, H Mameri, C W P Carvalho, Mariana Simões Larraz Ferreira. THE EXTRUSION ROLE IN PHENOLIC COMPOUND AND PROTEIN BIOACCESSIBILITY FOR SORGHUM BASED PRODUCTS: A INFOGEST PROTOCOL-BASED STUDY. 3rd International Congress on Bioactive Compounds, Dec 2024, UNICAMP, Brazil. hal-04770716

HAL Id: hal-04770716

<https://hal.inrae.fr/hal-04770716v1>

Submitted on 7 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



4th International Workshop on Bioactive Compounds:
Functional Foods as the Cornerstone of Healthy Nutrition:
Linking Academia and Industry
December 10th to 11th, 2024, UNICAMP

THE EXTRUSION ROLE IN PHENOLIC COMPOUND AND PROTEIN BIOACCESSIBILITY FOR SORGHUM BASED PRODUCTS: A INFOGEST PROTOCOL-BASED STUDY

¹MOURA, B. S.; ²OLIVEIRA, M. E. A. S.; ³FERREIRA, R. M.; ⁴CARVALHO, M. G. O.;
⁵MAMERI, H.; ⁶CARVALHO, C. W. P.; ⁷FERREIRA, M. S. L.

¹Laboratory of Bioactives, Food and Nutrition Graduate Program (PPGAN), Federal University of the State of Rio de Janeiro (UNIRIO), Rio de Janeiro, RJ, Brazil, beatrizmoura@edu.unirio.br

²Laboratory of Bioactives, Food and Nutrition Graduate Program (PPGAN), Federal University of the State of Rio de Janeiro (UNIRIO), Rio de Janeiro, RJ, Brazil, mariaeugenia@edu.unirio.br

³Laboratory of Bioactives, Food and Nutrition Graduate Program (PPGAN), Federal University of the State of Rio de Janeiro (UNIRIO), Rio de Janeiro, RJ, Brazil, renatamarenda@gmail.com

⁴Laboratory of Bioactives, Food and Nutrition Graduate Program (PPGAN), Federal University of the State of Rio de Janeiro (UNIRIO), Rio de Janeiro, RJ, Brazil, matheus.carvalho@unirio.br

⁵UMR IATE, Univ. Montpellier, INRAE, Institut-Agro Montpellier, Montpellier, France, hamza.mameri@inrae.fr

⁶Embrapa Agroindústria de Alimentos, Rio de Janeiro, RJ, Brazil, carlos.piler@embrapa.br

⁷Laboratory of Bioactives, Food and Nutrition Graduate Program (PPGAN), Federal University of the State of Rio de Janeiro (UNIRIO), Rio de Janeiro, RJ, Brazil, mariana.ferreira@unirio.br

Keywords: *In vitro* digestion; HPLC; Protein digestibility; *Sorghum bicolor*.

Summary: Sorghum (*Sorghum bicolor* L.) has sustainable agronomic characteristics and represents an outstanding source of phenolic compounds (PC) and proteins compared to other cereals. The objective of this work was to evaluate how extrusion influences the PC bioaccessibility and *in vitro* protein digestibility (IVPD) in sorghum-based gluten-free products. Two different sorghum-based products were analyzed: cooked commercial

pasta and breakfast cereal and classical physical properties were determined. Free (FPC) and bound (BPC) phenolic compounds were sequentially extracted in ethanolic solution (80%), followed by alkaline and acid hydrolysis and analyzed by Folin-Ciocalteu. *In vitro* digestion was performed by INFOGEST and PC and protein bioaccessibility were determined by Kjeldahl and HPLC-DAD methods, respectively. The breakfast cereal showed the expected water absorption and solubility, and the pasta showed a solid loss and swelling compatible with gluten-free products. The cereal showed a higher PC content than pasta ($4654.8 \pm 0,0$ against 183.0 ± 33.0 mg GAE/100 g) and BPC was the main fraction (6.5 times) in both products. After digestion, PC presented a significant reduction in cereal (44%) and pasta (76%). Seven PC were quantified in the samples by HPLC based on standards. The content of the gallic acid (53%) and caffeic acid (207%) increased after digestion, as well as the presence of catechins, identified only in the digested breakfast cereal, due to hydrolysis of condensed tannins. IVPD was determined in the remained pellet, commercial pasta showed an IVPD of 48%, while breakfast cereal was 60%. Cooking extruded product presented a higher digestibility, probably due to the improved protein solubility, facilitating the proteolysis, conferred by the thermomechanical process. The PC bioaccessibility corresponded to 60% on average, while the IVPD was about 54% in the sorghum extruded products. These results corroborated the functional and nutritional characteristics of sorghum-based products, confirming the extrusion impact on PC and protein digestibility.

Financial support: Federal University of the State of Rio de Janeiro; Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro-FAPERJ (E26/201.402/2024); CNPq; CAPES (code 001).

