



HAL
open science

Ageing, retirement and changes in vegetable consumption in France: findings from the prospective GAZEL cohort

Marie Plessz, Alice Guéguen, Marcel Goldberg, Sébastien Czernichow, Marie Zins

► **To cite this version:**

Marie Plessz, Alice Guéguen, Marcel Goldberg, Sébastien Czernichow, Marie Zins. Ageing, retirement and changes in vegetable consumption in France: findings from the prospective GAZEL cohort. *British Journal of Nutrition*, 2015, 114 (6), pp.979 - 987. 10.1017/s0007114515002615 . hal-04789920

HAL Id: hal-04789920

<https://hal.inrae.fr/hal-04789920v1>

Submitted on 19 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Journal *British Journal of Nutrition*

Title **Aging, retirement and changes in vegetable consumption in France: findings from the prospective GAZEL cohort**

Version Accepted June 12th 2015

Authors Marie Plessz^{1,2}
Alice Guéguen²
Marcel Goldberg^{2,3}
Sébastien Czernichow^{2,3,4}
Marie Zins^{2,3}

Corresponding author Marie Plessz
tel + 33 1 49 59 69 12
fax + 33 1 49 59 69 90
email marie.plessz@ivry.inra.fr

Authors' addresses ¹ UR1303 ALISS Alimentation et sciences sociales, INRA, Ivry sur Seine, France
² Inserm, Population-based Epidemiologic Cohorts Unit, UMS 011, Villejuif, France
³ University of Versailles Saint Quentin, Versailles, France
⁴ Department of Nutrition, APHP, A. Paré hospital, Boulogne-Billancourt, France

Running title Aging, retirement and vegetable consumption

Keywords Aging, vegetable consumption, retirement, multiple imputations, meal place, GAZEL prospective cohort

Comment citer ce document :

Plessz, M. (Auteur de correspondance), Gueguen, A., Goldberg, M., Czernichow, S., Zins, M. (2015). Ageing, retirement and changes in vegetable consumption in France: findings from the prospective GAZEL cohort. *British Journal of Nutrition*, 114 (06, September 2015), 979-987. DOI : 10.1017/S0007114515002615

1 **Abstract**

2 The aim of this paper was to describe the change in vegetable consumption with aging and the
3 transition to retirement. Study subjects were the participants in the GAZEL prospective cohort
4 (Gaz and Électricité de France) aged 40-49 years at inclusion in 1989 who retired between
5 1991 and 2008 (12,942 men and 2,739 women). Four food-frequency questionnaires were
6 completed from 1990 to 2009. We used multiple imputation by chained equations in order to
7 avoid dropping incomplete cases. The odds-ratio for eating vegetables everyday was estimated
8 as a function of aging, retirement status and the place of lunch before retirement through
9 generalized estimating equations. Analyses were stratified by sex and models were adjusted for
10 confounders, including current spousal status. In 1990, 17.7% of men and 31% of women
11 reported eating vegetables daily. The odds of consuming vegetables everyday increased with
12 aging for both men and women. The usual place of lunch was home for less than half the
13 sample before retirement and for almost every respondent after retirement. For those who
14 changed their place of lunch, the association between being retired and the odds of eating
15 vegetables daily was positive and significant. We found that in this cohort, vegetable
16 consumption increased with aging. Retirement had an indirect effect on vegetable consumption
17 mediated by changes in the place of lunch.

19 *Keywords*

20 Aging, vegetable consumption, retirement, multiple imputations, meal place, GAZEL
21 prospective cohort.

22

23 Vegetables are a major focus of current nutritional guidelines in developed countries because
24 higher intakes could help protect from a wide range of non-communicable diseases⁽¹⁾. Recent
25 findings suggest that only 12% of the Americans meet the official dietary guideline for
26 vegetables⁽²⁾ and that the British consumed on average 135g/d of vegetables, 100g/d below UK
27 guidelines⁽³⁾. In France vegetable intakes appear to be higher, close to the levels reported in
28 Southern European countries⁽⁴⁾, with 55% of the population eating more than 2 servings (160g)
29 daily⁽⁵⁾. Barriers to vegetable consumption include cost, poor nutritional knowledge, and
30 limited cooking time and skills⁽⁶⁻⁸⁾. Although people spend more time in food preparation in
31 France than in many other European countries⁽⁹⁾, these factors are also associated with lower
32 vegetable consumption^(5,10,11).

33 Cross-sectional surveys suggest a strong association between age and vegetable consumption in
34 France and elsewhere^(5,12-15). However, it remains unclear if age reflects the process of aging,
35 differences across birth cohorts or the impact of specific life-course transitions⁽¹⁶⁻¹⁹⁾ such as
36 retirement. Studies of the transition to retirement have shown its positive impact on health
37 outcomes such as self-rated health⁽²⁰⁾ and sleep⁽²¹⁾, but results are unclear regarding physical
38 activity^(22,23) or alcohol consumption⁽²⁴⁾. In a cohort of 1,200 Finnish civil servants, Helldán et
39 al⁽²⁵⁾ found an increase in healthy food habits in retired women but not in retired men. Food
40 budget tends to decline after retirement⁽²⁶⁾ along with a decline in eating out⁽²⁷⁾ and an increase
41 in time devoted to cooking⁽²⁸⁾. However, adverse effects of retirement were also reported, e.g.,
42 people who retired from strenuous jobs were found to gain weight and waist circumference
43 while diminishing their fruit and vegetable intakes⁽²⁹⁾. All these findings suggest that retirement
44 might affect food consumption and especially vegetable consumption but research on this
45 aspect has been very limited by now⁽³⁰⁾.

46 The aim of the paper was to assess the relative effects of both the aging process and transition
47 to retirement on the odds of daily vegetable consumption in a cohort with 19 years of follow-
48 up.

49

50 **Methods**

51 *Study design*

52 The GAZEL cohort study is an occupational cohort composed of 20,625 employees of the
53 French national Gas and Electricity Company, born in 1939-48 (men) and 1939-53 (women).
54 They were included in 1989 and have been followed up since then. Follow-up includes a
55 mailed self-administered questionnaire every year; medical data from the medical department
56 of the company; and linkage to the company's human resources database and to the national
57 deaths registry, as described in detail elsewhere⁽³¹⁾. The company pays the pensions to the
58 retirees and could thus provide mailing addresses to the investigators. For this reason loss to
59 follow-up is very low (see Figure 1). A qualitative food frequency questionnaire (FFQ) was
60 included in the questionnaire in 1990, 1998, 2004 and 2009⁽³²⁾. The GAZEL study was
61 approved by the national commission overseeing ethics in data collection in France
62 (“Commission Nationale Informatique et Liberté”, CNIL, #105728).

63 *Population*

64 We selected individuals satisfying the following criteria: (a) aged 40 to 49y in 1989 (birth years
65 1939-48); (b) still alive at the end of 2009 (exclusion of 1352 deaths and 349 lost to follow-up);
66 (c) retired after the first FFQ in 1990 and before the last FFQ in 2009; (d) aged 50 to 61 when
67 they retired; (e) never had long-standing illness or disability, according to company records.
68 This last restriction aimed at excluding individuals whose retirement was due to health issues.

69 *Food frequency questionnaire and outcome variable*

70 The qualitative FFQ was designed so as to capture eating habits (rather than nutritional intake).
71 Subjects had to report how often they ate foods from various food groups over a typical week,
72 from “never” to “daily or almost daily”⁽³³⁾. In 1990, the FFQ included 10 food groups, e.g.,
73 potatoes (fried, mashed...); green vegetables (fresh, canned or frozen...). In 1998 the “green
74 vegetables” item was replaced by “cooked vegetables as starter, soup or main dish (leeks,
75 cabbage, green beans ...)” and “crudités or raw vegetables (green salad, carrots, tomatoes,
76 radish, beet...)”. The FFQ did not change after 1998.

77 The outcome variable was defined as eating vegetables daily (*vs* less than daily). In accordance
78 with French dietary guidelines, we did not include potatoes in the “vegetables” group. We used
79 the “green vegetables” item in the 1990 FFQ and the “cooked vegetables” item from the 1998
80 to 2009 FFQs, because they appeared to be closest in content and distribution.

81 *Exposure variables*

82 The variables of interest were age in years and retirement status. For age we used a piecewise
83 linear function with two knots at ages 50 and 61 in order to capture non-linearities. Retirement
84 status was coded using date of retirement as provided by company records.

85 *Mediator*

86 The impact of retirement on vegetable consumption could be mediated by a change in the usual
87 place of lunch after retirement. The most frequent place for lunch (home *vs* away from home)
88 was collected in the FFQ. We retained the most frequent place for lunch at the last FFQ before
89 the subject's retirement.

90 *Control variables*

91 All analyses were stratified by sex.

92 The prevalence of diets followed for health reasons increases with age⁽⁵⁾. We controlled for
93 whether the subject declared a diet prescribed by a doctor at each FFQ. Men living with a
94 partner eat more vegetables^(14,34,35). Changes in spousal status or in the spouse's activity status
95 may act as a confounder of the subject's retirement and were therefore estimated at each FFQ
96 year (no spouse, economically active spouse or inactive spouse).

97 Additionally, we controlled for the birth cohort, coded in two categories (born 1939-43 or
98 1944-48). Educational level was coded in three categories: low (primary school or less, leaving
99 school before age 12), medium (vocational or technical secondary education) or high
100 (secondary school degree or higher education)⁽³⁶⁾.

101 *Models*

102 In a first step, we modelled the outcome as a function of age, retirement status, and the
103 following potential confounders: date of FFQ (1990 versus 1998-2009), current status
104 regarding doctor-prescribed diet, birth cohort, education level and status of the spouse (model
105 1). In a second step, we studied whether a change in the usual place of lunch could mediate the
106 association between retirement status and vegetable consumption. For this purpose, an
107 interaction term between the place of lunch before retirement and retirement status was
108 introduced (model 2).

109 We ran logistic regressions using Generalized Estimating Equations (GEE) with an
110 unstructured correlation matrix, in order to account for the repeated nature of the data.

111 In order to test whether the change in FFQ items affected the results, we estimated model 2
112 excluding the data from year 1990.

113 *Treatment of missing data*

114 We imputed the non-responses using multiple imputation by chained equations^(37,38). This
115 technique allows imputing missing information for several variables at a time, through an
116 iterative process (the chained equations). Moreover, running multiple imputations produces
117 between-imputation variance, which accounts for the precision or imprecision of the imputation
118 process.

119 In the GAZEL study, non-response to the annual questionnaire is linked with gender, age,
120 occupational status and retirement status, as well as poor health and unhealthy lifestyle⁽³⁹⁾.

121 These variables had very high response rates because they were collected at inception in 1989
122 or from the company records (occupational status, retirement, age). They are also known
123 predictors of vegetable consumption^(5,12-14). In order to account for the (assumed) non-response
124 mechanism, the imputation model included the covariates from the model of interest as well as
125 the following auxiliary variables: self-reported health at inclusion, smoking status at inclusion,
126 occupational status at age 35 (see online supplementary materials for more details). Other
127 auxiliary variables (BMI and alcohol consumption at baseline) were tested and discarded
128 because they did not improve the quality of the imputations while creating colinearity issues.

129 We imputed the missing data separately for men and women. This allowed the imputation
130 model to be fully compatible with the specifications of the model of interest. It also warranted
131 that, if the non-response mechanism or the sample size made the imputation less precise for
132 women, this would not affect the results for men. We generated 50 completed datasets, ran the
133 GEE on these files and combined the results using Rubin's rules⁽⁴⁰⁾, which take into account the
134 variability in estimates and standard errors between the imputed datasets⁽³⁸⁾. We also ran model
135 2 on the complete cases only (model 3). We carried out all statistical analyses with Stata
136 version 12 (commands `mi impute chained` and `mi estimate: gee`). Results were
137 reported as odds-ratios with 95% confidence intervals.

138

139 **Results**

140 *Study population*

141 Figure 1 provides a flowchart for the selection of participants in the study. The population
142 included 12,942 men and 2,739 women. Descriptive statistics and number of missing values for
143 men and women are presented in Table 1 (baseline characteristics) and Table 2 (time-varying
144 variables). Because the company has specific rules on retirement, retirement occurred on
145 average at age 55. The average year of retirement was 1999 for men and 2000 for women. In
146 the cohort, a higher proportion of women ate vegetables every day over the whole follow-up
147 period, had a low educational level and were single ($p < 0.001$). Among the available cases
148 (respondents who completed the questionnaires in a given year), between 1990 and 2009, the
149 daily consumption of vegetable increased from 17.7% to 33.4% for men and from 31% to
150 55.3% for women. The proportion of individuals without a spouse increased for both sexes.
151 Crude associations between the repeated measurements of vegetable consumption appear in the
152 online supplementary material.

153 *Missing values and imputation*

154 Only 4,332 men and 855 women had no missing values (complete cases). The number of
155 missing outcomes went up after 1990 (Table 2). Figure 2 displays the proportion of subjects
156 who declared they ate vegetables everyday according to year of measurement for both sexes,
157 among the complete cases (listwise deletion), the available cases (pairwise deletion) and after
158 imputation (50 completed datasets). The proportion of positive outcomes is lower after
159 imputation than among the complete cases and the gap gets wider over time although
160 confidence intervals overlap.

161 *Aging and retirement*

162 Table 3 reports the odds-ratios of daily consumption of vegetables. Model 1 showed the
163 evolution of vegetable consumption as subjects grew older and retired, based on the imputed
164 data. The odds of eating vegetables daily increased steadily by 6% per year of age for men
165 (95% CI 5%,7%), and 7% for women (95% CI 4%,10%) until age 61. After age 61 the increase
166 was smaller but remained statistically significant.

167 Model 1 also showed that being retired had a positive and significant effect on the odds of
168 eating vegetables every day for men, but the effect was not significant for women. For men,
169 retiring was equivalent to being three years older in terms of vegetable consumption.

170 *Place of lunch before retirement*

171 Based on the available cases of our sample (pairwise deletion), at the last FFQ before
172 retirement the most frequent place for lunch was away from home for 53.7% of men and 69.9%
173 of women (Table 1). After retirement only 1.5% of the respondents had most lunches away
174 from home, implying that most of those who used to lunch out started to lunch at home.
175 Model 2 in Table 3 shows that before retirement, men who had most of their lunches at home
176 ate vegetables significantly more often than those who had lunch away from home (OR= 1.28,
177 95% CI 1.18,1.39). The difference was smaller and not significant for women (OR= 1.16).
178 After retirement, there was no significant difference between the subjects who used to have
179 lunch away from home when working and those who did not.
180 Indeed, for men who had lunch mostly away from home before retirement, the effect of
181 retirement was positive and statistically significant (OR=1.31, 95% CI 1.17, 1.46). For these
182 men, getting retired increased the chances of daily vegetable consumption in the same
183 proportions as being five years older. By contrast, those who had lunch mostly at home when
184 working increased only slightly and non-significantly their vegetable intake upon retirement
185 (before retirement OR= 1.28 versus after retirement: 1.38). The same is true for women: those
186 who used to lunch out increased their vegetable consumption after retirement (OR= 1.21) while
187 for the others the odds-ratios before and after retirement were very close (1.16 versus 1.22).
188 Figure 3 displays the results of model 2 as predicted probabilities of eating vegetables daily if
189 the subjects had retired at age 56. In terms of probabilities of eating vegetables daily, the gap
190 between men and women grew larger over time. The transition to retirement implied no change
191 for those who had lunch at home at the end of their active life. There was a positive change for
192 those who had lunch away from home prior to retirement, so that after retirement the two
193 groups did not differ in their vegetable consumption.

194 *Robustness checks*

195 In order to assess the impact of the imputation process on the results we ran Model 2 on the
196 complete cases only (Model 3 in Table 3). Changes in estimate values were modest and non-
197 significant for men. Some coefficients varied more markedly for women, possibly due to the
198 smaller number of complete cases.

199 In 1990 the question on vegetables was about “green vegetables” while later on, the item was
200 “cooked vegetables”. We tested whether this change affected the estimates by running model 2
201 on the imputed data after excluding the 1990 FFQ, and found very similar results (online
202 supplementary material).

203 Discussion

204 We used four FFQ over 19 years of follow-up to assess the evolution of vegetable consumption
205 between ages 42 and 70, as subjects aged and retired. Our results show that for both men and
206 women, the process of aging was associated with gradually increasing odds of eating
207 vegetables every day. The increase in vegetable consumption observed upon the transition to
208 retirement appeared to be mediated by the place of lunch before retirement. These results hold
209 after imputing non-responses and taking into account the main potential confounders.

210 Age is associated with increased vegetable consumption in cross-sectional studies^(5,12-14). Our
211 study suggests that as a given birth cohort grows older, its members tend to eat vegetables more
212 often. This aging effect remains positive and significant after controlling for retirement status,
213 but also for current spousal status, current dieting prescription status, and birth cohort. The
214 aging effect is relatively stable over time, at least from 42 to 61 years of age, and may decline
215 slightly afterwards.

216 Retirement is a major life-course transition between midlife and old age. While the association
217 between transition to retirement and changes in health-related behaviour, such as alcohol
218 drinking and physical activity, have been studied before, there are only a few studies on
219 changes in food intake^(25,30,41). In this cohort study, retiring was associated with increased
220 vegetable consumption in those respondents whose meal environment changed upon
221 retirement. This is consistent with the life-course perspective on food consumption⁽¹⁶⁻¹⁹⁾.

222 Food eaten away from home seems less healthy than food eaten at home^(42,43). Evidence is less
223 clear on the nutritional quality of meals taken at work, e.g. in staff or university canteens⁽⁴⁴⁻⁴⁶⁾.

224 In France a nationally representative survey including a 7-day food diary found that 26% of the
225 lunches eaten by adults were taken away from home⁽⁴⁷⁾. For the US a similar figure (25%) was
226 found in the NHANES survey 2003-2004⁽⁴²⁾. In our study, those who had lunch away from
227 home while working ate vegetables less often, but vegetable consumption after retirement did
228 not differ according to the place of lunch *before* retirement.

229 This result may be related to the impact of the spouse on male respondents' food habits. In the
230 GAZEL study, men's BMI depended more on the spouse's social status than on the
231 respondent's occupational category⁽⁴⁸⁾. In the present study, the presence of a spouse was
232 associated with higher odds of eating vegetables every day for males. Women's vegetable
233 consumption was higher than men's and less affected by the presence of a spouse. This may be
234 linked to the fact that women have better nutritional knowledge⁽⁸⁾ and do most of the cooking in
235 European countries⁽⁴⁹⁾.

236 Our hypothesis was that getting retired may have a causal effect on vegetable consumption,
237 through changes in the place of lunch. In order to move from the measurement of an
238 association to the identification of a causal pathway, we have taken several precautions. A
239 reverse causality path from vegetable consumption to retirement was unlikely. We controlled
240 for several sources of confusion besides aging. A poor health condition during active life could
241 lead both to retiring earlier than expected and to adopting healthier eating habits, including
242 higher vegetable consumption. We therefore excluded from the study population all individuals
243 who had a long-standing illness or disability, according to the company records, and we
244 controlled for current dieting prescription status. It is also necessary to separate the subject's
245 retirement from his or her spouse's retirement: we controlled for the presence and the activity
246 status of the spouse at each FFQ date. Additionally we controlled for birth cohort, education
247 level and changes in the questionnaire.

248 Another potential source of confusion is the period of observation. Over 60% of the subjects
249 retired between 1998 and 2004. In 2001 France launched nutrition information campaigns
250 including the message to eat "five fruits and vegetables a day". According to cross-sectional
251 surveys, the share of the adult population aware of this nutritional guideline (2.5% in 2002)
252 increased 11-fold between 2002 and 2008⁽⁵⁾. However, based on the 24-hour recalls included in
253 the same surveys, the level of vegetable consumption had not changed much: the share of the
254 French population who ate five fruit and vegetable servings a day was estimated at 10.2% in
255 2002 and 13% in 2008^(5,50). Confusion between the measured effect of aging and the impact of
256 nutrition information campaigns during the observation period might play a role, but it is very
257 unlikely that it would explain all the aging and retirement effects.

258 An important limitation regarding the interpretation of the results is the population included in
259 the survey. Respondents were all employees of a large, state-owned French company. While
260 their social background was diverse (over 80% entered the firm as blue-collar or clerical
261 workers⁽³¹⁾), the subjects' risk of being laid-off was virtually null and their income did not drop
262 substantially after retirement. It is therefore likely that the impact of retirement would be lower
263 in a population exposed to unemployment or having less protective retirement status. Similarly,
264 it would be unwise to extend our conclusion to other age-groups or birth cohorts than those
265 included in this survey. Indeed, it appears that in the GAZEL cohort, the subjects born after
266 1943 had a higher BMI⁽³⁶⁾ and lower alcohol consumption⁽²⁴⁾. Other studies in France suggest
267 differences in lifestyle and food consumption across birth cohorts born after the Second World
268 War^(51,52). Additionally, in the GAZEL cohort, men are far more numerous than women. While
269 the results seemed robust for men, for women it was sometimes unclear whether the estimates

270 were non-significant because the associations were weaker or because the sample size reduced
271 the power of the tests.

272 Measures of vegetables (or fruit) intake are sensitive to survey designs and definitions. For
273 example, the French official dietary guideline of five fruits and vegetables a day is met by only
274 13% of the French population⁽⁵⁾, but around 45% of the French have intakes $\geq 400\text{g/d}$ ^(5,53,54). In
275 2008, 86% of the French respondents (aged 12-75) to a nutritional survey had eaten vegetables
276 at least once during the 24-hour recall⁽⁵⁾. In contrast, in our occupational cohort, in 2009 one
277 out of three men and 55% of the women reported that they usually ate (cooked) vegetables
278 every day. Rather than food intake, our short, qualitative and self-administered FFQ captured
279 food habits, namely, whether a food group is usually consumed on a daily basis. Misreporting
280 is possible, but measurement error induces bias in a longitudinal analysis only if it is time-
281 dependent. In our case, the FFQ did not change from 1998 to 2009. The FFQ changed between
282 1990 and 1998 but our robustness check showed that this did not affect our results.

283 Measurement error should therefore not be a concern in this study. Finally, since we do not
284 control for total food intake or BMI, it is possible that the subjects increased their consumption
285 of unhealthy foods along with their vegetable consumption. While our results indicate that
286 vegetable consumption complied more and more with nutritional guidelines as subjects grew
287 older, it remains to be shown whether their whole diet became healthier.

288 Major strengths of our study were a large sample and a long follow-up period, with very low
289 attrition: more than 16,000 subjects were followed up over 19 years, with four repeated
290 questionnaires on their food habits. To our knowledge, there is no comparable study on
291 vegetable consumption and aging. Moreover, due to the initial recruitment of people aged over
292 40 in a large, state-owned company, virtually every subject had a continuous work history in
293 the company and had retired during the survey period. Multiple imputation of missing values
294 prevented loss of power and reduced bias due to non-response. We were also able to address
295 several potential sources of confusion.

296 In developed countries, people aged 60 years and more represent nearly 23% of the population.
297 They face a life expectancy of 23 years at age 60⁽⁵⁵⁾ with an increasing risk of nutrition-related
298 chronic diseases. While population aging is a major public health concern for industrialized
299 societies, our results also suggest that aging and retiring may have a positive effect on
300 vegetable intake among older adults. This supports the idea that dietary interventions at the
301 time of retirement may make an effective and sustainable⁽⁵⁶⁾ contribution to healthy aging.

302

303 *Acknowledgements*

304 We thank EDF-GDF and the Population-based Epidemiologic Cohorts Unit (UMS 011
305 INSERM/UVSQ) for their contribution to building and providing the GAZEL database. We are
306 grateful to Dr. Séverine Gojard (INRA ALISS), Éléonore Herquelot (INSERM UMS 011) and
307 Diane Cyr (INSERM UMS 011) for their support and comments.

308 *Affiliations*

309 UR1303 ALISS Alimentation et sciences sociales, INRA, Ivry sur Seine, France (Marie Plessz)
310 Inserm, Population-based Epidemiologic Cohorts Unit, UMS 011, Villejuif, France (Marie
311 Plessz, Alice Guéguen, Marcel Goldberg, Sébastien Czernichow, Marie Zins)
312 Versailles St-Quentin University, UMS 011, F-94807, Villejuif, France (Marcel Goldberg,
313 Sébastien Czernichow, Marie Zins)
314 Department of Nutrition, APHP, A. Paré hospital, Boulogne-Billancourt, France (Sébastien
315 Czernichow)

316 *Authors' contributions*

317 MP, MG, MZ, and SC designed the research. MP and AG analysed the data. MP wrote the
318 paper. MZ and MG provided detailed information on the data. All authors revised the text and
319 approved the final draft.

320 *Financial support*

321 None

322 *Conflicts of interest*

323 The authors declare no conflict of interest.

324

References

1. World Health Organization (2003) *WHO fruit and vegetable promotion initiative – a meeting report*. Geneva: World Health Organization.
2. Krebs-Smith SM, Guenther PM, Subar AF *et al.* (2010) Americans do not meet federal dietary recommendations. *J Nutr* **140**, 1832-1838.
3. Maguire ER, Monsivais P (2015) Socio-economic dietary inequalities in uk adults: An updated picture of key food groups and nutrients from national surveillance data. *Br J Nutr* **113**, 181-189.
4. Orfanos P, Naska A, Trichopoulos D *et al.* (2007) Eating out of home and its correlates in 10 european countries. The European prospective investigation into cancer and nutrition (EPIC) study. *Public Health Nutr* **10**, 1515-1525.
5. Escalon H, Bossard C, Beck F (editors) (2009) *Baromètre santé nutrition 2008*. Saint-Denis: INPES.
6. Welch N, McNaughton SA, Hunter W *et al.* (2009) Is the perception of time pressure a barrier to healthy eating and physical activity among women? *Public Health Nutr* **12**, 888-895.
7. Crawford D, Ball K, Mishra G *et al.* (2007) Which food-related behaviours are associated with healthier intakes of fruits and vegetables among women? *Public Health Nutr* **10**, 256-265.
8. Baker AH, Wardle J (2003) Sex differences in fruit and vegetable intake in older adults. *Appetite* **40**, 269-275.
9. Warde A, Cheng S-L, Olsen W *et al.* (2007) Changes in the practice of eating. *Acta Sociologica* **50**, 363-385.
10. Plessz M, Gojard S (2015) Fresh is best? Social position, cooking, and vegetable consumption in France. *Sociology* **49**, 172–190.
11. Bertail P, Caillavet F (2008) Fruit and vegetable consumption patterns: A segmentation approach. *American Journal of Agricultural Economics* **90**, 827-842.
12. Lindström M, Hanson BS, Wirfält E *et al.* (2001) Socioeconomic differences in the consumption of vegetables, fruit and fruit juices. *Eur J Public Health* **11**, 51-59.
13. Roos G, Johansson L, Kasmel A *et al.* (2001) Disparities in vegetable and fruit consumption: European cases from the north to the south. *Public Health Nutr* **4**, 35-43.
14. Plessz M, Gojard S (2013) Do processed vegetables reduce the socio-economic differences in vegetable purchases? A study in France. *Eur J Public Health* **23**, 747-752.

15. Dubuisson C, Lioret S, Touvier M *et al.* (2010) Trends in food and nutritional intakes of French adults from 1999 to 2007: Results from the INCA surveys. *Br J Nutr* **103**, 1035-1048.
16. Devine CM, Connors M, Bisogni CA *et al.* (1998) Life-course influences on fruit and vegetable trajectories: Qualitative analysis of food choices. *Journal of Nutrition Education* **30**, 361-370.
17. Bisogni CA, Jastran M, Shen L *et al.* (2005) A biographical study of food choice capacity: Standards, circumstances, and food management skills. *J Nutr Educ Behav* **37**, 284-291.
18. Furst T, Connors M, Bisogni CA *et al.* (1996) Food choice: A conceptual model of the process. *Appetite* **26**, 247-266.
19. Plessz M, Dubuisson-Quellier S, Gojard S *et al.* (2014) How consumption prescriptions change food practices. Assessing the role of household resources and life course events. *Journal of Consumer Culture* **Online first February 4, 2014**.
20. Westerlund H, Kivimäki M, Singh-Manoux A *et al.* (2009) Self-rated health before and after retirement in France (GAZEL): A cohort study. *Lancet* **374**, 1889-1896.
21. Vahtera J, Westerlund H, Hall M *et al.* (2009) Effect of retirement on sleep disturbances: The GAZEL prospective cohort study. *Sleep* **32**, 1459-1466.
22. Sjösten N, Kivimäki M, Singh-Manoux A *et al.* (2012) Change in physical activity and weight in relation to retirement: The French GAZEL cohort study. *BMJ Open* **2**, e000522.
23. Slingerland AS, van Lenthe FJ, Jukema JW *et al.* (2007) Aging, retirement, and changes in physical activity: Prospective cohort findings from the globe study. *Am J Epidemiol* **165**, 1356-1363.
24. Zins M, Guéguen A, Kivimaki M *et al.* (2011) Effect of retirement on alcohol consumption: Longitudinal evidence from the French gazel cohort study. *PLoS ONE* **6**, e26531.
25. Helldán A, Lallukka T, Rahkonen O *et al.* (2012) Changes in healthy food habits after transition to old age retirement. *Eur J Public Health* **22**, 582-586.
26. Hurd MD, Rohwedder S (2006) *Some answers to the retirement-consumption puzzle*. National Bureau of Economic Research.
27. Chung S, Popkin BM, Domino ME *et al.* (2007) Effect of retirement on eating out and weight change: An analysis of gender differences. *Obesity* **15**, 1053-1060.
28. Stancanelli E, Soest AV (2012) Retirement and home production: A regression discontinuity approach. *Am Econ Rev* **102**, 600-605.

29. Nooyens AC, Visscher TL, Schuit JA *et al.* (2005) Effects of retirement on lifestyle in relation to changes in weight and waist circumference in dutch men: A prospective study. *Public Health Nutr* **8**, 1266-1274.
30. Zantinge EM, van den Berg M, Smit HtA *et al.* (2014) Retirement and a healthy lifestyle: Opportunity or pitfall? A narrative review of the literature. *Eur J Public Health* **24**, 433-439.
31. Goldberg M, Leclerc A, Bonenfant Sb *et al.* (2007) Cohort profile: The GAZEL cohort study. *Int J Epidemiol* **36**, 32-39.
32. Le Port As, Gueguen A, Kesse-Guyot E *et al.* (2012) Association between dietary patterns and depressive symptoms over time: A 10-year follow-up study of the GAZEL cohort. *PLoS ONE* **7**, e51593.
33. Goldberg M, Zins M (2015) The gazel cohort: 20 000 volunteers for medical research. Questionnaires. <http://www.gazel.inserm.fr/en/documentation/questionnaires.html> (accessed 01/19/2015).
34. Kamphuis CBM, Giskes K, Bruijn G-jD *et al.* (2006) Environmental determinants of fruit and vegetable consumption among adults : A systematic review. *Br J Nutr*, 620-635.
35. Bove CF, Sobal J, Rauschenbach BS (2003) Food choices among newly married couples: Convergence, conflict, individualism, and projects. *Appetite* **40**, 25-41.
36. Dugravot A, Sabia S, Stringhini S *et al.* (2010) Do socioeconomic factors shape weight and obesity trajectories over the transition from midlife to old age? Results from the French GAZEL cohort study. *Am J Clin Nutr* **92**, 16-23.
37. White IR, Royston P, Wood AM (2011) Multiple imputation using chained equations: Issues and guidance for practice. *Stat Med* **30**, 377-399.
38. Sterne J, A. C. , White IR, Carlin JB *et al.* (2009) Multiple imputation for missing data in epidemiological and clinical research: Potential and pitfalls. *Br Med J* **338**, b2393
39. Goldberg M, Chastang JFo, Zins M *et al.* (2006) Health problems were the strongest predictors of attrition during follow-up of the GAZEL cohort. *J Clin Epidemiol* **59**, 1213-1221.
40. Rubin DB (1987) *Multiple imputation for nonresponse in surveys*. vol. 307. New York: Wiley.
41. Akbaraly TN, Brunner EJ (2008) Socio-demographic influences on trends of fish consumption during later adult life in the whitehall II study. *Br J Nutr* **100**, 1116-1127.
42. Mancino L, Todd J, Lin B-H (2009) Separating what we eat from where: Measuring the effect of food away from home on diet quality. *Food Policy* **34**, 557-562.

43. Guthrie JF, Lin B-H, Frazao E (2002) Role of food prepared away from home in the american diet, 1977-78 versus 1994-96: Changes and consequences. *J Nutr Educ Behav* **34**, 140-150.
44. Kjollesdal MR, Holmboe-Ottesen G, Wandel M (2011) Frequent use of staff canteens is associated with unhealthy dietary habits and obesity in a norwegian adult population. *Public Health Nutr* **14**, 133-141.
45. Lachat CK, Huybregts LF, Roberfroid DA *et al.* (2009) Nutritional profile of foods offered and consumed in a belgian university canteen. *Public Health Nutr* **12**, 122-128.
46. Roos E, Sarlio-Lähteenkorva S, Lallukka T (2004) Having lunch at a staff canteen is associated with recommended food habits. *Public Health Nutr* **7**, 53-61.
47. Lafay L (editor) (2009) *Étude individuelle nationale des consommations alimentaires : 2006-2007 (INCA 2)*. Paris: AFSSA.
48. Ribet C, Lang T, Zins M *et al.* (2001) Do cardiovascular risk factors in men depend on their spouses' occupational category? *Eur J Epidemiol* **17**, 347-356.
49. Tai T-o, Treas J (2013) Housework task hierarchies in 32 countries. *European Sociological Review* **29**, 780-791.
50. Guilbert P, Perrin-Escalon H (editors) (2002) *Baromètre santé nutrition 2002*. Paris: INPES.
51. Chauvel L, Schröder M (2014) Generational inequalities and welfare regimes. *Soc Forces* **Advance access**, sot156.
52. Hébel P, Recours F (2007) Effets d'âge et de générations : Transformation du modèle alimentaire. *Cahiers de nutrition et de diététique* **42**, 297-303.
53. Castetbon K, Vernay M, Malon A *et al.* (2009) Dietary intake, physical activity and nutritional status in adults: The French nutrition and health survey (ENNS, 2006–2007). *Br J Nutr* **102**, 733-743.
54. Lioret S, Dubuisson C, Dufour A *et al.* (2010) Trends in food intake in French children from 1999 to 2007: Results from the INCA (étude individuelle nationale des consommations alimentaires) dietary surveys. *Br J Nutr* **103**, 585-601.
55. United Nations Organization (2001) *World population ageing 1950-2050*. New York: UN.
56. Lara J, Hobbs N, Moynihan PJ *et al.* (2014) Effectiveness of dietary interventions among adults of retirement age: A systematic review and meta-analysis of randomized controlled trials. *BMC Med* **12**, 60.

Tables

Table 1. Baseline characteristics: descriptive statistics and number of missing values in men (n=12,942) and women (n=2,739) in the GAZEL cohort, France 1989-2009.

	Men		Women	
	Column % ^a	N missing	Column % ^a	N missing
Educational level ^d		238		85
Low	18.9		31.9	
Medium	53.4		51.9	
High	27.7		16.2	
Birth cohort		0		0
1939-1943	41.4		40.1	
1944-1948	58.6		59.9	
Usual place of lunch before retirement ^{b, d}		2,947		772
Out	53.7		69.9	
Home	46.3		30.1	
Age in 1990 ^c	45.9 (2.8)	0	45.9 (2.9)	0
Year of retirement ^{c, e}	1999.3 (3.6)	0	1999.9 (3.7)	0
Age at the time of retirement ^{c, e}	54.9(2.4)	0	55.5 (2.7)	0

Abbreviations: FFQ, food frequency questionnaire; SD, standard deviation.

^a Column percentages based on available cases (pairwise deletion).

^b Measured at the last FFQ before retirement.

^c Continuous variables are expressed as mean (standard deviation in parentheses).

^d Difference between men and women statistically significant (chi² test: p<0.001).

^e Difference between men and women statistically significant (t-test: p<0.001).

Table 2. Sample characteristics: descriptive statistics^a and number of missing values for the variables with repeated measurements for men (n=12,942) and women (n=2,739) in the GAZEL cohort, France 1989-2009.

	Men				Women			
	1990	1998	2004	2009	1990	1998	2004	2009
Outcome : daily consumption of vegetables ^b								
Yes (%)	17.7	19.9	28.6	33.4	31.0	36.1	50.8	55.3
N missing	1,616	3,235	3,050	3,282	372	810	709	750
Spouse ^b (column %)								
No spouse	6.4	7.7	9.0	10.5	21.7	24.9	26.5	29.7
Spouse works	52.5	46.3	28.1	13.0	72.5	43.7	15.4	4.4
Spouse inactive	41.1	46.0	62.9	76.5	5.7	31.4	58.2	65.9
N missing	1,555	3,239	2,941	3,216	361	816	702	738
Diet prescribed by a doctor								
Yes (%)	15.8	8.0	8.6	6.9	16.3	8.3	9.6	6.6
N missing	1,794	5,443	4,003	5,117	454	1,203	890	1,136

^a Column percentages based on available cases (pairwise deletion).

^b Difference between men and women statistically significant each year (chi² test: p<0.001).

Comment citer ce document :

Plessz , M. (Auteur de correspondance), Gueguen, A., Goldberg, M., Czernichow, S., Zins, M. (2015). Ageing, retirement and changes in vegetable consumption in France: findings from the prospective GAZEL cohort. *British Journal of Nutrition*, 114 (06, September 2015), 979-987. DOI : 10.1017/S0007114515002615

Table 3. Multivariate models for daily vegetable consumption, men and women in the GAZEL cohort, France 1989-2009 (odds-ratios with 95% CI).

	Men						Women					
	Model 1		Model 2		Model 3		Model 1		Model 2		Model 3	
	OR	CI	OR	CI	OR	CI	OR	CI	OR	CI	OR	CI
Age: slope 42-50	1.06	1.04, 1.08	1.06	1.03, 1.08	1.08	1.05, 1.11	1.08	1.04, 1.11	1.07	1.04, 1.11	1.1	1.05, 1.17
Age: slope 50-61	1.06	1.05, 1.07	1.06	1.05, 1.07	1.07	1.05, 1.09	1.07	1.04, 1.10	1.07	1.04, 1.10	1.09	1.05, 1.13
Age: slope 61-70	1.03	1.02, 1.05	1.03	1.02, 1.05	1.03	1.01, 1.04	1.04	1.01, 1.06	1.04	1.01, 1.06	1.03	1.00, 1.07
Retired	1.19	1.08, 1.31					1.16	0.97, 1.39				
Not retired & lunches at home ^a			1.28	1.18, 1.39	1.32	1.16, 1.49			1.16	0.99, 1.35	1.31	1.01, 1.70
Retired & lunched out ^a			1.31	1.17, 1.46	1.25	1.07, 1.46			1.21	1.01, 1.46	1.18	0.89, 1.56
Retired & lunched at home ^a			1.38	1.23, 1.54	1.31	1.10, 1.56			1.22	0.97, 1.53	1.27	0.90, 1.81
FFQ 1990 (ref: years 1998-2009)	1.42	1.26, 1.59	1.41	1.26, 1.58	1.46	1.23, 1.74	1.44	1.16, 1.78	1.43	1.16, 1.77	1.56	1.13, 2.16
Diet prescribed by a doctor	1.29	1.19, 1.41	1.29	1.19, 1.41	1.33	1.18, 1.49	1.7	1.46, 1.99	1.7	1.46, 1.99	1.59	1.27, 2.00
Born 1944-48 (ref: 1939-43)	1.12	1.04, 1.19	1.12	1.05, 1.19	1.11	1.00, 1.23	1.3	1.13, 1.48	1.3	1.13, 1.49	1.58	1.26, 1.97
Education:	1.15	1.06, 1.25	1.15	1.06, 1.25	1.14	1.01, 1.29	1.09	0.95, 1.24	1.09	0.96, 1.24	1.02	0.82, 1.27

Comment citer ce document :

Plessz , M. (Auteur de correspondance), Gueguen, A., Goldberg, M., Czernichow, S., Zins, M. (2015). Ageing, retirement and changes in vegetable consumption in France: findings from the prospective GAZEL cohort. *British Journal of Nutrition*, 114 (06, September 2015), 979-987. DOI : 10.1017/S0007114515002615

medium (ref: low)													
Education: high	1.21	1.11, 1.32	1.22	1.12, 1.33	1.14	0.99, 1.31	1.42	1.19, 1.69	1.43	1.20, 1.70	1.39	1.04, 1.85	
Spouse works (ref: no spouse)	1.35	1.21, 1.50	1.35	1.21, 1.50	1.43	1.20, 1.69	1.03	0.89, 1.18	1.02	0.89, 1.17	1.15	0.91, 1.46	
Spouse inactive	1.62	1.47, 1.79	1.6	1.45, 1.77	1.74	1.48, 2.06	1.25	1.09, 1.43	1.25	1.09, 1.43	1.26	1.02, 1.56	
N. of observations	51768		51768		17328		10956		10956		3420		
N. of subjects	12942		12942		4332		2739		2739		855		

Abbreviations: FFQ, food frequency questionnaire. GEE, Generalized Estimating Equations.

Logistic regressions using Generalized estimating equations.

Model 1: controls + age + retirement status. Completed data (50 imputations).

Model 2: controls + age + interaction between retirement status and usual place of lunch before retirement. Completed data (50 imputations).

Model 3: same as Model 2 on complete cases (listwise deletion: subjects with zero missing value).

^a Lunch : the most frequent place of lunch declared at the last FFQ before retirement. Reference category: not retired and lunches out

Comment citer ce document :

Plessz , M. (Auteur de correspondance), Gueguen, A., Goldberg, M., Czernichow, S., Zins, M. (2015). Ageing, retirement and changes in vegetable consumption in France: findings from the prospective GAZEL cohort. *British Journal of Nutrition*, 114 (06, September 2015), 979-987. DOI : 10.1017/S0007114515002615

Figure legends

Figure 1

Flowchart describing the selection of the study population from the GAZEL cohort, France 1989-2009.

^a Lost to follow-up: subjects asked not to be contacted again or stopped being employed by EDF-GDF before they retired.

Figure 2

Proportion of subjects eating vegetables daily according to sex and year of measurement among complete cases (listwise deletion, white), available cases (pairwise deletion, light grey) and after imputation (dark grey) in the GAZEL cohort, France 1989-2009, with 95% CI.

Figure 3

Predicted probabilities of consuming vegetables daily for men and women according to age, retirement status and usual place of lunch before retirement in the GAZEL cohort, France 1989-2009.

Predicted probabilities of eating vegetables everyday in the GAZEL cohort study are based on model 2 (imputed data) assuming that retirement occurred at age 56 years adjusted for educational level, (high versus low), current doctor-prescribed dieting status (no versus yes), current marital status (no versus yes), FFQ 1998 or later.

Solid line: predicted probability for people having most lunches away from home before retirement.


Shaded area: 95% confidence interval for the solid line.

Dashed line: before retirement, alternative prediction for a person who had most lunches at home.

Abbreviations: FFQ, food frequency questionnaire. GEE: Generalized Estimating Equations.

Figures


Figure 1



Comment citer ce document :

Plessz , M. (Auteur de correspondance), Gueguen, A., Goldberg, M., Czernichow, S., Zins, M. (2015). Ageing, retirement and changes in vegetable consumption in France: findings from the prospective GAZEL cohort. *British Journal of Nutrition*, 114 (06, September 2015), 979-987. DOI : 10.1017/S0007114515002615


Figure 2



Comment citer ce document :

Plessz, M. (Auteur de correspondance), Gueguen, A., Goldberg, M., Czernichow, S., Zins, M. (2015). Ageing, retirement and changes in vegetable consumption in France: findings from the prospective GAZEL cohort. *British Journal of Nutrition*, 114 (06, September 2015), 979-987. DOI : 10.1017/S0007114515002615

Figure 3



Comment citer ce document :

Plessz, M. (Auteur de correspondance), Gueguen, A., Goldberg, M., Czernichow, S., Zins, M. (2015). Ageing, retirement and changes in vegetable consumption in France: findings from the prospective GAZEL cohort. *British Journal of Nutrition*, 114 (06, September 2015), 979-987. DOI : 10.1017/S0007114515002615