



HAL
open science

Zoonotic emergence at the animal-environment-human interface: the forgotten urban socio-ecosystems

Gauthier Dobigny, Serge Morand

► To cite this version:

Gauthier Dobigny, Serge Morand. Zoonotic emergence at the animal-environment-human interface: the forgotten urban socio-ecosystems. Peer Community Journal, 2022, 2, pp.e79. 10.24072/pcjournal.206 . hal-04935721

HAL Id: hal-04935721

<https://hal.inrae.fr/hal-04935721v1>

Submitted on 7 Feb 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License



Peer Community Journal

Section: Infections

RESEARCH ARTICLE

Published
2022-12-09

Cite as
Gauthier Dobigny and Serge
Morand (2022) *Zoonotic
emergence at the
animal-environment-human
interface: the forgotten urban
socio-ecosystems*, Peer
Community Journal, 2: e79.

Correspondence
gauthier.dobigny@ird.fr

Peer-review
Peer reviewed and
recommended by
PCI Infections,
<https://doi.org/10.24072/pci.infections.100005>



This article is licensed
under the Creative Commons
Attribution 4.0 License.

Zoonotic emergence at the animal-environment-human interface: the forgotten urban socio-ecosystems

Gauthier Dobigny^{1,2} and Serge Morand^{3,4}

Volume 2 (2022), article e79

<https://doi.org/10.24072/pcjournal.206>

Abstract

Zoonotic emergence requires spillover from animals to humans, hence animal-human interactions. A lot has been emphasized on human intrusion into wild habitats (e.g., deforestation, hunting) and the development of agricultural and farming activities. However, the highly human-modified urban and peri-urban socio-ecosystems are also of great concern due to huge amounts of pet, domesticated and wild animals (e.g., birds, rodents and bats) that live in very close contact to very dense human populations. This adds to the existence of wet wildlife markets, urban parks and wastelands, zoos and even labs, where spillover from wildlife to humans may also occur. Furthermore, cities are transport hubs that form hotspots of import/export of living resources including animals, thus potentially promoting rapid and wide-scale spread of reservoir and vectors, hence pathogens, as well as pathogen admixture through viral recombination or bacterial plasmid exchanges. Finally, cities are deeply modified environments where living organisms, including reservoirs, vectors and pathogens, undergo strong selective pressures, thus opening the gate to evolutionary novelties, hence potential new infectious threats. As such, we believe urban socio-ecosystems should be paid more attention in terms of drivers of zoonotic emergence in humans needing adapted surveillance and mitigation. Accordingly, we propose and discuss several avenues of research and examples of actions that could be tested or generalized (e.g., focus on hotspots of emergence risks like informal settlements, wet markets or transport hubs; development of participative surveillance programs; shift towards inter-sectoral academic courses; massive investment into education and community information) in order to operationalize effective zoonotic surveillance. We advocate that this would allow ones to add emergence-preventive and early warning to usual outbreaks response strategies, thus significantly improving our collective ability to prevent zoonotic emergence and subsequent pandemics.

¹IRD, UMR CBGP – Montpellier, France, ²Institut Pasteur de Madagascar, Unité Peste – Antananarivo, Madagascar, ³CNRS, UMR MiVEGEC – Montpellier, France, ⁴Kasetsart University, Faculty of Veterinary Technology – Bangkok, Thailand

Peer Community Journal is a member of the
Centre Mersenne for Open Scientific Publishing
<http://www.centre-mersenne.org/>

e-ISSN 2804-3871



CENTRE
MERSENNE

Introduction

During the last decades, (re)emergence of zoonotic pathogens have led to an increasing number of human epidemics, the current Covid-19 pandemics being the most recent illustration. An important amount of observational studies as well as meta-analyses of the available scientific literature clearly point towards the association of human footprints on nature with animal-to-human spillover of infectious agents (e.g., Wolfe et al., 2005; Plowright et al., 2011; Jones et al., 2013; Gottdenker et al., 2014; Johnson et al., 2020; White & Razgour, 2020). In particular, wildlife hunting and consumption, encroachment into pristine areas, deforestation and land-use switches towards agricultural land as well as industrial livestock farming have been associated with the emergence of highly deleterious diseases such as plague, Human Immunodeficiency Virus (HIV), Ebola, Nipah, influenza, Severe Respiratory Acute Syndrome (SARS), Middle-East Respiratory Syndrome (MERS) and possibly Severe Respiratory Acute Syndrome Coronavirus 2 (SARS-Cov-2).

Although urbanization has also been identified as a possible factor for zoonotic spillover (e.g., Santiago-Alarcon & MacGregor-Fors, 2020; Muloi et al., 2022) and amplification of some human epidemics (e.g., Olival et al., 2017), much research and calls-for-action have rather focused on the importance of surveillance and prevention of zoonotic emergences along landscapes interfaces, with special emphasis on tropical forests (Wolfe et al., 2005; Guégan et al., 2019) and agricultural and farming industries (Jones et al., 2013). We of course fully agree with these recommendations. However, several of them failed to include cities as of potential importance for zoonotic emergence in humans *per se* (e.g., see Table 1 in the seminal paper of Jones et al., 2013; Kreuder-Johnson et al., 2015; Dobson et al., 2020; see also the very limited place given to infectious diseases in the recent review of urban-associated diseases by Flies et al., 2019). Yet, despite existing studies on urban-associated zoonoses (see examples below), we believe that further attention should be paid to urban, peri-urban and sub-urban areas where massive though not fully evaluated contacts between wildlife, domestic animal and inhabitants occur on a daily basis. Cities are also expected to be efficient incubators of evolutionary novelties. Such a situation is probably extremely favorable to repeated chains of already known or still unknown microbe transmission, thus opening the gate for spillover events and potential adaptation of some pathogens to human-to-human transmission, hence (re)emergence and large subsequent epidemics and pandemics.

Cities as the most important human habitat

Urbanization is not a recent phenomenon. Large cities have existed in the past, and their development has been associated with wide-scale zoonotic epidemics (Moore, 2020). During the last century, urbanization has accelerated and, for the first time in human History, more than half of humanity now lives in cities, with Asia and Africa being currently the most urbanizing continents (UN Habitat, 2019) (Fig. 1 A-B). In the last twenty years (2001-2018), built-up areas (BUA) have been increasing by 5,300,000 km² (i.e., more than 1,100 standard soccer fields per day) with 9 out of the 10 countries showing the highest BUA expansion being of low- or middle-income ones (Sun et al., 2020). During the same period, 22 large cities (i.e., >100km²) have displayed population growth of more than 2 million inhabitants, with 15, 5 and 1 of them in Asia, Africa and South America, respectively (Sun et al., 2020) (Fig. 1 C-F). In most high-income countries, anticipated and supervised urban planning usually translates into regular sanitation devices, thus allowing for a proper management of garbage and sewage. Unfortunately, in urban settings from some low- and middle-income regions, urbanization goes faster than authorities' actions and resources allocation, thus leading to the creation and sprawling of wide, precarious and often informal urban settlements (UN Habitat, 2019). As such, millions of urban and peri-urban dwellers live within unhealthy areas characterized by waste accumulation, persistent standing waters, absence of drinkable water and electricity supply, lack of education and health services (Firdaus, 2012). In 2018, an estimate of 4.2 billion people were urban dwellers, with 20-25% (up to >60% in Africa) of them living in informal settlements (UN Habitat, 2019).

Thus, urban habitats must be considered as the most important environment of human life, at least for the century to come, since 6.7 billion human beings are expected to live in cities by 2050 (UN Habitat,

2019). Some scholars even argue that urbanization and its developing peripheral forms (e.g. ex-urbanization, sub-urbanization) influences the fate of most non-urban areas, far beyond the sole core cities (see the concept of “extended urbanization”; Tzaninis et al., 2021). Due to this matter of fact as well as the obviously radical environmental changes, hence strong selective pressures, that urbanization puts on biodiversity, urban evolutionary ecology has recently received an important renewal of interest (Alberti, 2015; Rivkin et al., 2018; Alberti et al., 2020; Verrelli et al., 2022). However, it may be that some kind of academic European and North American leadership in the field induces a Northern hemisphere-biased perception (e.g., Parsons et al., 2017, 2020; Alberti et al., 2020; see Box 2 in Verrelli et al., 2022 showing strong under-representation of Global South researchers in urban evolutionary ecology; see Gandy, 2021 for an emphasis on urban health ecology), and so a very partial view of what animal-human interactions within cities may include. This is especially critical since near 90% of extant urbanization takes place in the tropics, essentially in Asia and Africa (UN Habitat, 2019).

Also, cities are sometimes perceived as places of improved health following the existence of sanitation systems, the concentration of health services offer as well as more diversified and abundant food supply. However, such shortcut view fails to take into account the frequent and deep spatial and/or socio-economic inequalities which may be source of strong infectious risks, and may obscure the obvious needs of a broader comprehension of what urbanization may represent in terms of health (Ahmed et al., 2019; Santiago-Alarcon & MacGregor-Fors, 2020). As an example, when one comes to infectious diseases and emergence risks, cities are places of millions of animal / human interactions which, by essence, are of major concern.




Figure 1: Changes in rural (A) and urban (B) population size from 1960 to 2050 (note the sharp increase in urban population in tropical regions) and in the number and size of large cities from (C) 1950, (D) 1975, (E) 2000 and (F) 2025 (data from the World Bank, <https://databank.worldbank.org/source/world-development-indicators>)

Cities as places for millions of close animal-human interactions

Animals such as mammals and birds are intuitively associated with natural habitats, remote and/or protected areas that remain almost free of human disturbance. Yet, urban and peri-urban habitats, though deeply modified and densely populated by people, may shelter a wide spectrum of biodiversity, including livestock and poultry (Fig. 2). This is particularly true in “extended cities” of low- and middle-income countries, many of which lie in the intertropical zone and encompass wide zones or urban, peri-urban, sub-urban (i.e., peripheral but developing independently of the core city) and ex-urban (i.e., peripheral zones dedicated to human activities such as trade centers, airports, etc ... that densify and finally sprawl) areas (Connolly et al., 2021).

Pet and domestic animals

Animal pets that may move around (such as dogs or cats) are plentiful in human settlements, including cities, and may play a major role in some diseases' ecology (e.g., Munide et al., 2021). For instance, felids are the definitive host of *Toxoplasma gondii*; expectedly, free-living cats play a major role in the spreading of this highly human prevalent parasite within cities (e.g., in Brazil, 40% of Sao Paulo cats seropositive; Mereiles et al., 2004). Stray dogs may reach very large numbers in some urban and peri-urban areas, thus sometimes greatly contributing to rabies virus circulation and transmission within cities (e.g., Tenzin et al., 2015; Castillo-Neyra et al., 2017). In addition, many urban dwellers in Africa, South America or Asia keep domestic animals such as cattle, pigs, goats, sheep, rabbits, poultries or pigeons in their households where they represent crucial food and economic resources. For instance, 56.6% of households in Banjul, Gambia, shelter poultry, while 47% of households in Dakar, Senegal, have small ruminants (Wilson, 2018). In Nairobi, Kenya, livestock is estimated to be over 1.3 million individuals: small-scale breeders represent 50-80% of cattle owners and are especially numerous in the informal settlements, with the latter harboring 5% of dairy animals of the urban sub-counties. There, infrastructures are described as “poor and dirty” while many inhabitants even declare to share their house with their cattle (Alarcon et al., 2017). Free-range scavenging throughout insalubrious areas such as dumping sites, backyard slaughtering as well as outdoor left-over of dead animals also contribute to poor hygienic conditions (Alarcon et al., 2017). As an example, Kenyan urban farmers may sell poultry carcasses to pig farmers when massive deaths occur (Alarcon et al., 2017). Such practices are of course risky since domestic animals have been shown to play a central role in known human-mammal-virus associations network (Wells et al., 2019) and to share the highest number of viruses with our species (Johnson et al., 2020). As an example, pigs and poultry are well known domestic animals which may be influenza virus-amplifying bridge species that favor transmission to human (reviewed in Chan et al., 2013). Poultry and livestock play an important role in *E. coli* population structure at the household level across Nairobi, and inhabitants in direct contact with them had higher chance to share the same *E. coli* isolates, thus pointing towards livestock involvement in human transmission (Muloi et al., 2022).

Wild animals including bats and rodents

Although this does not seem to translate into a higher relative number of zoonotic pathogens found in urban mammals, a recent meta-analysis suggests that wild urban-adapted mammal species tend to shelter an overall higher pathogen richness than their non-urban counterparts (Albery et al., 2021). Wild mammals, which were suggested as the source of transmission to humans of 86 (out of 95; 91%) zoonotic viruses (Kreuder-Johnson et al., 2015), are numerous in cities (Soulsbury & White, 2015). First, some cities sprawl directly into wild environments, thus favoring wildlife urban foraging. For instance, small but also very large South American or African cities develop through direct encroachment of tropical forests (e.g. Rio de Janeiro, Sao Paulo, Iquitos in South America; Franceville in Africa) and their fringes may be visited by a wide range of free-ranging wild animals. Many records (e.g., Barkham, 2017) demonstrate that sometimes large and unexpected animals regularly visit peri-urban zones, when not core cities: foxes in Europe, coyotes and mountain lions in North America, leopards in India, hyenas in Ethiopia, raccoons, elephants, monkeys, wild boars, etc. The list could be long and may reflect a growing phenomenon, i.e.

wildlife being constrained to cope with urban sprawling and/or taking advantage of easily accessible and abundant food resources through human-provide food or dumping sites (e.g., Newsome et al., 2015; see also Reed & Bonter, 2018 for a study on increased mammalian abundance on supplemented sites). This could have important consequences in terms of infectious diseases ecology. For instance, sub-urban residential habitats may provide increased food resources, thus fueling white-tailed deer densities, a keystone host for ticks that transmit borreliosis in the US (Diuk-Wasser et al., 2020). Raccoons, reservoirs for rabies virus, also display great eco-ethological adaptation to contrasted urban landscapes and resource availability levels, thus allowing them to occur throughout some North American cities (Gross et al., 2012). Second, some species (or populations of some species) have even adapted further and evolved a fully urban lifestyle; this includes a wide range of bird and mammal species. The latter are usually generalist, rapidly reproducing and highly adaptable species (Santini et al., 2019) which can proliferate and reach particularly elevated densities within urbanized environments (e.g., Moller et al., 2012; Tait et al., 2016). Among them, two taxonomic groups may be particularly critical for the zoonotic emergence risks, namely rodents and bats, whose urban communities may display very high densities and very large population size that are expected to greatly favor pathogen transmission (Lloyd-Smith et al., 2005).

Many studies have identified bats as major potential sources of zoonotic pathogens, especially RNA viruses (e.g., Olival et al., 2017; Nieto-Rabiela et al., 2019; Wells et al., 2019; reviewed in Hayman, 2016). Bats are usual urban dwelling mammals, including in large metropolises. In a meta-analysis, a (probably underestimated) total of 84 bat species were identified in Brazilian cities, 70% of which may roost (and 12% roosting exclusively) within human infrastructures, and at least 32.1% of which harbored zoonotic pathogens (Nunes et al., 2017). In many tropical towns, thousands of bats gather in trees that stand along streets, above houses, markets and squares where millions of people live and work every day (Ayivor et al., 2017). Among other species, the straw-coloured fruit bat (*Eidolon helvum*) is known to shelter several zoonotic viral lineages (e.g. Paramyxoviruses: Baker et al., 2013; Rotavirus: Esona et al., 2010; Lyssaviruses: Suu-Ire et al., 2017; Ebolaviruses: De Nys et al., 2018). The species was found to form one single continental Africa-wide panmictic population (Peel et al., 2013). This suggests that viral transmission could be facilitated across its whole geographical range (as supported by similar virus seroprevalence values observed in West, Central and East Africa; Peel et al., 2013) which encompasses many tropical African cities, including very large metropolises where colonies routinely reach several thousand individuals (Peel et al., 2016). In Eastern Australian cities, fruit bats of the *Pteropus* genus may form roosts of several thousands of individuals. While globally decreasing in Eastern Australia, the flying-foxes populations show a shift towards more urban lifestyle, thus increasing conflicts with humans (Westcott et al., 2015) and spillover of Hendra paramyxoviruses (Plowright et al., 2011). In addition, maybe unexpectedly, direct contacts between wild bats and human may not be rare within cities. As an example, 31 cases of human injuries due to bats were reported in the heart of Salvador, Brazil, between 2012 and 2015 (Santos et al., 2019), thus reflecting a poorly documented but potentially important source of spillover of bat-borne pathogens to humans. The West African Ebola epidemics, that affected >28,000 and killed >11,000 persons between 2013 and 2015, most probably started in a kid playing in close proximity to a tree that sheltered insectivorous bats (Mari-Saez et al., 2015). Though unknown, we suspect that such promiscuity-mediated bat-to-human transmission of RNA viruses occurrences may be quite high in many urban settlements in the world.

Although viral network analyses often identify bats as major sources of zoonotic viruses, especially RNA viruses (Nieto-Rabiela et al., 2019; Wells et al., 2019; Johnson et al., 2020), a higher proportion of zoonotic viruses are found in rodents than in bats (Nieto-Rabiela et al., 2019). The former ones are known to be reservoirs for a wide panel of known zoonotic pathogens (reviewed in Meerburg et al., 2009), with 400 million human infections by pathogens whose ecology involve rodents each year (Colombe et al., 2019). A very recent study on 50 zoonotic viruses ranked rodent-specific Lassa Mammarenavirus and Seoul Orthohantavirus as the first and fourth biggest risks of animal-to-human viral spill-over (Grange et al., 2021). Though representing only a limited fraction of global rodent specific diversity, the urban-adapted rodent species were recently demonstrated to share significantly more pathogens and parasites with humans than their non-urban counterparts (Gibb et al., 2020). Urban rodent assemblages are usually simplified and dominated by highly competitive, often invasive species, such as rats and mice that may replace native taxa within core cities (e.g. Dalecky et al., 2014; Garba et al., 2014b; Hima et al., 2019; Blasdell et al., 2022). They also consist in particularly competent species (Gibb et al., 2020). For instance,

Norway rats from New York were found to shelter an important viral diversity, including many undescribed viruses that belong to genera pathogenic for humans (e.g., Firth et al., 2014). In a similar manner, black rats are abundant in central Kuching, Malaysia, and host a wide panel of zoonotic pathogens, especially pathogens with environmental transmission (Blasdell et al., 2022). Among existing rodent-borne diseases with possible shifts to human-to-human transmission, the recent plague epidemics in Madagascar also reminds us about the risk associated with this deadly bacterial disease, especially in large and overcrowded urban areas (e.g., Majunga: Boisier et al., 1997; Antananarivo: Randremanana et al., 2019). Lassa hemorrhagic fever is usually considered as a rural disease, but its rodent reservoir, *Mastomys natalensis*, is widespread in many African cities (e.g. Garba et al., 2014b) where it may display quite high prevalence (e.g. close to 23.3% in N'Zerekore, Guinea; Karan et al., 2019). Although the 'historical strain' of SARS-CoV-2 was poorly infective in mice and rats due to its poor ability to bind to the murine ACE2 cellular receptor, some very recent pieces of evidence suggest that laboratory mice may be able to replicate some of its variants (at least the B.1.1.7 and other N501Y-carrying ones) in their lungs at high levels, suggesting that SARS-CoV-2 evolution may occur in commensal rodents as secondary hosts with potential subsequent spill-back to humans (Shuai et al., 2021; see also Wei et al., 2021 for a similar scenario involving the Omicron variant). Urban rodents were also shown to be important reservoirs for a wide panel of parasites: for instance, 15.3% of Durban rodents carry *Angiostrongylus cantonensis* (Archer et al., 2017); in Kuala Lumpur, 28.4% of rats are infected by *Hymenolepis nana* (Mohd Zaïn et al., 2012) while up to 94% of *H. diminuta* prevalence was observed in rats from some Ibadan areas, Nigeria (Akinbodae et al., 1981). Worms are not necessarily emerging pathogens, but they are under-documented though massively infecting urban human dwellers (Jariwala et al., 2017) and they are expected to remain a major public health issue in cities where rodents and humans coexist in high numbers. The absolute numbers of rats and mice in the different cities of the world remain astonishingly unknown, but they are obviously millions in many of them. In Chicago, the Bureau of Rodent Control of Chicago, USA, registered between 35,000 and 49,000 rodent-associated complaints per year between 2008 and 2017 (Murray et al., 2018). In only five boroughs of New York City, USA, 514 rodent bites were reported to the Department of Health in less than four years (Childs et al., 1998). Exceptionally high numbers of rat-seeing regularly trigger public reactions and subsequent press releases in most European and North American cities, thus reflecting the abundance of rodents in these urban settlements. Yet, in most of them, underground sewers provide some kind of physical separation between rats and human beings, thus probably greatly limiting direct human / rodent contacts. Unfortunately, in many low- or middle-income countries, sanitation systems are lacking or very poorly developed, and when they do exist, they often consist in large open sewers. In informal settlements, garbage management is very limited or inexistent, thus providing omnipresent, sustainable and open-air food resources for rodents all year long. As a direct consequence, rats, mice and other commensal rodents are plentiful in most poor urban settings, especially informal ones, where they regularly forage in close contact with people, and even nest inside houses. For instance, in the Pau da Lima favela, Salvador, Brazil, 109 out of 185 (59%) of households are infested by Norway rats (Costa et al., 2014). In Luang Prabang, Lao DR, 61.5% of interviewed inhabitants declared rodent sighting indoor within the week preceding the survey (Promkerd et al., 2008). During a two year-long study in Cotonou, Benin, small mammals were trapped in 91.8% of the investigated households (Dossou, Houéménou & Dobigny, unpublished results). In Niamey, Niger, rodents were trapped in 125 out of the 170 households investigated (73.5%), while rodent-associated issues were reported in 164 of them (96.5%) by the inhabitants (Garba et al., 2014a). In the same survey, 5% of interviewed inhabitants declared to have been bitten by wild rodents at least once at home (Garba et al., 2014a). In Osogbo, capital of Osun State in SW Nigeria where Lassa fever occurs each year, 55.4% of 500 informal settlements inhabitants declare seeing rodents moving around freely in their own houses, with 43.3% having observed at least one rodent in their home within the last 24 hours before the interview, and 20.2% having already eaten wild rats (Olalekan, 2015).

Altogether, these studies illustrate how abundant pathogen-carrying wildlife, especially small flying and terrestrial mammals, may be in cities where human density reaches its maximum. This sets the scene for billions of everyday wildlife-human interactions, hence spill-over opportunities (Fig. 2).




Figure 2: Graphical summary of the main hypothetical trends in terms of animal-human interaction and zoonotic processes that we believe are expected along a rural-urban gradient.

Captive animals, wildlife meat and wildlife meat markets

Kreuder-Johnson et al. (2015) pointed towards the risk of viral emergence in places where taxonomically diverse mammal species coexist in close promiscuity and where trans-species transmission is plausible. They mainly quoted such risks in zoos, labs -that usually lie within cities- as well as wildlife meat markets. Relying on current data of two viruses that display only rodent-to-human transmission in nature, namely the Lymphocytic Chorio-Meningitis Mammarenavirus and the Seoul Orthohantavirus, Childs et al. (2019) alerted on the importance of pet-, lab- and even transplant-associated risks of zoonotic viral spillover. Accidental human contaminations within research laboratories where infectious agents are investigated and manipulated are not infrequent (Peng et al., 2018), with 100-275 potential releases of pathogen each year in the US (Kaiser, 2014). Famous examples of lab-escapees or wrong uses include agents of SARS (Normile, 2004), anthrax, small pox, flu, etc (Owens, 2014). Among other hypotheses, a lab-escapee as a possible origin for the ongoing Covid-19 pandemics remains to be fully addressed (Harrison & Sachs, 2022).

Manipulation and consumption of wildlife meat have also been identified as a major risk for zoonotic spillovers (Wolfe et al., 2005; Katani et al., 2019), with wildlife markets being identified as potential spillover hotspots due to numerous interspecific contacts and poor sanitary and hygienic conditions (e.g., Saylor et al., 2021). Although wildlife consumption may be more important in rural areas, especially remote ones (Kurpiers et al., 2015), urban wildlife markets are not rare and surprisingly under-represented in studies that quantify the demand for wildlife meat (McNamara et al., 2019). Yet, urban demand for wildlife meat is underpinned by a wide panel of socio-cultural drivers (Shairp et al. 2016; Chausson et al., 2019) and is likely to increase in the coming decades (Kurpiers et al., 2015; McNamara et al., 2019). This may imply intensifying rural-to-urban socio-economic connections, from hunters to city-dwelling consumers. As an example, Kamins et al. (2011) showed that ca. 130,000 bats were sold each year along a 400 km-long economic network in Ghana (see other African examples in Kurpiers et al., 2015). It should be kept in mind that Ebola virus remains viable on monkey carcasses for at least a week, and viral RNA is detectable for up to 10 weeks (Prescott et al., 2015). In addition, many urban markets group living wild animals, thus probably greatly enhancing the chances of infectious pathogens spillover as well as the number of persons potentially involved, thus favoring epidemics early stages. For instance, it is plausible that SARS coronavirus has jumped from wild horseshoe bats to small carnivores, and then to humans in Chinese wet markets (Guan et al., 2003; Lau et al., 2005). Likewise, though still uncertain, the hypothesis of the emergence of SARS-CoV-2 at Wuhan wet market, China, would be another example of the potential role played by such places in zoonotic chains of transmission from wildlife to humans (Li et al., 2020).

Interestingly, many of the animals listed here above belong to a panel of mammalian lineages, namely Artiodactyla, Carnivora, Chiroptera, Eulipotyphla and Rodentia, that were all shown to have a critical role in zoonotic emergence, especially viral emergence (Olival et al., 2017; Nieto-Rabiela et al., 2019; Wells et

al., 2019). Such a biodiversity, together with very close promiscuity between animals and people, both at very high densities, result in constant interspecific contacts within cities around the world. Urban habitats may even offer the highest number of opportunities of wildlife / domestic animals / human contacts. This, of course, provides optimal conditions for pathogen exchanges between people, domestic and wild animals, hence zoonotic emergence and subsequent epidemics.

Arthropod vectors

Many infectious diseases are vector-borne (usually involving diptera, coleoptera, fleas, mites, lice or ticks; reviewed in Huttington et al., 2016; Semenza & Suk, 2018) with 40% of zoonotic viruses of wild animals being vector-transmitted to humans (Kreuder-Johnson et al., 2015) and the majority of emerging vector-borne diseases being transmitted by ticks and mosquitos (Swei et al., 2020). Although recent meta-analyses suggest that complex parasitological cycles may have higher chances than simple cycle to be disrupted within cities, especially in carnivores and primates (Werner & Nunn, 2020), many vectors of primary health importance show clear trends of adaptation to urban ecological conditions (e.g., Triatomine bugs: Parra-Henao et al., 2021; *Anopheles* spp.: Azrag & Mohammed, 2018; Takken & Lindsay, 2019; *Aedes* spp.: Wilke et al., 2020), sometimes taking advantage of remnant forests, urban parks and increasing city greenings (e.g. ticks and tick-borne diseases: Lydecker et al., 2019; Heylen et al., 2019; typhus-transmitting chiggers: Wulandary et al., 2021; *Aedes* spp.: Huynh & Minakawa, 2022). Accordingly, several studies have clearly pointed towards the existence of urban zoonotic cycles of vector-transmitted pathogens, thus suggesting increasing city-specific infectious risks (e.g., tick-borne *Borrelia*: Reye et al., 2010; mosquito-transmitted Zika virus: Musso & Gubler, 2016; Gandy, 2021; triatomine-borne *Trypanosoma cruzi*: Dye-Braumuller et al., 2019). The agents of murine (*Rickettsia typhi*) and cat-flea (*R. felis*) typhus were detected in rodents, tree-shrews and shrews from seven urban parks of Bangkok, Thailand (Rungroj et al., 2021), while urban parks have also been associated with 20.5% of human cases of scrub typhus (due to another Rickettsial bacterium *Orientia tsutsugamushi*) in Seoul, South Korea (Park et al., 2015).

The great heterogeneity of urban landscapes may deeply impact the arthropod vectors distribution and biology, including transmission characteristics such as competence, biting rate or incubation time, among others, thus making incidence of vector-borne infectious diseases quite evenly distributed within and among cities (review in LaDeau et al., 2015). The wildland-urban interface has been identified as a unique ecotone critical for ticks due to habitat heterogeneity hence increased ecological opportunities (the “intermediate disturbance hypothesis”), thus potentially leading to increased tick-borne diseases risk (Diuk-Wasser et al., 2020). There is little doubt that, in the decades to come, increasing processes of land-use change, especially peri-urbanization, sub-urbanization and ex-urbanization will be accompanied by a wide range of vector species proliferation in close proximity to high densities of both animals (see above) and humans, thus contributing to the growing range of infectious diseases that may emerge in core cities or their peripheries.

Cities as incubators of evolutionary novelties

By essence, cities are the socio-ecosystems with the highest human densities and population growth, and they are the receptacles for large human rural-urban or post-conflicts migrations; consequently, they are privileged places for epidemic amplification of human-human transmitted pathogens (Connolly et al., 2021). The 2014-2015 West African epidemics of Ebola fever, which was responsible for a death toll that had never been faced before, was essentially due to the unprecedented migration of human cases from rural areas into urban centers: at the outbreak peak, 51% of the cases identified in Liberia were from the capital city, Monrovia (Fallah et al., 2018; see also Wilkinson & Leach, 2015, and Gandy, 2021).

Importantly, even before human contamination, one also expects that very high numbers of available hosts, as may typically be the case within cities, translate into more intensive pathogen multiplication, hence genetic mutation and subsequent evolutionary changes. This should be particularly true for viruses, as illustrated by the SARS-CoV-2 variants evolution (Blackstone et al., 2020; Bandy & Wermer, 2021). As such, cities may be particularly favorable for human pathogen evolution simply due to higher number of infections, hence increased genetic change opportunities. Unfortunately, little is known about these aspects (Verrelli et al., 2022).

In addition, cities are very deeply modified habitats and display radically reorganized landscapes structure and connectivity, highly constrained hydrographic networks, usually very elevated levels of air, water and soil pollution, fully reshaped species communities and interactions, and even locally modified microclimates (Grimm et al., 2008). For instance, in central France, urban temperatures may be up to 3-4°C higher than their peripheral counterparts, and heat islands effect induces up to extra 3-4°C in some circumstances (Richard et al., 2021). While the average increase in global temperatures (i.e., 1.5-4°C in the coming decades) is expected to be associated with enhanced infectious risks (Hess et al., 2020), the resulting impacts of local climatic anomalies such as urban heat islands on processes like pathogen replication, reservoirs and vector population dynamics, hosts immunity and competence, hence infectious diseases transmission and trans-species zoonotic spillover, remain largely unknown (LaDeau et al., 2015; see e.g., Misslin et al., 2016, for an example on *Aedes* mosquito-transmitted dengue; Mironova et al., 2019, for an example on *Plasmodium vivax* malaria).

Another very poorly understood aspect of city life is the effect of air, soil and water pollution on genetic mutation rate in pathogens, and the consequences in terms of multiplication, contagiousness, host-specificity and virulence. Following overconsumption and subsequent release of antibiotics in the urban environment (Buelow et al., 2021), repeated exposure of pet and domestic animals as well as urban wildlife to a wide spectrum of antibacterial molecules is highly probable, and would greatly contribute to antibiotic-resistance evolution. Moreover, commensal microbiotes are inter-related with the environmental microbial biodiversity as well as widely impacted by lifestyle (Flandroy et al., 2018). Accordingly, although a lot remains to be described and understood, it is expected that urban lifestyle and habitats induce important changes in animal symbiotic microbial communities, thus potentially driving many evolutionary changes that may have consequences in terms of health (Flandroy et al., 2018; Mills et al., 2019), for instance through impacts on reservoir immune systems and their capacity of acquiring zoonotic agents (Pickard et al., 2017), or on vector ability to multiply and transmit pathogens (Yin et al., 2020).

From what precedes, urban socio-ecosystems' characteristics are expected to induce very strong selective pressures on living organisms, thus setting the stage for very new eco-evolutionary trajectories and scenarios in the decades to come (Alberti et al., 2020). This of course involves reservoir and vector species as well as zoonotic pathogens (Bradley & Alitzer, 2006). A recent meta-analysis suggested that urban environment may disrupt transmission cycle of some parasites, especially those displaying complex cycles (Werner & Nunn, 2020). However, some zoonoses tend to urbanize (e.g., leptospirosis: Lau et al., 2010; Blasdel et al., 2019; scrub typhus: Park et al., 2015), sometimes following the abundance of a few but highly competent reservoir species that are well-adapted to deeply human-modified habitats (Morand et al., 2019; Gibb et al., 2020). We still do not know much about evolutionary fate of zoonotic systems under urban selective regimes (Verrelli et al., 2022), and it may be cautious and wise to keep aware that one cannot really predict what may emerge from urban-driven novelties in terms of parasite diversity, hosts/parasite associations and, ultimately, zoonotic emergence.

Another crucial characteristic of cities is that they are transport hubs, so critical nodes for dissemination of reservoirs, vectors and pathogens, hence long-distance expansion of infectious diseases (Bradley & Alitzer, 2006; Alirol et al., 2011; Neiderud, 2015; e.g. plague: Zietz & Dunkelberg, 2004; hantavirus: Lin et al., 2012, Castel et al., 2021). This adds to the increasing supply of cities with living animals for food human consumption, as well as to the many urban-rural linkages through human movements, thus opening the gate to human- and animal-mediated import of rural pathogens into cities (Alirol et al., 2012; Ahmed et al., 2019; Connolly et al., 2021). Such multi-scale redistributions of reservoirs, vectors and pathogens (i.e., the "global biological cauldron"; Gandy, 2021) may greatly foster the evolution of host-pathogen interactions and dynamics to some extent that remain poorly understood and quantified.

In addition, increased exchanges between and within cities may deeply impact microbial interactions. First, they may favor exchange of genetic material between sometimes highly divergent lineages of bacteria. In particular, bacterial communities may share antibiotic-resistant plasmids, thus potentially conferring antibiotic multi-resistance to wide diversity of pathogens such as gastro-enteric ones (Partridge et al., 2018), especially at the human-livestock-wildlife interface (Hassel et al., 2019). Taking the risk associated with plague (Stenseth et al., 2008) as an example, the identification of antibiotic-resistant strains of *Yersinia pestis* in Malagasy black rats, including rats from the capital city Antananarivo (Cabanel et al., 2018) and the probable horizontal transfers within flea mid gut (Galimand et al., 2006) provide another example that undoubtedly deserves a particular attention. Second, cities may also favor the

coexistence of viruses, hence recombination between viral strains which facilitates host jumps and immune evasion (Vijaykrishna et al., 2015; Gandy, 2021). This may be particularly true for a wide panel of coronaviruses and influenza A viruses that actively circulate among bats and birds, respectively (reviewed in Chan et al., 2013). Much less documented but potentially worrisome, the presence of influenza A viruses in bats (Tong et al., 2012) as well as in 11% of Norway rats from Boston, USA (Cummings et al., 2019), probably deserves special attention, too.

Call for some actions

From what precedes, it appears that already known as well as still unknown microbes may arrive to, and get amplified in cities, or even emerge *in situ* following local evolutionary innovations as well as a very high number of daily interactions between humans and animals, both wild and domestic. There is little doubt that these multiple animal-human contacts provide excellent conditions for “microbial chatter” (extending the “viral chatter” concept proposed by Wolfe et al., 2005), i.e. iteration of animal-human human-animal transmission events until the pathogen’s adaptation opens the gate to human-to-human infectious routes and subsequent rapid and large-scale epidemics. As such, we believe that, together with the already well-identified risks associated with deforestation and zones of intensifying agricultural and livestock or poultry farming activities, urban areas as places of zoonotic emergence should be considered as another priority in terms of surveillance and risk mitigation measures. As a first step, this requires a better knowledge of the urban-specific zoonotic danger.

A better understanding of pathogen evolution at the animal-human interface within extended urban socio-ecosystems

Following the rapid urbanization and increase in urban lifestyle, the chances of zoonotic spillovers within urban and peri-urban regions will probably increase (Blasdell et al., 2019). However, the monitoring of all cities and urban zones is unrealistic. In order to prioritize urban zoonotic risk and guide surveillance networks’ implementation, research is needed to identify which urban features and/or activities are the most favorable ones for disease emergence. Indeed, a lot remains to be understood about the selective pressures and resulting adaptive trajectories that may shape urban-dwelling organisms’ ecology and evolution. For instance, recent experimental credit has been given to the pivotal role of selection on the distribution of *E. coli* and their resistance to antibiotics across an African city (the so-called “everything is everywhere, the environment selects” hypothesis; Muloi et al., 2022). Such studies are very useful for our understanding of pathogen ecology at the human-animal interface within urban environments and should be extended to other geographic contexts and wider panels of zoonotic pathogens. More generally, increased eco-evolutionary research is crucial in order to anticipate and to cope with the new zoonotic threats to come (Roche et al., 2020). In this context, moving towards studies that consider urban ‘extended’ socio-ecosystems in their full complexity and diversity would be very useful, as recently recommended in the fields of urban health (Flies et al., 2019), urban ecology (Alberti et al., 2020) and urban political ecology (Gandy, 2021; Tzaninis et al., 2021). In particular, it appears much too simplistic to consider cities as one single hence homogeneous ecosystem (e.g., urban vs. non-urban conditions; Blasdell et al., 2022), as what is sometimes done in meta-analyses that are yet published in very high standard scientific journals (e.g., Werner & Nunn, 2020; Albery et al., 2021). Cities are still too often perceived as essentially human socio-ecosystems that are to be opposed to natural environments. Clearly, health ecology investigations in cities must take extended urban socio-environmental heterogeneity and historical trajectories into much better consideration (Gandy, 2021; Blasdell et al., 2022; Verrelli et al., 2022).

From response to epidemics towards densified surveillance and prevention networks

The Covid19 crisis has deeply questioned social models and their resilience abilities. A recent UN report (UN Habitat, 2021) highlights the importance of pre-existing and well established institutionalized bodies and governance to set up and drive appropriate response policies against the epidemics in cities of the world. It also shows that, in addition to task forces that were created *ad hoc*, local networks drawn from preceding crises were successfully re-leveraged in some countries (e.g., DRC and its Ebola experience). However, it may be that the Covid19 pandemics have exacerbated some limits of response-based systems that may be insufficient to respond quickly enough in our extremely connected world, and should be

supplemented by prevention-targeting actions. Such a shift towards prevention rather than sole response is not a new idea (e.g., Patz et al., 2004), but its operationalization through inter-sectorial One Health approaches still remains limited.

From a surveillance perspective (in cities, but also in other socio-environments), human sciences should take the scene early in the process of building One Health actions. In particular, they should have a leading role in showing biomedical experts and health ecologists which socio-ecosystems (hence where, when, who and what) to investigate from the biological and medical perspectives. Indeed, one priority should be to identify fine-scale habitats and associated uses as well as historical, demographic and socio-environmental dynamics that are associated with the most extensive microbial interaction and evolution, as well as with human / animal contacts that are the most significant in terms of spillover potential. From there, “sentinel spaces” and “sentinel uses” (rather than sentinel species) could be targeted for the monitoring of microbial chatting, thus constituting a solid science-guided basis for emergence-preventive strategies. For instance, a recent surveillance system of 15 zoonotic diseases was implemented and evaluated in three counties of Western Kenya where husbandry practices are currently moving from subsistence to intensive market-oriented production (Falzon et al., 2019). It builds on hierarchical sampling protocols of sentinel markets, slaughterhouses and human health centers. Like others (e.g., MonkeyFeverRisk on tick-borne viral Kyasanur Forest Disease that affects low-income communities in the Western Ghats, India; Burthe et al., 2021; <https://www.monkeyfeverrisk.ceh.ac.uk/>), this remarkably integrated device involves veterinarians and medical doctors, founded on a good understanding of the local socio-economic context and bets on a long-term confidence relationship between academic, operational and community stakeholders. Likewise, in-depth analysis of urban livestock keeping uses and supply chains permits the clear understanding of economic constraints and opportunities that characterize urban livestock economy in Southern cities, thus pointing towards an urgent need to accompany urban farmers in animal management in order to reach both food security and safety (Alarcon et al., 2017). Other emblematic One Health programs were conducted on rabies (such as in Chad, Mali and Côte d’Ivoire: reviewed in Lechenne et al., 2021) and have led to important improvement in terms of both human and animal surveillance as well as of showing the interest of some strategies over others. For instance, they were useful in identifying some sources of improvement, such as the good reporting of data in main cities -where labs usually exist- that contrasts with the mass of knowledge and data that are collected in peripheral areas but fail to reach national hence international monitoring systems (Lechenne et al., 2021). Robust epidemiologically structured sampling procedures focusing on the human-animal interface within cities are also very informative to identify scales at which microbial spillover may occur (e.g., household vs. city; Muloi et al., 2022), and so at which scales preventive actions may be tailored. Consequently, this type of long-term inter-sectorial programs should be strongly encouraged by funding bodies and then put forward in order to convince socio-economic stakeholders and decision makers to invest in their generalization and institutionalization.

More specifically, interventional research could fuel similar programs by targeting strategic urban sites (e.g., wildlife meat markets, informal settlements) as well as urban/non-urban interfaces. The new forms of extended urbanization, that usually take place at (peri-urbanization), when not beyond (sub- and ex-urbanization) core cities’ margins, especially in the global South, should also be carefully scrutinized due to potentially high levels of wild and domestic biodiversity together with dense, rapidly growing and vulnerable human populations (Diuk-Wasser et al., 2020; Connolly et al., 2021).

In addition, as discussed above, entry points and transport hubs also deserve a particular attention. For instance, dedicated actions could be launched to train and enroll transport and trans-border stakeholders (e.g. seaport and port industrial networks, custom services, import/export firms), taking advantage of their operational “savoir-faire” and tools which would probably be of great help in the surveillance of transport means, hubs and networks, hence of the prevention of long distance dissemination of zoonotic agents. In essence, this is the purpose of the International Health Regulation (WHO, 2005) that is in fact inherited from quite old international debates (Howard-Jones, 1975) and texts – e.g., the 1903 International Sanitary Convention already preconized rat control on ships, and desinsectization was added in the subsequent versions in 1912 and 1923 (Lembrez, 1966). It could be interesting to capitalize on concepts and approaches developed for biological invasion management (e.g., see the special issue of Biological Invasions on Early Detection and Rapid Response of invasive organisms, eds. Meyerson, A. & Simberloff, D., 2020). For instance, collaborative platforms of surveillance managed jointly by academic, institutional and operational

stakeholders could be implemented within transport stations, ports and seaports (e.g., Port Platform of Environmental Surveillance currently settled in Cotonou seaport, Benin; Adamjy et al., 2020).

A special role for local community-based networks?

Prevention and early warning before spillover and subsequent spread of emerging pathogens cannot be achieved without a dense surveillance network at the closest from the field. The only realistic option is to take advantage of already existing operational networks, starting with local veterinary and health care centers and staffs. Surveying cities, where human / animal interactions are daily and widespread, requires massive human resources. In such a context, community-based early alert systems intuitively appear as a promising way to leverage surveillance networks within urban settings (e.g., Toledo-Romani et al., 2007; Costa et al., 2014; Urioste-Stone et al., 2015; Carvalho Parente et al., 2017; Ouédraogo et al., 2018; Valdez et al., 2020). This requires strong inhabitants' trust in order to reach ownership, hence long-term enrollment and implication in participative actions which remains a critical challenge (e.g., Webb Hooper et al., 2019; Valdez et al., 2020). The latter should be facilitated by political goodwill and supportive leadership through a national structure dedicated to One Health actions (Falzon et al., 2019). Participative approaches are promising and further experimental attempts should be made in a wide panel of socio-cultural contexts, especially urban ones, to consolidate the corpus of knowledge and "savoir-faire" in such actions which may be particularly challenging in cities (Ridde et al., 2014; Saré et al., 2018). Indeed, though usually beneficial, community-based approaches to health improvement may have limited or incomplete effects (Nickel & von dem Knesebeck, 2020). For instance, limitations of community-based actions against urban vectors or reservoirs may arise from the lack or poor engagement of local champions/leaders, especially in absence of financial incentives (e.g., Espino et al., 2012; Saré et al., 2018 for examples in Cuba, Philippines and Burkina-Faso). This may sometimes be overcome through more adapted communication, iterative processes, local capacity building as well as political support and investment in order to help the poorest communities to overpass the socio-economic constraints that prevent them to self-adapt for disease prevention purposes (Nickel & von dem Knesebeck, 2020; Asaaga et al., 2021). Such debates highlight the crucial importance of the evaluation of these types of interventional researches (Ridde et al., 2020), especially within cities where heterogeneous socio-economic contexts may require specific community-based approaches (e.g., formal vs. informal areas: Ridde et al., 2014) and where the choice of the leaders may be particularly critical (Espino et al., 2012; Saré et al., 2018). This implies that more knowledge, hence research, about the real efficiency and adaptability of participatory programs aiming at reducing animal-human contacts and mitigating zoonotic risks within urban areas are conducted. It also raises the issue of rapid sharing of knowledge from the communities to the Health authorities in charge, whoever they might be, in order to trigger immediate and coordinated higher-level responses.

Wide-scale empowerment on the One Health concept

Thinking forwards, as many people as possible need to be empowered, especially young generations. To do so, generalizing the teaching of One Health concept and approach in educational and academic courses is probably the only way to achieve massive, efficient, wide-scale and long term capacity building and awareness raising at a reasonable cost. In universities, this will mean to adapt existing or create *de novo* courses in a transdisciplinary manner, far beyond the sole biomedical and environment sciences, i.e. including socio-economical, anthropological, urban planning, logistical and political aspects. Accordingly, the creation of Centers (or Networks) of Excellence in Ecology and Health Research and Training housed in regional universities and/or research institutes had already been advocated more than fifteen years ago (Patz et al., 2004); unfortunately, to our knowledge, they are still rare (e.g., One Health Center of Excellence, University of Florida, USA; One Health Frontier Graduate School of Excellence, Hokkaido University, Japan; One Health Research Center of Excellence for Research and Training, Sun Yat-sen University, Guangzhou, China; One Health European Joint Program, European Union network).

Large-scale education is also a critical issue to question and potentially reshape people's habits and uses since the latter factors are central in the zoonotic spill-over process (Epstein, 2020; Hassel et al., 2021). This will require huge funding investments in education in universities and schools as well as at the community level (e.g., Hancke & Suarez, 2014). Importantly, it should benefit from significant inputs from socio-anthropologists, linguists (to transfer specialized concepts and semantics to local languages), experts in education and communication sciences.

For such investments to be prioritized, an important lobbying towards national and local political leaders will be necessary. Media support through adapted channels (e.g., national TVs, community radios, social media, social workers' campaigns) will be another important way to reach a maximum of people and stakeholders.

Urban management and poverty reduction

Last but not least, one must recognize that many zoonotic emergence risks are associated with poverty. As a consequence, efficient prevention and early response against nascent epidemics cannot be achieved without full political involvement and support. Decline of infectious diseases in US and European cities within the last two centuries probably owes a lot to concerted efforts of urban planners and public health stakeholders who work together towards wide-scale sanitation and waste management services, something that has been rather rare in the Global South where attention is rarely paid to infectious diseases, especially zoonoses, when it comes to urban planning (Ahmed et al., 2019). Obviously, eradication of insalubrious urban areas would immediately translate into great improvement in terms of infectious diseases burden (G. Suzan, in Robin, 2021). As such, following the One Health approach, a real dialogue between urban policy makers and health stakeholders is required to make significant progress towards better sanitation and living conditions, hence decreased zoonotic risks, for a vast number of poor urban dwellers.

Acknowledgements

We are grateful to many of our colleagues and students for the fascinating discussions that allowed us to fuel the present manuscript. The final version of this manuscript greatly benefitted from the inputs of three reviewers, namely E. Dumonteil, N.L. Gottdenker and a third anonymous contributor.

Preprint version 3 of this article has been peer-reviewed and recommended by Peer Community in Infections (<https://doi.org/10.24072/pci.infections.100005>).

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Funding

SM is supported by the French ANR project FutureHealthSEA (ANR-17-CE35-0003-01). GD is supported by the SCARIA and HANTARIS projects funded by the Belmont Forum (CRA *Pathways to Sustainability*) and the French National Agency for Sanitary Security of Food, Environment and Work (ANSES), respectively.

References

- Acharya KP, Acharya N, Phuyal S, Upadhyaya M, Lasee S (2020) One-health approach: a best possible way to control rabies. *One Health*, **10**, e100161. <https://doi.org/10.1016/j.onehlt.2020.100161>
- Adamjy T, Aholou S, Mourlon M, Dobigny G (2020) La gouvernance des risques liés aux invasions biologiques : l'exemple du Bénin. *Sciences, Eaux et Territoires*, **HS70**. <https://doi.org/10.3917/set.hs1.0001j>
- Ahmed S, Davila DJ, Allen A, Mordechai H, Tacoli C, Fèvre EM (2019) Does urbanization make emergence of zoonosis more likely? Evidence, myths and gaps. *Environment and Urbanization*, **31**, 443-460. <https://doi.org/10.1177/0956247819866124>
- Akinbodae O, Dipeolu O, Ogunji F, Adegoke G (1981) The parasites obtained and bacteria isolated from house rats (*Rattus rattus* Linnaeus 1758) caught in human habitations in Ibadan, Nigeria. *International Journal of Zoonoses*, **8**, 26-32.

- Alarcon P, Fèvre EM, Muinde P, Murungi PM, Kiambi S, Akoko J, Rushton J (2017) Urban livestock keeping in the city of Nairobi: diversity of production systems, supply chains and their disease management and risks. *Frontiers in Veterinary Sciences*, **4**, e171. <https://doi.org/10.3389/fvets.2017.00171>
- Alberti M (2015) Eco-evolutionary dynamics in an urbanizing planet. *Trends in Ecology and Evolution*, **30**, 114-126. <https://doi.org/10.1016/j.tree.2014.11.007>
- Alberti M, Palkovacs EP, Des Reoches S, De Meester L, Brans KI, Govaert L, Grimm NB, Harris NC, Hendry AP, Schell CJ, Szulkin M, Munshi-South J, Urban MC, Verrelli BC (2020). The complexity of urban eco-evolutionary dynamics. *BioScience*, **70**, 772-793. <https://doi.org/10.1093/biosci/biaa079>
- Albery GF, Carlson CJ, Cohen LE, Eskew EA, Gibb R, Ryan SJ, Sweeny AR, Becker DJ (2021) Urban-adapted mammal species have more known pathogens. *bioRxiv preprint*. <https://doi.org/10.1101/2021.01.02.425084>
- Alirol E, Getaz L, Chappuis F, Loutan L (2011) Urbanisation and infectious diseases in a globalised world. *Lancet Infectious Diseases*, **10**, 131-141. [https://doi.org/10.1016/S1473-3099\(10\)70223-1](https://doi.org/10.1016/S1473-3099(10)70223-1)
- Archer C, Appleton C, Mukaratirwa S, Lamb J, Schoeman M (2017) Endo-parasites of public health importance recovered from rodents in the Durban metropolitan area, South Africa. *South African Journal of Infectious Diseases*, **1**, 1-10. <https://doi.org/10.1080/23120053.2016.1262579>
- Asaaga FA, Rahman M, Kalegowda SD, Mathapati J, Savanur I, Srinivas PN, Seshadri T, Narayanswamy D, Kiran SK, Oommen MA, Young JC, Purse BV (2021) 'None of my ancestors ever discussed this disease before!': how disease information shapes adaptive capacity of marginalized rural populations in India. *PLoS Neglected Tropical Diseases*, **15**, e0009265. <https://doi.org/10.1371/journal.pntd.0009265>
- Ayivor JS, Ohemeng F, Tweneboah Lawson E, Waldman L, Leach M, Ntiamoah-Baidu Y (2017) Living with bats: the case of Ve Golokuati township in the Volta Region of Ghana. *Journal of Environment and Public Health*, e5938934. <https://doi.org/10.1155/2017/5938934>
- Azrag RS, Mohammed BH (2018) *Anopheles arabiensis* in Sudan: a noticeable tolerance to urban polluted larval habitats associated with resistance to Temephos. *Malaria Journal*, **17**, 204. <https://doi.org/10.1186/s12936-018-2350-1>
- Baddal B, Kahir N (2020) Co-infection of MERS-CoV and SARS-CoV-2 in the same host: a silent threat. *Journal of Infections and Public Health*, **12**, 1251-252. <https://doi.org/10.1016/j.jiph.2020.06.017>
- Baker KS, Tood S, Marsh GA, Cramer G, Barr J, Kamins AO, Peel AJ, Yu M, Hayson DTS, Nadjm B, Mtve G, Amos B, Reyburn H, Nyarko E, Suu-Ire R, Murica PR, Cunningham AA, Wood JLN, Wang LF (2013) Novel, potentially zoonotic paramyxoviruses from the African straw-colored fruit bat *Eidolon helvum*. *Journal of Virology*, **87**, 1348-1358. <https://doi.org/10.1128/JVI.01202-12>
- Bandoy D, Weimer B (2021) Analysis of SARS-CoV-2 genomic epidemiology reveals disease transmission coupled to variant emergence and allelic variation. *Scientific Reports*, **11**, e7380. <https://doi.org/10.1038/s41598-021-86265-4>
- Barkham P (2017) Urban beasts: how wild animals have moved into cities. *The Guardian*, 20th May, 2017. <https://www.theguardian.com/environment/2017/may/20/urban-beasts-how-wild-animals-have-moved-into-cities> (accessed 29 Nov 2021)
- Blackstone NW, Blackstone SR, Berg AT (2020) Variation and multilevel selection of SARS-CoV-2. *Evolution*, **74**, 2429-2434. <https://doi.org/10.1111/evo.14080>
- Blasdell KR, Morand S, Perera D, Firth C (2019) Association of rodent-borne *Leptospira* spp. with urban environments in Malaysian Borneo. *PLoS Neglected Tropical Diseases*, **13**, e0007141. <https://doi.org/10.1371/journal.pntd.0007141>
- Blasdell KR, Morand S, Laurance SGW, Doggett SL, Hahs A, Trinh K, Perera D, Firth C (2022). Rats and the city: implications of urbanization on zoonotic disease risk in Southeast Asia. *Proceedings of the National Academy of Sciences USA*, **119**, e2112341119. <https://doi.org/10.1073/pnas.2112341119>
- Bradley CA, Alitzer S (2006) Urbanization and the ecology of wildlife diseases. *Trends Ecology and Evolution*, **22**, 95-102. <https://doi.org/10.1016/j.tree.2006.11.001>
- Boisier P, Rasolomaharo M, Ranaivoson G, Rasoamanana B, Rakoto L, Andrianirina Z, Andriamahefazafy B, Chanteau S (1997) Urban epidemic of bubonic plague in Majunga, Madagascar: epidemiological aspects. *Tropical Medicine and International Health*, **2**, 422-427. <https://doi.org/10.1111/j.1365-3156.1997.tb00163.x>

- Buelow E, Ploy MC, Dagot C (2021) Role of pollution on the selection of antibiotic resistance and bacterial pathogens in the environment. *Current Opinion in Microbiology*, **64**, 117-124. <https://doi.org/10.1016/j.mib.2021.10.005>
- Burthe SJ, Schäfer SM, Asaaga FA, Balakrishnan N, Mudassar Chanda M, Darshan N, Hoti SL, Kiran SK, Sechradi T, Srinivas P, Vanak AT, Purse BV (2021) Reviewing the ecological evidence base for management of emerging tropical zoonoses: Kyasanur Forest Disease in India as a case study. *PLoS Neglected Tropical Diseases*, **15**, e0009243. <https://doi.org/10.1371/journal.pntd.0009243>
- Cabanel N, Bouchier C, Rajerions M, Carniel E (2018) Plasmid-mediated doxycycline resistance in a *Yersinia pestis* strain isolated from a rat. *International Journal of Antimicrobial Agents*, **51**, 249-254. <https://doi.org/10.1016/j.ijantimicag.2017.09.015>
- Carvalho Parente C, Bezerra FSM, Parente PI, Dias-Neto RV, Xavier SCC, Ramos Jr AN, Carvahlo-Costa FA, Lima MM (2017). Community-based entomological surveillance reveals urban foci of Chagas disease vectors in Sobral, State of Ceará, Northeastern Brazil. *PLoS ONE*, **12**, e0170278. <https://doi.org/10.1371/journal.pone.0170278>
- Castillo-Neyra R, Zegarra E, Monroy Y, Bernedo RF, Cornejo-Rosello I, Paz-Soldan VA, Levy MZ (2017) Spatial association of canine rabies outbreak and ecological urban corridors, Arequipa, Peru. *Tropical Medicine and Infectious Diseases*, **2**, e38. <https://doi.org/10.3390/tropicalmed2030038>
- Castel G, Kant R, Badou S, Etougbétché J, Dossou HJ, Gauthier P, Houéménou G, Smura T, Sironen T, Dobigny G (2021) Genetic characterization of Seoul virus in the seaport of Cotonou, Benin. *Emerging Infectious Diseases*, **27**, 2704-2706. <https://doi.org/10.3201/eid2710.210268>
- Chan JF-W, Kai-Wang T, Tse H, Jin D-Y, Yuen K-Y (2013) Interspecies transmission and emergence of novel viruses: lessons from bats and birds. *Trends in Microbiology*, **21**, 544-555. <https://doi.org/10.1016/j.tim.2013.05.005>
- Chausson AM, Rowcliffe JM, Escouflaire L, Wieland M, Wright JH (2019) Understanding the sociocultural drivers of urban bushmeat consumption for behavior change interventions in Pointe Noire, Republic of Congo. *Human Ecology*, **47**, 179-191. <https://doi.org/10.1007/s10745-019-0061-z>
- Childs JE, McLafferty SL, Sadek R, Miller GL, Khan AS, DuPree ER, Advani R, Mills JN, Glass GE (1998) Epidemiology of rodent bites and prediction of rat infestation in New York City. *American Journal of Epidemiology*, **148**, 78-87. <https://doi.org/10.1093/oxfordjournals.aje.a009563>
- Childs JE, Klein SL, Glass GE (2019) A case study of two rodent-borne viruses: not always the same old suspects. *Frontiers in Ecology and Evolution*, **7**, e35. <https://doi.org/10.3389/fevo.2019.00035>
- Cleaveland S, Lankester F, Townsend S, Lembo T, Hampson K (2014) Rabies control and elimination: a test case for One Health. *Veterinary Records*, **175**, 188-193. <https://doi.org/10.1136/vr.g4996>
- Colombe S, Jancloues M, Rivière A, Bertherat E (2019) A new approach to rodent control to better protect human health. *WHO Weekly Epidemiological Records*, **17**, 197-212. <https://apps.who.int/iris/handle/10665/312103>
- Connolly C, Keil R, Ali SH (2021) Extended urbanisation and the spatialities of infectious disease: demographic change, infrastructure and governance. *Urban Studies*, **58**, <https://doi.org/10.1177/0042098020910873>
- Costa F, Ribeiro GS, Felzemburgh RDM, Santos N, Reis RB, Santos AC, Bittencourt D, Fraga M, Araujo WN, Santana C, Childs JE, Reis MG, Ko AI (2014) Influence of household rat infestation on *Leptospira* transmission in the urban slum environment. *PLoS Neglected Tropical Diseases*, **8**, e3338. <https://doi.org/10.1371/journal.pntd.0003338>
- Cummings CO, Hill NJ, Puryear WB, Rogers B, Mukherjee J, Leibler JH, Rosenbaum MH, Runstadler JA (2019) Evidence of Influenza A in wild Norway rats (*Rattus norvegicus*) in Boston, Massachusetts. *Frontiers in Ecology and Evolution*, **7**, e36. <https://doi.org/10.3389/fevo.2019.00036>
- Dalecky A, Bâ K, Piry S, Lippens C, Diagne CA, Kane M, Sow A, Diallo M, Niang Y, Konecny A, Sarr N, Charbonnel N, Granjon L, Duplantier J-M, Brouat C (2015) Range expansion of the invasive house mouse *Mus musculus domesticus* in Senegal, West Africa: a synthesis of trapping data over three decades, 1983-2014. *Mammal Review*, **45**, 176-190. <https://doi.org/10.1111/mam.12043>
- De Nys DM, Kingebeni PM, Keita AK, Butel C, Thaurignac G, Villabona-Arenas C-J, Lemarcis T, Geraerts M, Vidal N, Esteban A, Bourgarel M, Roger F, Leendertz F, Diallo R, Ndimbo-Kumugo S-P, Nsio-Mbeta J, Tagg N, Koivogui L, Toure A, Delaporte E, Ahuka-Mundeye S, Muembe Tamfum J-J, Mpoudi-Ngole E, Ayouba A, Peeters M (2018) Survey of Ebola viruses in frugivorous and insectivorous bats in Guinea,

- Cameroon and the Democratic Republic of the Congo, 2015-2017. *Emerging Infectious Diseases*, **24**, 2228-2240. <https://doi.org/10.3201/eid2412.180740>
- Diuk-Wasser MA, VanAcker MC, Fernandez MP (2020) Impact of land use changes and habitat fragmentation on the eco-epidemiology of tick-borne diseases. *Journal of Medical Entomology*, **58**, 1546-1564. <https://doi.org/10.1093/jme/tjaa209>
- Dobson AP, Pimm SL, Hannah L, Kaufman L, Ahumada JA, Ando AW, Bernstein A, Busch J, Daszak P, Engelmann J, Kinnaird MF, Li BV, Loch-Tenzelides T, Lovejoy T, Nowak K, Roehrdanz PR, Vale MM (2020) Ecology and economics for pandemic prevention. *Science*, **369**, 379-381. <https://doi.org/10.1126/science.abc3189>
- Dye-Braumuller KC, Gorchakov R, Gunter SM, Nielsen DH, Roachell WD, Wheless A, Debboun M, Murray K, Nolan MS (2019) Identification of Triatomines and their habitats in a highly developed urban environment. *Vector Borne Zoonotic Diseases*, **19**, 265-273. <https://doi.org/10.1089/vbz.2018.2352>
- Epstein JH, Anthony SJ, Islam A, Kilpatrick AM, Ali Khan S, Balkey MD, Ross N, Smith I, Zambrana-Torrel C, Tao Y, Islam A, Lan Quan P, Olival KJ, Uddin Khan MS, Gurley ES, Hossein MJ, Field HE, Fielder MD, Brises T, Rahman M, Broder CC, Crameri G, Wang L-F, Luby SP, Lipkin WI, Daszak P (2020) Nipah virus dynamics in bats and implications for spillover to humans. *Proceedings of the National Academy of Sciences, USA*, **117**, 29190-29201. <https://doi.org/10.1073/pnas.2000429117>
- Esona MD, Mijatovic-Rustempasic S, Conrardy C, Tong S, Kuzmin IV, Agwanda B, Breiman BF, Banyai K, Niezgodna M, Rupprecht CE, Gentsch JR, Bowen MD (2010) Reassortant group A Rotavirus from straw-colored fruit bat (*Eidolon helvum*). *Emerging Infectious Diseases*, **16**, 1844-1852. <https://doi.org/10.3201/eid1612.101089>
- Espino PF, Marco J, Salazar NP, Salazar F, Mendoza Y, Velasco A (2012) Community-based dengue vector control: experiences in behavior change in Metropolitan Manila, Philippines. *Pathogens and Global Health*, **8**, 455-460. <https://doi.org/10.1179/2047773212Y.0000000061>
- Fallah MP, Skrip LA, Enders J (2018) Preventing rural to urban spread of Ebola: lessons from Liberia. *The Lancet*, **392**, 279-280. [https://doi.org/10.1016/S0140-6736\(18\)31435-1](https://doi.org/10.1016/S0140-6736(18)31435-1)
- Falzon LC, Alumasa L, Amanyia F, Kang'ethe E, Kariuki S, Momanyi K, Muinde P, Murungi MK, Njoroge SM, Ogendo A, Ogola J, Rushton J, Woolhouse MEJ, Fèvre EM (2019) One Health in action: operational aspects of an integrated surveillance system for zoonoses in Western Kenya. *Frontiers in Veterinary Science*, **6**, 252. <https://doi.org/10.3389/fvets.2019.00252>
- Firdaus G (2012) Urbanization, emerging slums and increasing health problems: a challenge before the nation: an empirical study with reference to state of Uttar Pradesh in India. *Journal of Environmental Research and Management*, **3**, 146-159.
- Firth C, Bhat M, Firth MA, Williams SH, Frye MJ, Simmonds P, Conte J, Ng J, Garcia J, Bhuvana NP, Lee B, Che X, Quan P-L, Lipkin WI (2014) Detection of zoonotic pathogens and characterization of novel viruses carried by commensal *Rattus norvegicus* in New York City. *mBio*, **5**, e1933-14. <https://doi.org/10.1128/mBio.01933-14>
- Flandroy L, Poutahidis T, Berg G, Clarke G, Dao MC, Decaestecker E, Furman E, Haahtela T, Massart S, Plovier H, Sanz Y, Rook G (2018) The impact of human activities and lifestyles on the interlinked microbiota and health of humans and of ecosystems. *Science of the Total Environment*, **627**, 1018-1038, <https://doi.org/10.1016/j.scitotenv.2018.01.288>
- Flies EJ, Mavo S, Zosky GR, Mantziorise E, Williams C, Erig R, Brooka BW, Buette JC (2019) Urban-associated diseases: candidate diseases, environmental risk factors, and a path forward. *Environment International*, **133**, e105187. <https://doi.org/10.1016/j.envint.2019.105187>
- Galimand M, Carniel E, Courvalin P (2006) Resistance of *Yersinia pestis* to antimicrobial agents. *Antimicrobial Agents and Chemotherapy*, **50**, 3233-3236. <https://doi.org/10.1128/AAC.00306-06>
- Gandy M (2021) The zoonotic city: urban political ecology and the pandemic imaginary. *International Journal of Urban Geography Research*, <https://doi.org/10.1111/1468-2427.13080>
- Garba M, Kane M, Gagaré S, Kadaouré I, Sidikou R, Rossi JP, Dobigny G (2014a) Local perception of rodent-associated problems in Sahelian urban areas: a survey in Niamey, Niger. *Urban Ecosystems*, **17**, 573-584. <https://doi.org/10.1007/s11252-013-0336-x>
- Garba M, Dalecky A, Kadaouré I, Kane M, Hima K, Véran S, Gagaré S, Gauthier P, Tatard C, Rossi JP, Dobigny G (2014b) Spatial segregation between invasive and native commensal rodents in an urban

- environment: a case study in Niamey, Niger. *PLoS ONE*, **9**, e110666. <https://doi.org/10.1371/journal.pone.0110666>
- Gibb R, Redding RW, Chin KQ, Donnelly CA, Blackburn TM, Newbold T, Jones KE (2020) Zoonotic host diversity increases in human-dominated ecosystems. *Nature*, **584**, 398-402. <https://doi.org/10.1038/s41586-020-2562-8>
- Gottdenker NL, Streicker DG, Faust CL, Carroll CR (2014) Anthropogenic land use change and infectious diseases: a review of the evidence. *EcoHealth*, **11**, 619-632. <https://doi.org/10.1007/s10393-014-0941-z>
- Grange ZL, Goldstein T, Johnson CK, Anthony S, Gilardi K, Daszak P, Olival KJ, O'Rourke T, Murray S, Olsong SH, Togami E, Vidal G, Expert Panel, PREDICT consortium, Mazet JAK (2021) Ranking the risk of animal-to-human spillover for newly discovered viruses. *Proceedings of the National Academy of Sciences, USA*, **118**, e2002324118. <https://doi.org/10.1073/pnas.2002324118>
- Grimm NB, Faeth SH, Golubiewski NE, Redman CL, Wu J, Bai X, Briggs JM (2008) Global changes and the ecology of cities. *Science*, **319**, 756-760. <https://doi.org/10.1126/science.1150195>
- Gross J, Elvinger F, Hungerford LL, Gehrt SD (2012) Raccoon use of the urban matrix in the Baltimore Metropolitan area, Maryland. *Urban Ecosystems*, **15**, 667-682. <https://doi.org/10.1007/s11252-011-0218-z>
- Guan Y, Zheng BJ, He YQ, Zhuang ZX, Cheung CL, Luo SW, Li PH, Zhang LJ, Guan YJ, Butt KM, Wong KL, Chan KW, Lim W, Shortridge KF, Yuen KY, Peiris JSM & Poon LLM (2003) Isolation and characterization of viruses related to the SARS coronavirus from animals in southern China. *Science*, **302**, 276-278. <https://doi.org/10.1126/science.1087139>
- Hancke D, Suarez OV (2014) Environmental health education in schools as strategy for rodent control: an experience in a shantytown of Buenos Aires, Argentina. *EcoHealth*, **11**, 133-140. <https://doi.org/10.1007/s10393-013-0879-6>
- Harrison NL, Sachs JD (2022) A call for an independent inquiry into the origin of the SARS-CoV-2 virus. *Proceedings of the National Academy of Sciences, USA*, **119**, e220276911. <https://doi.org/10.1073/pnas.2202769119>
- Hassel JM, Ward MJ, Muloi D, Bettridge JM, Robinson TP, Kariuki S, Ogendo A, Kiiru J, Imboma T, Kang'Ethe EK, Öghren EM, Williams NJ, Begon M, Woolhouse MEJ, Fèvre EM (2019) Clinically relevant antimicrobial resistance at the wildlife-livestock-human interface in Nairobi: an epidemiological study. *The Lancet Planetary Health*, **3**, e259-269. [https://doi.org/10.1016/S2542-5196\(19\)30083-X](https://doi.org/10.1016/S2542-5196(19)30083-X)
- Hassel JM, Bettridge JM, Ward MJ, Ogendo A, Imboma T, Muloi D, Fava F, Robinson TP, Begon M, Fèvre EM (2021) Socio-ecological drivers of vertebrate biodiversity and human-animal interfaces across an urban landscape. *Global Change Biology*, **27**, 781-792. <https://doi.org/10.1111/gcb.15412>
- Hayman DTS (2016) Bats as viral reservoirs. *Annual Review of Virology*, **3**, 77-99. <https://doi.org/10.1146/annurev-virology-110615-042203>
- Hess J, Boodram L-LG, Paz S, Ibarra AMS, Wasserheit JN, Lowe R (2020) Strengthening the global response to climate change and infectious disease threats. *Climate Change and Communicable Diseases*, **371**, m3081. <https://doi.org/10.1136/bmj.m3081>
- Heylen D, Lasters R, Adriaensen F, Fonville M, Sprong H, Matthysen E (2019) Ticks and tick-borne diseases in the city: role of landscape connectivity and green space characteristics in a metropolitan area. *Science of the Total Environment*, **670**, 941-949. <https://doi.org/10.1016/j.scitotenv.2019.03.235>
- Hima K, Houéménou G, Badou S, Garba M, Dossou H-J, Etougbétché J, Gauthier P, Artige E, Fossati-Gaschnard O, Gagaré S, Dobigny G, Dalecky A (2019) Native and invasive small mammals in urban habitats along the commercial axis connecting Benin and Niger, West Africa. *Diversity*, **11**, e238. <https://doi.org/10.3390/d11120238>
- Howard-Jones N (1975) The scientific background of the International Sanitary Conferences, 1851-1938. *History of International Health*, World Health Organization, Geneva, Switzerland.
- Huntington MK, Allison J, Nair D (2016) Emerging vector-borne diseases. *American Family Physician*, **94**, 551-557.
- Huynh TTT, Minawaka N (2022) A comparative study of dengue virus vectors in major parks and adjacent residential areas in Ho Chi Minh City, Vietnam. *PLoS Neglected Tropical Diseases*, **16**, 1, e0010119. <https://doi.org/10.1371/journal.pntd.0010119>

- Jariwala S, Redding L, Hewitt D (2017) The severely under-recognized public health risk of strongyloidiasis in North American cities: a One Health approach. *Zoonoses and Public Health*, **64**, 8, 579-588. <https://doi.org/10.1111/zph.12371>
- Jones BA, Grace D, Kock R, Alonso S, Rushton J, Said MY, McKeever D, Mutu F, Young J, McDermott J, Pfeiffer DU (2013) Zoonosis emergence linked to agricultural intensification and environmental change. *Proceedings of the National Academy of Sciences, USA*, **110**, 8399-8404. <https://doi.org/10.1073/pnas.1208059110>
- Johnson CK, Hitchens PL, Pandit PS, Rushmore J, Evans TS, Young CCW, Doyle MM (2020) Global shifts in mammalian population trends reveal key predictors of virus spillover risk. *Proceedings of the Royal Society of London B*, **287**, e20192736. <https://doi.org/10.1098/rspb.2019.2736>
- Kaiser J (2014) The catalyst. *Science*, **345**, 1112-1115. <https://doi.org/10.1126/science.345.6201.1112>
- Kamins AO, Restif O, Ntiamo-Baidu Y, Suu-Ire R, Hayman DT, Cunningham AA, Wood JLN et al. (2011) Uncovering the fruit bat bushmeat commodity chain and the true extent of fruit bat hunting in Ghana, *West African Biology and Conservation*, **144**, 3000-3008. <https://doi.org/10.1016/j.biocon.2011.09.003>
- Karan LS, Makenov MT, Korneev MG, Sacko N, Boumbaly S, Bayandin RB, Gladysheva AV, Kourouma K, Toure AH, Kartashov M, Shipovalov AV, Porshakov AM, Koulibaly M, Boiro MY (2019) Lassa Virus in the host rodent *Mastomys natalensis* within urban areas of N'zerekore, Guinea. *bioRxiv*. <https://doi.org/10.1101/616466>
- Katani R, Schilling MA, Lyimo B, Tonui T, Cattadori IM, Eblate E, Martin A, Estes AB, Buza T, Rentsch D, Davenport KW, Hovde BT, Lyimo S, Munuo L, Stomeo F, Tiambo C, Radzio-Basu J, Mosha F, Hudson PJ, Buza JJ, Kapur V (2019) Microbial diversity in bushmeat samples recovered from the Serengeti ecosystem in Tanzania. *Scientific Reports*, **9**, 18086. <https://doi.org/10.1038/s41598-019-53969-7>
- Kreuder-Johnson C, Hitchens PL, Evans TS, Goldstein T, Thomas K, Clements A, Joly DO, Wolfe ND, Daszak P, Karesh WB, Mazet JK (2015) Spillover and pandemic properties of zoonotic viruses with high host plasticity. *Scientific Reports*, **5**, e14830. <https://doi.org/10.1038/srep14830>
- Kurpiers LA, Schulte-Herbrüggen B, Ejotre I, Reeder SAM (2015) Chap. 24: bushmeat and emerging infectious diseases: lessons from Africa. In: *Problematic Wildlife*, ed. Angelici, F.M. (ed.). Springer International Publishing Switzerland. https://doi.org/10.1007/978-3-319-22246-2_24
- LaDeau SL, Allan BF, Leisnham PT, Levy MZ (2015) The ecological foundations of transmission potential and vector-borne disease in urban landscape. *Functional Ecology*, **29**, 889-901. <https://doi.org/10.1111/1365-2435.12487>
- Lau SKP, Woo PCY, Li KSM, Huang Y, Tsoi HW, Wong BHL, Wong SSY, Leung SY, Chan KH, Yuen KY (2005) Severe acute respiratory syndrome coronavirus-like virus in Chinese horseshoe bats. *Proceedings of the National Academy of Sciences, USA*, **102**, 14040-14405. <https://doi.org/10.1073/pnas.0506735102>
- Lau CL, Smythe LD, Craig SB, Weinstein P (2010) Climate change, flooding, urbanisation and leptospirosis: fuelling the fire? *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **104**, 631-638. <https://doi.org/10.1016/j.trstmh.2010.07.002>
- Lechenne M, Traoré A, Hattendorf J, Kallo V, Oussiguere A, Tetchi M, Moyengari R, Traoré AK, Moto DD, Ngandolo R, Bonfoh B, Zinsstag K (2021) Increasing rabies data availability: the example of a One Health research project in Chad, Côte d'Ivoire and Mali. *Acta Tropica*, **215**, e105808. <https://doi.org/10.1016/j.actatropica.2020.105808>
- Lembrez J (1966) Rodent control on ships and in ports of Europe, America and Middle East. *WHO Vector Control*, **66.217**, 123-128.
- Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, Ren R, Leung KSM, Lau EHY, Wong JY, Xing X, Xiang N, Wu Y, Li C, Chen Q, Li D, Liu T, Zhao J, Liu M, Tu W, Chen C, Jin L, Yang R, Wang Q, Zhou S, Wang R, Liu H, Luo Y, Liu Y, Shao G, Li H, Tao Z, Yang Y, Deng Z, Liu B, Ma Z, Zhang Y, Shi G, Lam TTY, Wu JT, Gao GF, Cowling BJ, Yang B, Leung GM, Feng Z (2020) Early transmission dynamics in Wuhan, China, of a novel coronavirus-infect pneumonia. *New England Journal of Medicine*, **382**, 1199-1207. <https://doi.org/10.1056/NEJMoa2001316>
- Lin XD, Guo WP, Wang W, Zou Y, Hao ZY, Zhou DJ, Jong X, Qu YG, Li MH, Tian HF, Wen JF, Plyusnin A, Xu J, Zhang YZ (2012) Migration of Norway rats resulted in the worldwide distribution of Seoul hantavirus today. *Journal of Virology*, **88**, 972-981. <https://doi.org/10.1128/JVI.00725-11>

- Lloyd-Smith JO, Cross PC, Briggs CJ, Gaugherty M, Getz WM, Latto J, Sanchez MS, Smith AB, Swei A (2005) Should we expect population threshold for wildlife disease? *Trends in Ecology and Evolution*, **20**, 511-519. <https://doi.org/10.1016/j.tree.2005.07.004>
- Lydecker W, Hochuli DF, Banks PB (2019) Peri-urban black rats host a rich assembly of ticks and healthier rats have more ticks. *Ticks and Tick-Borne Diseases*, **10**, 749-753. <https://doi.org/10.1016/j.ttbdis.2019.03.007>
- Marí Saéz A, Weiss S, Nowak K, Lapeyre V, Zimmermann F, Dux A, Kühl HS, Kaba M, Regnaut S, Merkel K, Sachse A, Thiesen U, Villányi L, Boesch C, Dabrowski PW, Radonić A, Nitsche A, Leendertz SA, Petterson S, Becker S, Krähling V, Couacy-Hymann E, Akoua-Koffi C, Weber N, Schaade L, Fahr J, Borchert M, Gogarten JF, Calvignac-Spencer S, Leendertz FH (2015) Investigating the zoonotic origin of the West African Ebola epidemic. *EMBO Molecular Medicine*, **7**, 17-23. <https://doi.org/10.15252/emmm.201404792>
- McNamara J, Fa JE, Ntiamoa-Baidu Y (2019) Understanding drivers of urban bushmeat demand in a Ghanaian market. *Biological Conservation*, **239**, e108291. <https://doi.org/10.1016/j.biocon.2019.108291>
- Meerburg B, Singleton GR, Kijlstra H (2009) Rodent-borne diseases and their risks for public health. *Critical Reviews in Microbiology*, **35**, 221-270. <https://doi.org/10.1080/10408410902989837>
- Mereiles L, Galisteo Jr A, Pompeu E, Andrade Jr H (2004) *Toxoplasma gondii* spreading in an urban area evaluated by sero-prevalence in free-living cats and dogs. *Tropical Medicine and International Health*, **9**, 8, 876-881. <https://doi.org/10.1111/j.1365-3156.2004.01280.x>
- Mills GS, Brookes JD, Gellie NJC, Liddicoat C, Lowe AJ, Sydnor HR, Thomas T, Weinstein P, Weyrich LS, Breed MF (2019) Relating urban biodiversity to human health with the 'holobiont' concept. *Frontiers in Microbiology*, **10**, e550. <https://doi.org/10.3389/fmicb.2019.00550>
- Mironova V, Shartova N, Beljaev A, Varentsov M, Grishchenko M (2019) Effects of climate change and heterogeneity of local climates on the development of malaria parasite (*Plasmodium vivax*) in Moscow megacity region. *International Journal of Environmental Research and Public Health*, **16**, 694. <https://doi.org/10.3390/ijerph16050694>
- Misslin R, Telle O, Daudé E, Vaguet A, Paul RE (2016) Urban climate versus global climate change - what makes the difference for dengue? *Annals of the New York Academy of Science*, **1382**, 56-72. <https://doi.org/10.1111/nyas.13084>
- Mohd Zain S, Behnke J, Lewis J (2012) Helminth communities from two urban rat populations in Kuala Lumpur, Malaysia. *Parasites and Vectors*, **5**, e47. <https://doi.org/10.1186/1756-3305-5-47>
- Moller AP, Diaz M, Flensted-Jensen E, Grim T, Ibanez-Alamo JD, Jokimäki J, Mänd R, Marko G, Tryjanowski P (2012) High urban population density of birds reflects their timing of urbanization. *Oecologia*, **170**, 867-875. <https://doi.org/10.1007/s00442-012-2355-3>
- Moore P (2020) *Le petit livre des grandes épidémies*. Belin eds., 144 pages. ISBN: 9782701193649.
- Morand S, Blasdell K, Bordes F, Buchy F, Carcy B, Chaisiri K, Chaval Y, Claude J, Cosson J-F, Desquesnes M, Jittapalpong S, Jiyipong T, Karnchanabanthoen A, Pornpan P, Rolain J-M, Tran A-L (2019) Changing landscapes of Southeast Asia and rodent-borne diseases: decreased diversity but increased transmission risks. *Ecology Applications*, **29**, e01886. <https://doi.org/10.1002/eap.1886>
- Muinde P, Bettridge JM, Sousa FM, Dürr S, Dohoo IR, Berezowski J, Mutwiri T, Odinga CO, Fèvre EM, Falzon LC (2021) Who let the dog out? Exploring the spatial ecology of free-roaming dogs in western Kenya. *Ecology and Evolution*. <https://doi.org/10.1002/ece3.7317>
- Muloi DM, Wee BA, McClean DMH, Ward MJ, Pankhurst L, Phan H, Ivens AC, Kivali V, Kiyong'a A, Ndinda C, Gitahi N, Ouko T, Hassell JM, Imboma T, Akoko J, Murungi MK, Njoroge SM, Muinde P, Nakamura Y, Alumasa L, Furmaga E, Kaitho T, Ohgren EM, Amnaya F, Ogendo A, Wilson D, Bettridge JM, Kiiru J, Kyobutungi C, Tacoli C, Kang'ethe EK, Davila JD, Kariuki S, Robinson TP, Rushton J, Woolhouse MEJ, Fèvre M (2022) Population genomics of *Escherichia coli* in livestock-keeping households across a rapidly developing urban landscape. *Nature Microbiology*, **7**, 581-589. <https://doi.org/10.1038/s41564-022-01079-y>
- Murray MH, Fyffe R, Fidino M, Byers KA, Rios MJ, Mulligan MP, Magle SB (2018) Public complaints reflect rat relative abundance across diverse urban neighborhoods. *Frontiers in Ecology and Evolution*, **6**, e189. <https://doi.org/10.3389/fevo.2018.00189>

- Musso D, Gubler DJ (2016) Zika Virus. *Clinical Microbiology Reviews*, **29**, 487-524. <https://doi.org/10.1128/CMR.00072-15>
- Nickel S, von dem Knesebeck O (2020) Do multiple community-based interventions on health promotion tackle health inequalities? *International Journal of Equity in Health*, **19**, e157. <https://doi.org/10.1186/s12939-020-01271-8>
- Nieto-Rabiela F, Wiradtsudakul A, Suzan G, Rico-Chavez O (2019) Viral networks and detection of potential zoonotic viruses in bats and rodents: a worldwild analysis. *Zoonoses and Public Health*, **66**, 655-666. <https://doi.org/10.1111/zph.12618>
- Neiderud CJ (2015) How urbanization affects the epidemiology of emerging infectious diseases. *Infections Ecology and Epidemiology*, **5**, e27060. <https://doi.org/10.3402/iee.v5.27060>
- Newsome SD, Garbe HM, Wilson EC, Gehrt SD (2015) Individual variation in anthropogenic resource use in an urban carnivore. *Oecologia*, **178**, 115-128. <https://doi.org/10.1007/s00442-014-3205-2>
- Nickel S, von dem Knesebeck O (2020) Effectiveness of community-based health promotion interventions in urban areas: a systematic review. *Journal of Community Health*, **45**, 419-434. <https://doi.org/10.1007/s10900-019-00733-7>
- Normile D (2004) Infectious diseases. Mounting lab accidents raise SARS fears. *Science*, **304**, 659-661. <https://doi.org/10.1126/science.304.5671.659>
- Nunes H, Lopes Rocha F, Cordeiro-Estrela P (2017) Bats in urban areas of Brazil: roosts, food resources and parasites in disturbed environments. *Urban Ecosystems*, **20**, 953-969. <https://doi.org/10.1007/s11252-016-0632-3>
- Olival KJ, Hosseini PR, Zambrana-Torrel C, Ross R, Bogich TL, Daszak P (2017) Host and viral traits predict zoonotic spillover from mammals. *Nature*, **546**, 612-646. <https://doi.org/10.1038/nature22975>
- Ouédraogo S, Benmarhnia T, Bonnet E, Somé P-A, Barro AS, Kafando Y, Soma DD, Dabiré RK, Saré D, Fournet F, Ridde V (2018). Evaluation of effectiveness of a community-based intervention for control of dengue virus vector, Ouagadougou, Burkina Faso. *Emerging Infectious Diseases*, **24**, 1859-1867. <https://doi.org/10.3201/eid2410.180069>
- Owens B (2014) Anthrax and smallpox errors highlight gaps in US biosafety. *The Lancet*, **384**, 294. [https://doi.org/10.1016/S0140-6736\(14\)61246-0](https://doi.org/10.1016/S0140-6736(14)61246-0)
- Parente CC, Bezerra FSM, Parente PI, Dias-Neto RV, Xavier SCC, Ramos Jr AN, Carvalho-Costa FA, Lima MM (2017) Community-based entomological surveillance reveals urban foci of Chagas disease vectors in Sobral, State of Ceará, Northeastern Brazil. *PLoS One*, **12**, e0170278. <https://doi.org/10.1371/journal.pone.0170278>
- Park SW, Ha NY, Ryu B, Bang JH, Song H, Kim Y, Kim G, Oh M, Cho NH, Lee JK (2015) Urbanization of scrub typhus disease in South Korea. *PLoS Neglected Tropical Diseases*, **9**, e3814. <https://doi.org/10.1371/journal.pntd.0003814>
- Parra-Henao G, Garzon-Jimenez SP, Bernal-Rosas Y, Olivera MJ, Salgado M, Torres-Garcia OA (2021) Risk factors for triatominae infestation in a municipality of Colombia. *Therapeutic Advances in Infectious Diseases*, **8**, e20499361211030068. <https://doi.org/10.1177/20499361211030068>
- Parsons MH, Banks PB, Deutsch MA, Corrigan RF, Munshi-South J (2017) Trends in urban rat ecology: a framework to define the prevailing knowledge gaps and incentives for academia, pest management professionals (PMPs) and public health agencies to participate. *Journal of Urban Ecology*, **3**, 1-8. <https://doi.org/10.1093/jue/jux005>
- Parsons MH, Jardine CM, Crowther MS, Himsforth CS (2020) Trends in urban rodent monitoring and mitigation: improving our understanding of population and disease ecology, surveillance and control. *Frontiers in Ecology and Evolution*, **7**, e522. <https://doi.org/10.3389/fevo.2019.00522>
- Partridge SR, Kwong SM, Firth N, Jensen SO (2018) Mobile genetic elements associated with antimicrobial resistance. *Clinical Microbiology Reviews*, **31**, e00088-17. <https://doi.org/10.1128/CMR.00088-17>
- Patz JA, Daszak P, Tabor GM, Aguirre AA, Pearl M, Epstein J, Wolfe ND, Kilpatrick AM, Foutopoulos J, Molyneux D, Bradley DJ and members of the Working Group on Land Use Change and Disease Emergence (2004) Unhealthy landscapes: policy recommendations on land use change and infectious disease emergence. *Environmental and Health Perspectives*, **112**, 1092-1098. <https://doi.org/10.1289/ehp.6877>
- Peel AJ, Sargan DR, Baker KS, Hayman DTS, Barr JA, Crameri G, Suu-Ire R, Broder CC, Lembo T, Wang LF, Fooks AR, Rossiter SJ, Wood JLN, Cunningham AA (2013) Continent-wide panmixia of an African fruit

- bat facilitates transmission of potentially zoonotic viruses. *Nature Communications*, 4, e2770. <https://doi.org/10.1038/ncomms3770>
- Peel AJ, Wood JLN, Baker K, Breed AC, De Carvalhos A, Fernandez-Loras A, Gabrieli HS, Gembu GC, Kakengi VA, Kaliba PM, Kityo RM, Lembo Y, Mba FE, Ramos D, Rodriguez-Prieto I, Suu-Ire R, Cunningham AA, Hayman DTS (2017) How does Africa's most hunted bat vary across the continent? Population traits of the straw-coloured fruit bat (*Eidolon helvum*) and its interactions with humans. *Acta Chiropterologica*, 19, 77-92. <https://doi.org/10.3161/15081109ACC2017.19.1.006>
- Peng H, Bilal M, Iqbal HMN (2018) Improved biosafety and biosecurity measures and/or strategies to tackle laboratory-acquired infections and related risks. *International Journal of Environmental Research and Public Health*, 15, 2697. <https://doi.org/10.3390/ijerph15122697>
- Pickard JM, Zeng MY, Caruso R, Núñez G (2017) Gut microbiota: role in pathogen colonization, immune responses, and inflammatory disease. *Immunological Reviews*, 279, 70-89. <https://doi.org/10.1111/imr.12567>
- Plowright RK, Foley P, Field HE, Dobson AP, Foley JE, Eby P, Daszak P (2011) Urban habituation, ecological connectivity and epidemic dampening: the emergence of Hendra virus from flying foxes (*Pteropus* spp.). *Proceedings of the Royal Society of London B*, 278, 3703-3712. <https://doi.org/10.1098/rspb.2011.0522>
- Prescott J, Bushmaker T, Fischer R, Miazgowicz K, Judson S, Munster VJ (2015) Postmortum stability of Ebola virus. *Emerging Infectious Diseases*, 21, 856-859. <https://doi.org/10.3201/eid2105.150041>
- Promkerd P, Khoprasert Y, Virathavone P, Thoumabout M, Sirisak O, Jäkel T (2008) Factors explaining the abundance of rodents in the city of Luang Prabang, Lao PDR, as revealed by field and household surveys. *Integrative Zoology*, 3, 11-20. <https://doi.org/10.1111/j.1749-4877.2008.00069.x>
- Randremanana R, Andrianaivoarimanana V, Nikolay B, Ramasindrazana B, Paireau J, Ten Bosch QA, Rakotondramanga JM, Rahajandraibe S, Rahelinirina S, Rakotomanana F, Rakotoarimanana FM, Randriamampionona LB, Razafimbiana V, De Dieu Randria MJ, Raberahona M, Mikaty G, Le Guern AS, Rakotonjanabelo LA, Ndiaye CF, Rasolofo V, Bertherat E, Ratsitorahina M, Cauchemez S, Baril L, Spiegel A, Rajerison M (2019) Epidemiological characteristics of an urban plague epidemic in Madagascar, August-November, 2017: an outbreak report. *Lancet Infectious Diseases*, 19, 537-545. [https://doi.org/10.1016/S1473-3099\(18\)30730-8](https://doi.org/10.1016/S1473-3099(18)30730-8)
- Reed JH, Bonter DN (2018) Supplementing non-target taxa: bird feeding alters the local distribution of mammals. *Ecological Applications*, 28, 761-770. <https://doi.org/10.1002/eap.1683>
- Reye AL, Hübschen JM, Sausy A, Muller CP (2010) Prevalence and seasonality of tick-borne pathogens in questing Ixodes ricinus ticks from Luxembourg. *Applied Environmental Microbiology*, 76, 2923-31. <https://doi.org/10.1128/AEM.03061-09>
- Richard Y, Pohl B, Rega M, Pergaud J, Thevenin T, Emery J, Dudek J, Vairet T, Zito S, Chateau-Smith C, (2021) Is Urban Heat Island intensity higher during hot spells and heat waves (Dijon, France, 2014-2019)? *Urban Climate*, 35, e100747. <https://doi.org/10.1016/j.uclim.2020.100747>
- Ridde V, Pérez D, Robert E (2020) Using implementation science theories and frameworks in global health. *BMJ Global Health*, 5, e002269. <https://doi.org/10.1136/bmjgh-2019-002269>
- Ridde V, Rossier C, Soura AB, Bazié F, Kadio K (2014) A community-based approach to indigent selection is difficult to organize in a formal neighbourhood in Ouagadougou, Burkina Faso: a mixed methods exploratory study. *International Journal of Equity in Health*, 13, e31. <https://doi.org/10.1186/1475-9276-13-31>
- Robin M-M (2021) *La fabrique des pandémies : préserver la biodiversité, un impératif pour la santé planétaire*. Cahiers Libres, Eds. La Découverte, Paris, France. 327 pages.
- Roche B, Garchitorena A, Guéegan J-F, Arnal A, Roiz D, Morand S, Zambrana-Torrel C, Suzan G, Daszak P (2020) Was the COVID-19 pandemic avoidable? A call for a "solution-oriented" approach in pathogen evolutionary ecology to prevent future outbreaks. *Ecology Letters*, 23, 1557-1560. <https://doi.org/10.1111/ele.13586>
- Rungroj A, Chaisiri K, Paladsing Y, Morand S, Junjhon J, Blacksell SD, Ekcharyawat P (2021) Prevalence and molecular characterization of Rickettsia spp. from wild small mammals in public parks and urban areas of Bangkok Metropolitan, Thailand. *Tropical Medicine and Infectious Diseases*, 6, e199. <https://doi.org/10.3390/tropicalmed6040199>
- Rivkin LR, Santangelo JS, Johnson MTJ (2018) A roadmap for urban evolutionary ecology. *Evolutionary Applications*, 12, 384-398. <https://doi.org/10.1111/eva.12734>

- Santiago-Alarcon D, MacGregor-Fors I (2020) Cities and pandemics: urban areas are ground zero for the transmission of emerging human infectious diseases. *Journal of Urban Ecology*, **6**, <https://doi.org/10.1093/jue/juaa012>
- Santini L, Gonzalez-Suarez M, Russo D, Gonzalez-Voyer A, von Hardenberg A, Ancillotto L (2019) One strategy does not fit all: determinants of urban adaptation in mammals. *Ecology Letters*, **22**, 365-376. <https://doi.org/10.1111/ele.13199>
- Santos NJ, Carneiro IO, Borges Carneiro AJ, Santos da Paz E, Brito RL, Franke CR (2019) Eco-epidemiological approach in an urban community: environmental factors associated with injuries in humans caused by bats. *Journal of Infectious Developing Countries*, **13**, 1045-1051. <https://doi.org/10.3855/jidc.11904>
- Saré D, Pérez D, Somé P-A., Kafando Y, Barro A, Ridde V (2018) Community-based dengue control intervention in Ouagadougou: intervention theory and implementation fidelity. *Global Health Research*, **Pol. 3**, e21. <https://doi.org/10.1186/s41256-018-0078-7>
- Saylor KE, Mouichec MM, Lucas A, Mclvera DJ, Matsidae A, Clarya C, Maptue VT, Euren JD, LeBretonc M, Tamoufe U (2021) Market characteristics and zoonotic disease risk perception in Cameroon bushmeat markets. *Social Science and Medicine*, **268**, e113358. <https://doi.org/10.1016/j.socscimed.2020.113358>
- Semenza JC, Suk JE (2018) Vector-borne diseases and climate change: a European perspective. *FEMS Microbiology Letters*, **365**, 244. <https://doi.org/10.1093/femsle/fnx244>
- Shairp R, Verissimo D, Fraser I, Challender D, MacMillan D (2016) Understanding urban demand for wild meat in Vietnam: implications for conservation actions. *PLoS One*, **11**, e0134787. <https://doi.org/10.1371/journal.pone.0134787>
- Shuai H, Chan JF-W, Yuen TT-T, Yoon C, Hu J-C, Hu B, Yang D, Wang Y, Hou Y, Huang X, Chai Y, Chan CC-S, Poon VK-M, Lu L, Zhang R-Q, Chan W-M, Ip JD, Chu AW-C, Hu Y-F, Cai J-P, Chan K-H, Zhou J, To KK-W, Yuen K-Y, Chu H (2021) Emerging SARS-CoV-2 variants expand species tropism to murines. *eBio Medecine*, **73**, e103643. <https://doi.org/10.1016/j.ebiom.2021.103643>
- Soulsbury C, Piran C, White CL (2015) Human-wildlife interactions in urban areas: a review of conflicts, benefits and opportunities. *Wildlife Research*, **42**, 541-553. <https://doi.org/10.1071/WR14229>
- Stenseth NC, Atshabar BB, Begon M, Belmain SR, Bertherat E, Carniel E, Gage KL, Leirs H, Rahalison L (2008) Plague: past, present, and future. *PLoS Medicine*, **5**, e3. <https://doi.org/10.1371/journal.pmed.0050003>
- Sun L, Chen J, Li Q (2020) Dramatic uneven urbanization of large cities throughout the world in recent decades. *Nature Communications*, **11**, e5366. <https://doi.org/10.1038/s41467-020-19158-1>
- Suu-Ire RD, Fooks AR, Banyard AC, Selden D, Amposah-Mensah K, Riesle S, Ziekah MY, Ntiama-Biadiu Y, Wood JLN, Cunningham AA (2017) Lagos Bat Virus infection dynamics in free-ranging straw-colored fruit bats (*Eidolon helvum*). *Tropical Medicine Infectious Diseases*, **8**, e25, <https://doi.org/10.3390/tropicalmed2030025>
- Swei A, Couper LI, Coffrey LL, Kapan D, Bennett S (2020) Patterns, drivers and challenges of vector-borne disease emergence. *Vector-Borne and Zoonotic Diseases*, **20**, 159-170. <https://doi.org/10.1089/vbz.2018.2432>
- Takken W, Lindsay S (2019) Increased threat of urban malaria from *Anopheles stephensi* mosquitoes, Africa. *Emerging Infectious Diseases*, **25**, 1431-1433. <https://doi.org/10.3201/eid2507.190301>
- Tait J, Perotto-Baldivieso HL, McKeown A, Westcott DA (2014) Are flying-foxes coming to town? Urbanisation of the spectacled flying-fox (*Pteropus conspicillatus*) in Australia. *PLoS One*, **9**, e109810. <https://doi.org/10.1371/journal.pone.0109810>
- Tenzin T, Ahmed R, Debnath RC, Garba A, Mat Y (2015) Free-roaming dog population estimation and status of the dog population management and rabies control program in Dhaka city, Bangladesh. *PLoS Neglected Tropical Diseases*, **9**, e0003784. <https://doi.org/10.1371/journal.pntd.0003784>
- Toledo Romani ME, Vanlerberghe V, Perez D, Lefevre P, Ceballos E, Bandera D, Gil AB, Van der Stuyft P (2007) Achieving sustainability of community-based dengue control in Santiago de Cuba. *Social Science and Medicine*, **64**, 976-988. <https://doi.org/10.1016/j.socscimed.2006.10.033>
- Tong S, Li Y, Rivaille P, Conrardy C, Alvarez DA, Chen LM, Recueno S, Ellison JA, Davis CT, Yors IA, Turmelle AS, Moran D, Rogers S, Shi M, Tao Y, Weil MR, Tang K, Rowe LA, Sammons S, Xu X, Frace M, Lindblade KA, Cox NJ, Anderson LJ, Rupprecht CE, Donis RO (2012) A distinct lineage of influenza virus A from bats. *Proceedings of the National Academy of Sciences USA*, **109**, 4269-4274. <https://doi.org/10.1073/pnas.1116200109>

- Tzaninis Y, Mandler T, Kaika M, Keil R (2021) Moving urban political ecology beyond the 'urbanization of nature'. *Progress in Human Geography*, **45**, 229-252. <https://doi.org/10.1177/0309132520903350>
- UN Habitat (2019) *World urbanization prospects: the 2018 revision*. United Nations Habitat, New York, USA. ST/ESA/SER.A/420.
- UN Habitat (2021) *Cities and pandemics: towards a more just, green and healthy future*. United Nations Human Settlements Program, UN Habitat, Nairobi, Kenya.
- De Urioste-Stone SM, Pennington PM, Pellecer E, Aguilar TM, Samayoa G, Perdomo HD, Enriquez H, Juarez JG (2015) Development of a community-based intervention for the control of Chagas disease based on peridomestic animal management: an eco-bio-social perspective. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **109**, 159-167. <https://doi.org/10.1093/trstmh/tru202>
- Valdez D, Keys H, Ureña K, Cabral D, Camilo F, Ogando EC, Mercedes L, Noland GS, Blount SB, Lavery JV, Desir L, Puello J (2020) Malaria outbreak response in urban Santo Domingo, Dominican Republic: lessons learned for community engagement. *Revista Panamericana de Salud Publica*, **44**, e92. <https://doi.org/10.26633/RPSP.2020.92>
- Verrelli BC, Alberti M, Des Roches S, Harris NC, Hendry AP, Johnson MTJ, Savage AM, Charmantier A, Gotanda KM, Govaert L, Miles S, Rivkin LR, Winchell KM, Brans KI, Correa C, Diamond S, Fitzhugh B, Grimm NB, Hugues S, Marzluff JM, Munshi-South J, Rojas C, Santangelo JS, Schell CJ, Schweitzer JA, Szulkin M, Urban MC, Zhou Y, Ziter C (2022) A global horizon scan for urban evolutionary ecology. *Trends in Ecology and Evolution*, **3038**, <https://doi.org/10.1016/j.tree.2022.07.012>
- Vijaykrishna D, Mukerji R, Smith GJD (2015) RNA virus reassortment: an evolutionary mechanism for host jumps and immune evasion. *PLoS Pathogens*, **11**, e1004902. <https://doi.org/10.1371/journal.ppat.1004902>
- Webb Hooper M, Mitchell C, Marshall VJ, Cheatham C, Austin K, Sanders K, Krishnamurthi S, Grafton LL (2019) Understanding multilevel factors related to urban community trust in healthcare and research. *International Journal of Environmental Research and Public Health*, **16**, e3280. <https://doi.org/10.3390/ijerph16183280>
- Wei C, Shan K-J, Wang W, Zhang S, Huan Q, Qian W (2021) Evidence for a mouse origin of the SARS-CoV-2 Omicron variant. *Journal of Genetics and Genomics*, **998**, <https://doi.org/10.1016/j.jgg.2021.12.003>
- Wells K, Morand S, Wardeh M, Baylis M (2019) Distinct spread of DNA and RNA viruses among mammals amid prominent role of domestic species. *Global Ecology and Geography*, **29**, 470-481. <https://doi.org/10.1111/geb.13045>
- Werner CS, Nunn CL (2020) Effect of urban habitat use on parasitism in mammals: a meta-analysis. *Proceedings of the Royal Society of London B*, **287**, e20200397. <https://doi.org/10.1098/rspb.2020.0397>
- Westcott DA, Mc Kneown A, Hazel P, Parsons J, Jurdak R, Kusy B, Sommer P, Zhao K, Dobbie M, Heersink D, Camey P (2015) *Implementation of the national flying-fox monitoring program*. Australian Government, Rural Industries Research and Development Corporation, Publication No. 15/101, Project No. 008241, 190 pages.
- White RJ, Razgour O (2020) Emerging zoonotic diseases originating in mammals: a systematic review of effects of anthropogenic land-use change. *Mammal Review*. <https://doi.org/10.1111/mam.12201>
- WHO (2005) *International Health Regulation*. World Health Organization, Geneva, Switzerland. <https://www.who.int/publications/i/item/9789241580410>
- WHO, FAO and OIE (2019) *Taking a multisectoral, One Health approach: a tripartite guide to addressing zoonotic diseases in countries*. Joint Risk Assessment Operational Tool: an operational tool of the Tripartite Zoonoses Guide. <https://apps.who.int/iris/handle/10665/325620>
- Wilke ABB, Vasquez C, Carvajal A, Medina J, Chase C, Cardenas G, Mutebi JP, Petrie WD, Beier JC (2020) Proliferation of *Aedes aegypti* in urban environments mediated by the availability of key aquatic habitats. *Scientific Reports*, **10**, e12925. <https://doi.org/10.1038/s41598-020-69759-5>
- Wilkinson A, Leach M (2015) Briefing: Ebola- myths, realities and structural violence. *African Affairs*, **114**, 136-148. <https://doi.org/10.1093/afraf/adu080>
- Wilson RT (2018) Domestic livestock in African cities: production, problems and prospects. *Open Urban Study in Demography Journal*, **4**, 1-14. <https://doi.org/10.2174/2352631901804010001>
- Wolfe ND, Daszak P, Kilpatrick AM, Burke DS (2005) Bushmeat hunting, deforestation, and prediction of zoonotic disease emergence. *Emerging Infectious Diseases*, **11**, 1822-1827. <https://doi.org/10.3201/eid1112.040789>

- Wulandary SA, Paladsing Y, Saesim W, Charoennitiwat V, Sonthayanon P, Kumlert R, Morand S, Sumruayphol S, Chaisiri K (2021) High prevalence and low diversity of chigger infestation in small mammals found in Bangkok Metropolitan parks. *Medical and Veterinary Entomology*, **35**, 4, 534-546. <https://doi.org/10.1111/mve.12531>
- Yin C, Sun P, Yu X, Wang P, Cheng G (2020) Roles of symbiotic microorganisms in arboviral infection of arthropod vectors. *Trends in Parasitology*, **36**, 607-615. <https://doi.org/10.1016/j.pt.2020.04.009>
- Zietz BP, Dunkelberg H (2004) The history of the plague and the research on the causative agent *Yersinia pestis*. *International Journal of Environmental Health and Hygiene*, **207**, 165-178. <https://doi.org/10.1078/1438-4639-00259>